

16. 04. 2004



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

EP 04 / 4076

REC'D 28 JUN 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03019626.5

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk



Anmeldung Nr:  
Application no.: 03019626.5  
Demande no:

Anmeldetag:  
Date of filing: 04.09.03  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

NeuroNova AG  
Kraepelinstrasse 2  
80804 München  
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se referer à la description.)

Means and methods for diagnosing and treating affective disorders

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT RO SE SI SK TR LI

### **Means and methods for diagnosing and treating affective disorders**

The present invention relates to nucleic acid molecules, preferably genomic sequences, encoding an ATP-gated ion channel P2X7R which contain a mutation in the 5'UTR or 3'UTR regions, a mutation in exon 3, 5, 6, 8 or 13 or in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 or a deletion in exon 13, which allow to diagnose affective disorders. The invention further relates to polypeptides encoded by said nucleic acid molecules vectors and host cells comprising said nucleic acid molecules as well as to methods for producing polypeptides encoded by said nucleic acid molecules. The present invention also provides antibodies specifically directed to polypeptides encoded by said nucleic acid molecules and aptamers specifically binding said nucleic acid molecules.

Additionally, primers for selectively amplifying said nucleic acid molecules are provided in the present invention as well as kits, compositions, particularly pharmaceutical and diagnostic compositions comprising said nucleic acid molecules, vectors, polypeptides, aptamers, antibodies and/or primers. Moreover, the present invention relates to methods for diagnosing affective disorders associated with a non-functional P2X7R protein, an altered ATP-gating of the P2X7R protein, an over- or underexpression of the P2X7R protein or associated with the presence of any one of the aforementioned nucleic acid molecules or polypeptides encoded thereby. Additionally, the present invention relates to uses and methods for treating affective disorders employing a functional or non-functional ATP-gated ion-channel P2X7R.

The present invention also relates to uses of modulators of P2X7R activity for treating affective diseases.

Furthermore, the present invention also relates to methods for identifying and characterizing compounds which are capable of specifically interacting with or altering the characteristics of the polypeptides of the present invention as well as to methods for the production of pharmaceutical compositions.

Up to 10% of persons visiting a physician are afflicted with an affective disorder (also known as behavioural disorder, mood disorder). Nonetheless, most cases remain

undiagnosed or inadequately treated. Affective disorders include among others, depression, anxiety, and bipolar disorder. These diseases are well described in the literature; see, for example, Diagnostic and Statistical Manual of Mental Disorders-4th Edition Text Revision (DMS-IV-TR), American Psychiatric Press, 2000.

Depression, also known as unipolar affective disorder, is characterized by a combination of symptoms such as lowered mood, loss of energy, loss of interest, feeling of physical illness, poor concentration, altered appetite, altered sleep and a slowing down of physical and mental functions resulting in a relentless feeling of hopelessness, helplessness, guilt, and anxiety. The primary subtypes of this disease are major depression, dysthymia (milder depression), and atypical depression. Other important forms of depression are premenstrual dysphoric disorder and seasonal affective disorder. Present treatment of depression consists of psychotherapy, antidepressant drugs, or a combination of both. Most antidepressive drugs target the transport of the neurotransmitters serotonin and/or norepinephrine, or the activity of the enzyme monoamine oxidase. They include: Selective serotonin-reuptake inhibitors (e.g., fluoxetine, paroxetine, sertraline, fluvoxamine), tricyclic antidepressants (e.g., amitriptyline, imipramine, desipramine, nortriptyline), monoamine oxidase inhibitors (e.g., phenelzine, isocarboxazid, tranylcypromine), and designer antidepressants such as mirtazapine, reboxetine, nefazodone. However, all existing antidepressive drugs possess shortcomings such as long latency until response, high degree of non-responders and undesirable side effects (Holsboer, Biol. Psychol. 57 (2001), 47-65). Therefore, a need exists in the medical community for new antidepressive drugs with improved pharmacological profile (Baldwin, Hum. Psychopharmacol. Clin. Exp. 16 (2001), S93-S99).

Anxiety disorders are defined by an excessive or inappropriate aroused state characterized by feelings of apprehension, uncertainty, or fear. They are classified according to the severity and duration of their symptoms and specific affective characteristics. Categories include: (1) Generalized anxiety disorder, (2) panic disorder, (3) phobias, (4) obsessive-compulsive disorder, (5) post-traumatic stress disorder, and (6) separation anxiety disorder. The standard treatment for most anxiety disorders is a combination of cognitive-behavioural therapy with antidepressant medication. Additional medications include benzodiazepines and buspirone.



Bipolar disorder, also known as manic-depression, is characterized by mood swings between periods of mania (i.e. mood elevation including exaggerated euphoria, irritability) and periods of depression. Bipolar disorder is classified according to the severity of the symptoms. Patients diagnosed with bipolar disorder type I suffer from manic or mixed episodes with or without major depression. In Bipolar Disorder type II, patients have episodes of hypomania and episodes of major depression. With hypomania the symptoms of mania (euphoria or irritability) appear in milder forms and are of shorter duration. The current drugs used to treat bipolar disorders are lithium, valproate and lamotrigine, which stimulates the release of the neurotransmitter glutamate. As with antidepressive drugs, they take weeks to become effective and can result in undesirable side effects, for example, high levels of lithium in the blood can be fatal.

Compelling evidence suggest that affective disorders are biological diseases. However, there are no laboratory tests or other procedures that a common physician can use to make a definitive diagnosis. Instead, a specially trained physician or psychiatrist must diagnose the illness based on a group of symptoms that occur together. This process is often time consuming and laborious requiring several visits for the physician to perform a careful history of the symptoms that the patient is currently experiencing as well as any symptoms he or she has had in the past. Therefore, an easy and effective method for the accurate diagnosis of affective disorders is of high interest to the medical community (Wittchen et al., J. Clin. Psychiatry 62, suppl. 26 (2001), 23-28).

Most patients afflicted with affective disorders have family antecedents and identical twins studies suggest a strong genetic component. For example, genetic mapping on an isolated population of the central valley of Costa Rica suggests a locus for severe bipolar disorder at chromosome 18q22-q23 (Freimer et al., Nature Genetics 12 (1996), 436-441). Moreover, genetic studies performed on the Old Order Amish population suggest that genes on chromosomes 6, 13, and 15 may contribute to the susceptibility of bipolar affective disorder (Ginns et al., Nature Genetics 12 (1996), 431-435). Recently, a genome-wide search in a homogenous population found in the Saguenay/Lac-St-Jean region of Quebec suggests the presence of a major locus for bipolar disorder on chromosome 12q23-q24 (Morissette et al., Am. J. Med. Genet. (Neuropsychiatr. Genet.) 88 (1999), 567-587). Susceptibility loci on chromosomes 5

and 21 were also found in this study. Other groups report minimal evidence for linkage in the region of 12q23 (Kelsoe et al., Proc. Natl. Acad. Sci. USA 98 (2001), 585-590; Sklar, Annu. Rev. Genomics Hum. Genet. 3 (2002), 371-413). Given the various loci mentioned in the above studies (e.g., links to chromosomes 5, 6, 12, 13, 15, 18, 21), a definite genetic link for affective diseases remains to be found.

Thus, although several genes have been assumed to be linked with affective disorders as mentioned hereinabove, however, no clear correlation has so far been shown. Since no well-suited medication nor diagnosis on a molecular level for affective disorders is available, there is a need for identifying a gene whose mutations cause the whole spectrum of affective disorders as well as for providing medicaments and methods for diagnosis and treatment of affective disorders.

Thus, the technical problem underlying the present invention is to provide means and methods for diagnosis and treating affective disorders.

The solution to said technical problem is achieved by providing the embodiments characterized in the claims.

Accordingly, the present invention relates to a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:

- (a) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 5'UTR region corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;
- (b) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of the ATP-gated ion channel P2X7R, wherein in the exon as indicated in column "Exon" of the following Table A the amino acid residue as indicated in column "Amino acid residue" of Table A corresponding to the position as indicated in column "Position in wild-type" of Table A of the wild-type ATP-gated ion channel P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue

**Table A**

<b>Exon</b>	<b>Amino acid residue</b>	<b>Position in wild-type</b>
exon 3	R (Arg)	117
exon 5	G (Gly)	150
exon 6	E (Glu)	186
exon 6	L (Leu)	191
exon 8	R (Arg)	270
exon 13	I (Ile)	568
exon 13	R (Arg)	578

- (c) a nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in exon 5 or 8 corresponding to position 32548 or position 37633 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide
- (d) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted;
- (e) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R, wherein in the intron as indicated in column "Intron" of the following Table B the nucleotide as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1 is replaced by another nucleotide

**Table B**

<b>Intron</b>	<b>REPLACED NUCLEOTIDE</b>	<b>Position in wild-type</b>
intron 1	G	3166
intron 1	C	24778
intron 1	C	24830
intron 3	A	26308
intron 3	G	26422
intron 4	G	32394
intron 4	T	32434
intron 5	A	32783
intron 6	G	35641
intron 6	A	35725
intron 6	T	36001

intron 7	G	36378
intron 7	T	36387
intron 7	G	36398
intron 9	C	47214
intron 11	T	47563
intron 12	C	54307
intron 12	G	54308

- (f) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 3'UTR region corresponding to position 55169, 55170, 55171, 55917 or 54925 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;
- (g) a nucleotide sequence comprising at least 20 or 21 nucleotides and comprising the mutations or deletions as defined in any one of (a) to (f);
- (h) a nucleic acid sequence comprising a nucleotide sequence as shown in any one of SEQ ID NOs: 13 to 51;
- (i) a nucleic acid sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NOs: 5 to 12;
- (j) a nucleotide sequence which hybridizes to a nucleotide sequence defined in any one of (a) to (g) or to the nucleotide sequence of (h) and having a mutation as defined in any one of (a) to (f); and
- (k) a nucleic acid sequence being degenerate as a result of the genetic code to the nucleic acid sequence as defined in (j).

It has surprisingly been found that mutations in the P2X7R gene which encodes the ATP-gated ion channel P2X7R can cause the whole spectrum of affective disorders. Six different mutations in the 5'UTR of the P2X7R gene, seven different mutations in exons 3, 5, 6, 8 and 13 of the P2X7R gene leading to an amino acid replacement of the corresponding amino acid in the wild-type sequence of P2X7R depicted in SEQ ID NO: 3 or 4 and two mutations in exons 5 and 8 of said gene, respectively, leading to a replacement of a nucleotide by another nucleotide, a deletion of nucleotides in exon 13 of said gene, 18 mutations in introns 1, 3, 4, 5, 6, 7, 9, 11 and 12 and 5 mutations in the 3'UTR of the P2X7R gene have been identified to co-segregate with the affection status in 41 unrelated families affected with affective disorders. The

term "affective disorder" when used in the context of the present invention means to include, but is not limited to, depression, anxiety, unipolar disorder, bipolar disorder type I, bipolar disorder type II, mania, attention deficit hyperactive disorder, substance abuse, and any other disorders affecting the normal behaviour, or mood of an individual.

Each mutation causes alterations that can explain affective disorders as shown in the Examples hereinbelow.

P2X7R is an ATP-gated ion channel belonging to the P2X ionotropic channel family. The gene was first isolated from rat brain (Surprenant et al., (1996), 272, 735-738; Genbank accession number NM\_019256) and subsequently from a human monocyte library (Rassendren et al., J. Biol. Chem. 272 (1997), 5482-5486; Genbank accession numbers NM\_002562, Y09561) by virtue of its sequence homology with the other members of the P2X family. It was later found that P2X7R corresponded to the unidentified P2Z receptor which mediates the permeabilising action of ATP on mast cells and macrophages (Dahlgvist and Diamant, Acta Physiol. Scand. 34 (1974), 368-384; Steinberg and Silverstein, J. Biol. Chem. 262 (1987), 3118-3122; Gordon, Biochem. J. 233 (1986), 309-319). The P2X7R has two hydrophobic membrane-spanning domains, an extracellular loop, and forms transmembrane ion channels. P2X7 receptors seem to function only in homooligomeric form and bear a pharmacological profile markedly different from other P2X homo- or heteromers (North and Surprenant, Annual Rev. Pharmacology Toxicology 40 (2000), 563-580). P2X7R requires levels of ATP in excess of 1 mM to achieve activation, whereas other P2X receptors activate at ATP concentrations of  $\leq 100$   $\mu$ M (Steinberg et al., J. Biol. Chem. 262 (1987), 8884-8888; Greenberg et al., J. Biol. Chem. 263 (1988), 10337-10343) 32). While all P2X receptors demonstrate non-selective channel-like properties following ligation, the channels formed by the P2X7R can rapidly transform into pores that can allow the passage of molecules of up to 900 Dalton (Virginio et al., J. Physiol. 519 (1999), 335-346).

P2X7R is expressed in hematopoietic cells, mast cells and macrophages (Surprenant et al., Science 272 (1996), 3118-3122), where it is organized in tetrameric or hexameric form (Kim et al., J. Biol. Chem. 276 (2001), 23262-23267). P2X7R is inter alia involved in the regulation of the immune function and inflammatory response.

Activation of P2X7R by ATP in macrophages is associated with mitogenic stimulation of T cells (Baricordi et al., *Blood* 87 (1996), 682-690), the release of cytokines such as interleukin-1 $\beta$  (Griffiths et al., *J. Immunol.* 154 (1995), 2821-2828), and formation of macrophage polykations (Falzone et al., *J. Clin. Invest.* 95 (1995), 1207-1216). Stimulation of the P2X7R with ATP can also result in cell death by triggering massive transmembrane ion fluxes (particularly influx of Ca<sup>2+</sup> and Na<sup>+</sup>, and efflux of K<sup>+</sup>) and the formation of non-selective plasma membrane pores (Di Virgilio et al., *Cell Death Differ.* 5 (1998), 191-199).

In the brain, P2X7R was originally thought to be restricted to microglia (resident macrophage of the brain) and ependymal cells rather than neurons (Collo et al., *Neuropharmacology* 36 (1997), 1277-1283) suggesting a role of P2X7R in neurodegeneration. However, P2X7R has since been found in neurons of the rat retina (Brandle et al., *Brain Research Molecular Brain Res.* 62 (1998), 106-109), cochlear ganglion cells (Brandle et al., *Neuroscience Letters* 273 (1999), 105-108), and presynaptic terminals of neurons throughout the brainstem and spinal cord (Deuchards et al., *J. Neurosci.* 21 (2001), 7143-7152). Subsequent studies also suggest that P2X7R regulates the release of neurotransmitters such as glutamate and GABA in neurons of the hippocampus (Armstrong et al., *J. Neuroscience* 22 (2002), 5938-5945, Sperlágh et al., *J. Neurochem.* 81 (2002), 1196-1211). Organisation of P2X7R in glial cells and astrocytes of the brain appears monomeric (Kim et al., *J. Biol. Chem.* 276 (2001), 23262-23267).

Several agonists and antagonists of P2X7R have been identified. Brilliant Blue (Jiang et al., *Mol. Pharmacol.* 58 (2000), 82-88), the isoquinolines 1-[*N,O*-Bis(5-isoquinolinesulfonyl)-*N*-methyl-*L*-tyrosyl]-4-phenylpiperazine and *N*-[1-[*N*-methyl-*p*-(5-isoquinolinesulfonyl) benzyl]-2-(4-phenylpiperazine)ethyl]-5-isoquinolinesulfonamide (Humphreys et al., *Mol. Pharmacol.*, 54 (1998), 22-32), adamantane derivatives (WO 99/29660, WO 99/29661, WO 00/61569, WO 01/42194, WO 01/44170, WO 01/44213), substituted phenyl compounds (WO 00/71529), piperidine and piperazine derivatives (WO 01/46200) are antagonists of P2X7R while Oxidized ATP (oATP) acts as an irreversible inhibitor of the receptor (Chen et al., *J. Biol. Chem.*, 268 (1993), 8199-8203). Some of these antagonists are presently being evaluated for the treatment of inflammatory, immune, and cardiovascular diseases. BzATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate (C<sub>24</sub>H<sub>24</sub>N<sub>5</sub>O<sub>15</sub>P<sub>3</sub>)) acts as agonist of

P2X7R (North and Surprenant, Annu. Rev. Pharmacol. Toxicol. 40 (2000), 563-580). WO 99/55901 describes a method for identifying compounds that modulate the activity of a mammalian purinoreceptor selected from the group consisting of P2X2, P2X3, P2X4, P2X5, P2X6 and P2X7 and suggests a role of said purinoreceptors in therapy of behavioural disorders such as epilepsy, depression and aging-associated degenerative diseases.

Mutant mice lacking P2X7R are healthy, fertile and demonstrate no overt phenotype. However, in contrast to their wild-type counterparts, LPS-activated peritoneal macrophages from *P2X7R*<sup>-/-</sup> animals fail to generate mature interleukin-1 $\beta$  (IL-1 $\beta$ ) when challenged with ATP suggesting an inability of peritoneal macrophages to release IL-1 in response to ATP (Solle et al, J. Biol. Chem. 276 (2001), 125-132). A detailed behavioural study of the P2X7R<sup>-/-</sup> mice was not performed. In humans, a Glu-496 to Ala polymorphism leads to the loss of P2X7 function (Gu et al., J. Biol. Chem. 276 (2001), 11135–11142) and is associated with B-cell chronic lymphocytic leukaemia (Thunberg, et al, The Lancet 360 (2002), 1935-1939). Additional polymorphs in the putative P2X7R promoter region, and coding region have been reported (Li et al., FEBS Lett. 531 (2002), 127-131; EP 1199372).

Despite the abundant literature concerning P2X7R, a role in affective disorders has never been suggested or alluded to in the prior art.

Before the present invention is described in detail, it is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described herein as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art.

Preferably, the terms used herein are defined as described in "A multilingual glossary of biotechnological terms: (IUPAC Recommendations)", Leuenberger, H.G.W, Nagel, B. and Kölbl, H. eds. (1995), Helvetica Chimica Acta, CH-4010 Basel, Switzerland).

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integer or step.

Several documents are cited throughout the text of this specification. Each of the documents cited herein (including all patents, patent applications, scientific publications, manufacturer's specifications, instructions, etc.), whether supra or infra, are hereby incorporated by reference in their entirety. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the", include plural referents unless the context clearly indicates otherwise. Thus, for example, reference to "a reagent" includes one or more of such different reagents, and reference to "the method" includes reference to equivalent steps and methods known to those of ordinary skill in the art that could be modified or substituted for the methods described herein.

In accordance with the present invention, the term "nucleic acid sequence" means the sequence of bases comprising purine- and pyrimidine bases which are comprised by nucleic acid molecules, whereby said bases represent the primary structure of a nucleic acid molecule. Nucleic acid sequences include DNA, cDNA, genomic DNA, RNA, synthetic forms and mixed polymers, both sense and antisense strands, or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those skilled in the art.

When used herein, the term "polypeptide" means a peptide, a protein, or a polypeptide which encompasses amino acid chains of a given length, wherein the amino acid residues are linked by covalent peptide bonds. However, peptidomimetics of such proteins/polypeptides wherein amino acid(s) and/or peptide bond(s) have been replaced by functional analogs are also encompassed by the invention as well as other than the 20 gene-encoded amino acids, such as selenocysteine. Peptides, oligopeptides and proteins may be termed polypeptides. The terms polypeptide and



protein are often used interchangeably herein. The term polypeptide also refers to, and does not exclude, modifications of the polypeptide, e.g., glycosylation, acetylation, phosphorylation and the like. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature.

The term "position" used in accordance with the present invention means the position of either an amino acid within an amino acid sequence depicted herein or the position of a nucleotide within a nucleic acid sequence depicted herein.

The term "ATP-gated ion channel P2X7R", in accordance with this invention, denotes a polypeptide which can be classified as a member of the P2X ionotropic receptor family. They are also known as purinergic receptors. P2X receptors are ligand-gated ion channels. The ligand for these receptors may be ATP and/or another natural nucleotide such as ADP, UTP and UDP, or a synthetic nucleotide such as 2-methylthioATP. The criteria for the classification are: (1) a sequence homology that is higher than 39% across the family or different species; (2) signal transduction mechanism involving ion conductance (Khakh et al., Pharmacol Rev. 253 (2001), 107-18). Accordingly, the term "ATP-gated ion channel P2X7R" is interchangeable with the terms "ionotropic receptor" or "purinergic receptor". Preferably, the term "ATP-gated ion channel P2X7R" denotes a polypeptide which can be classified as an ATP-gated ion channel P2X7R on the basis of one or more structural and/or functional characteristics, preferably those described above. Structural characteristics refer to certain structural features which allow to classify a polypeptide as being a P2X7R protein. One such feature is the amino acid sequence. In the context of the present invention a polypeptide is classified as an ATP-gated ion channel P2X7R if it shows a certain degree of sequence identity over its own length to the amino acid sequence of the human P2X7R protein depicted in SEQ ID NO: 3 or 4. This degree of sequence identity is at least 40%, more preferably at least 50%, even more preferably at least 60%, at least 70%, at least 80%, at least 90% or at least 95%. It is particularly preferred that the degree of sequence identity is at least 65%.

Moreover; structural characteristics of P2X7R proteins are two hydrophobic membrane-spanning domains, an extra cellular loop which could be analyzed by using the program TMPRED (Hofmann Biol. Chem. 347 (1993), 166) or TMHMM (Krogh J. Mol. Bio. 305 (2001), 567-580). Additionally, P2X7R may exist as a single polypeptide, as dimer, tetramer or the like.

Thus, in the context of the present invention a protein is preferably classified as a P2X7R protein if it displays at least one of the above-mentioned structural characteristics. Functional characteristics refer to properties related to the biological activity of the P2X7R protein. In particular, P2X7R is an ATP-gated ion channel which allows calcium and sodium ions to pass from extracellular solution to intracellular solution, and allows potassium ions to pass from intracellular to extracellular solution. Moreover, the ATP-gated ion channel P2X7R forms naturally a homooligomeric form. The characteristics of P2X7R receptor proteins can be determined as mentioned hereinbelow. The term "ATP-gated ion channels P2X7R" comprises functional and non-functional forms of the ATP-gated ion channels P2X7R. A functional ATP-gated ion channel P2X7R is understood to be a P2X7R protein which has at least one of the above-mentioned functional characteristics which can be measured by methods known in the art. A non-functional ATP-gated ion channel P2X7R is a protein which can be classified as a P2X7R protein due to structural characteristics as described above but which has lost at least one, preferably all, functional characteristics of a P2X7R protein as described above. Non-functionality of the P2X7R protein can, e.g., be determined by measuring whether calcium and sodium ions can flow into cells or whether potassium ions can exit from cells. Thus, it is possible to determine the occurrence of a mutation in the ATP-gated ion channel P2X7R by measuring either calcium and/or sodium influx or efflux of cells. Cells harbouring a mutation in the P2X7R gene show an altered ion influx and/or efflux in comparison to cells harbouring a wild-type P2X7R protein.

Additionally, there are different methods that could be used to determine whether the P2X7R is functional or non-functional, for example, altered. One method consists of measuring the rate of ATP-induced incorporation of ethidium into cells, e.g. cells isolated from an individual. Ethidium is incorporated into the cells through P2X7R pores, when the pore formation is activated by ATP. Cells are then incubated with or

without ATP in the presence of ethidium, then they are analyzed by flow cytometry. Ethidium fluorescence is measured and compared in the presence or absence of ATP. If the P2X7R has lower activity, the ethidium fluorescence induced by ATP will be lower than in control cells. Such a method was used to verify P2X7R activity in isolated B-lymphocytes and T-lymphocytes from leukaemia patients (Wiley et al., Lancet 359 (2002), 1114-1119). Briefly, isolated cells are incubated in 1 ml of Hepes buffered potassium chloride at 37°C with continuous stirring. Ethidium is then added at a concentration of 25 mol/l, followed 40 seconds later by the addition of 10 µl of 100 mmol/l ATP stock. Cells are analyzed at 1,000 events/s by flow cytometry using a Coulter Elite flow cytometer (Coulter, Hialeah, FL) with argon laser excitation at 488 nm. Fluorescent emission was collected using a 590-nm long-pass filter. The linear mean channel fluorescence intensity for each gated subpopulation over successive 5-s intervals was analyzed with the use of Win-MDI software (Joseph Trotter, version 2.7) and plotted against time.

Another method of determining P2X7R activity is to measure calcium entry into isolated cells incubated with fluorescent dyes that emit only upon binding to calcium. The cells have to be loaded with the dye and then the calcium entry has to be stimulated. Examples of such dyes include Fura-2, Calcium green, calcium orange, calcium crimson (all available from Molecular Probes). Methods of measuring calcium transport are well known in the art; see for example, Takahashi et al., Physiol Rev. 79 (1999), 1089-1125. Furthermore, calcium entry into the cells produces changes in the membrane electric potential. This changes can be measured by electrophysiology (patch clamp) or by using dyes which are sensitive to voltage change. Such methods are also well known in the art, see for example, Gonzalez et al., DDT 4 (1999), 431-439; González and Tsien, Chemistry & Biology 4 (1997), 269-277; González and Tsien, Biophysical Journal 69 (1995), 1272-1280.

Yet another method is to measure uptake of <sup>133</sup>Ba21. Ba21 is a good surrogate for Ca21 and once inside the cell is neither pumped nor sequestered by transport ATPases. Ba21 uptake can be measured over 60 s using <sup>133</sup>BaCl2 (final concentration, 0.2 mM). At time 0, a prewarmed stock solution of <sup>133</sup>Ba21 (0.4 mM and 1 µCi/ml) is added in equal volumes to prewarmed isolated cells in 150 mM KCl with HEPES (pH 7.4) at 37°C. ATP (1 mM) is added either 10 minutes before or simultaneously with the <sup>133</sup>Ba21 isotope. Aliquots of 0.8 ml are taken at time points

between 0 and 60 s and are immediately mixed with 0.2 ml of ice-cold 50 mM MgCl<sub>2</sub> (in KCl-HEPES medium) that had been previously layered over 250 µl of oil mixture (di-n-butyl phthalate and di-iso-octyl phthalate, 7:3 vol/vol) and then centrifuged at 8,000 g for 30 s. The supernatants and the oil are aspirated, and the cell pellets are counted in a Wallac Wizard 3 automatic gamma-counter or in any other suitable gamma measuring unit.

The present invention is based on the finding that mutations of different kinds in the P2X7R gene are linked to the occurrence of affective disorders. The first type of mutations are mutations in the 5'UTR. Examples of such mutations are single nucleotide replacements at positions corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as described in SEQ ID NO: 1.

The position with respect to nucleotide sequences mentioned herein refer to the sequence shown in SEQ ID NO: 1. This sequence represents the nucleic acid sequence of the P2X7R gene encoding the ATP-gated ion channel P2X7R. It is possible for the skilled person to identify the position in the genomic sequence corresponding to a position in SEQ ID NO: 1 by aligning the sequences. Moreover, the exact locations of the exons and introns are indicated in SEQ ID NO: 1 hereinbelow. Additionally, the person skilled in the art is able to identify exons and introns of the P2X7R gene by comparing SEQ ID NO: 1 with SEQ ID NO: 2 which shows the cDNA sequence of the P2X7R gene.

Preferably, at position 362 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a thymine (T) is replaced by another nucleotide, preferably a purine base. More preferably, at said position said thymidine is replaced by a pyrimidine base. Particularly preferred, said thymine is replaced by a cytosine (C).

At position 532 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a thymine (T) is preferably replaced by another nucleotide, preferably a pyrimidine base. More preferably, at said position said thymine is replaced by a purine base. Particularly preferred, said thymidine is replaced by a guanine (G).

The adenine (A) residues at positions 1100 and 1122, respectively, in the 5'UTR of the genomic sequence of the P2X7R gene is preferably replaced by a pyrimidine base. More preferably, said adenine is replaced by a purine base and particularly preferred said adenine is replaced by a guanine (G).

At position 1171 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a cytidine (C) is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, said cytidine is replaced by a purine base and even more preferred, said cytidine is replaced by a guanine (G).

The guanine at position 1702 in the 5'UTR of the genomic sequence of the gene P2X7R depicted in SEQ ID NO: 1 is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, said guanine is replaced by a purine base and particularly preferred it is replaced by an adenine (A).

A second type of mutation found in the P2X7R gene are mutations in exons which lead to amino acid substitutions in the corresponding amino acid sequence. These are the mutations listed under item (b), supra. In this context, the term "an amino acid residue as indicated in column 'Amino acid residue' of Table A corresponding to position X of the wild-type ATP-gated ion channel P2X7R as depicted in column 'Position in wild-type' " has the following meaning: The amino acid residue in question would be located at position X in the sequence of SEQ ID NO: 3 or 4 if the sequence in which said amino acid residue occurs is compared and aligned with the amino acid sequence of SEQ ID NO: 3 or 4. The amino acid sequence shown in SEQ ID NO: 3 or 4 is the amino acid sequence of the human P2X7R gene and is used as a reference sequence in the present invention.

In order to determine whether an amino acid residue or nucleotide residue in a given P2X7R sequence corresponds to a certain position in the amino acid sequence or nucleotide sequence of SEQ ID NO: 1, 3 or 4, the skilled person can use means and methods well-known in the art, e.g., alignments, either manually or by using computer programs such as those mentioned further down below in connection with the definition of the term "hybridization" and degrees of homology.

For example, BLAST2.0, which stands for Basic Local Alignment Search Tool (Altschul, Nucl. Acids Res. 25 (1997), 3389-3402; Altschul, J. Mol. Evol. 36 (1993), 290-300; Altschul, J. Mol. Biol. 215 (1990), 403-410), can be used to search for local sequence alignments. BLAST produces alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in determining exact matches or in identifying similar sequences. The fundamental unit of BLAST algorithm output is the High-scoring Segment Pair (HSP). An HSP consists of two sequence fragments of arbitrary but equal lengths whose alignment is locally maximal and for which the alignment score meets or exceeds a threshold or cutoff score set by the user. The BLAST approach is to look for HSPs between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. The parameter E establishes the statistically significant threshold for reporting database sequence matches. E is interpreted as the upper bound of the expected frequency of chance occurrence of an HSP (or set of HSPs) within the context of the entire database search. Any database sequence whose match satisfies E is reported in the program output.

Analogous computer techniques using BLAST (Altschul (1997), loc. cit.; Altschul (1993), loc. cit.; Altschul (1990), loc. cit.) are used to search for identical or related molecules in nucleotide databases such as GenBank or EMBL. This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score which is defined as:

$$\frac{\% \text{sequence identity} \times \% \text{maximum BLAST score}}{100}$$

100

and it takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

As mentioned above, the second group of mutations identified in the P2X7R gene are mutations in the exons of the P2X7R gene which lead to amino acid substitutions. In this respect SEQ ID NO 2 shows the cDNA sequence of the P2X7R gene. In exon 3 at position 117 of the corresponding wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 an arginine (R) residue is replaced by another amino acid residue, preferably by an aliphatic, acidic or basic amino acid residue. More preferably, by an aromatic amino acid residue which is particularly preferred to be a tryptophane (W). The resulting polypeptide is shown in SEQ ID NO: 5.

In exon 5 at position 150 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 a glycine (G) residue is replaced by another amino acid residue, preferably by an aliphatic, aromatic or acidic amino acid residue. More preferably, by a basic amino acid residue and particularly preferred by an arginine (R). The resulting polypeptide is shown in SEQ ID NO: 6.

At position 186 in exon 6 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 a glutamate residue (E) is replaced by another amino acid residue, preferably by an aliphatic, aromatic or acidic amino acid residue. More preferably, said glutamate is replaced by a basic amino acid residue which is particularly preferred a lysine (K). The resulting polypeptide is shown in SEQ ID NO: 7.

In exon 6 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 at position 191 a leucine residue (L) is replaced by another amino acid residue. Said amino acid residue is preferably an aliphatic, acidic or basic amino acid residue. More preferably, said amino acid residue is an aromatic amino acid residue which is particularly preferred to be a proline (P). The resulting polypeptide is shown in SEQ ID NO: 8.

In exon 8 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 at position 270 an arginine residue (R) is replaced by another amino acid residue. Said amino acid residue is preferably an aromatic, acidic or basic amino acid residue. More preferably, said amino acid residue is an aliphatic amino acid residue

which is particularly preferred to be a cysteine (C). The resulting polypeptide is shown in SEQ ID NO: 9.

At position 568 in exon 13 of the wild-type amino acid residue of P2X7R depicted in SEQ ID NO: 3 or 4 an isoleucine (I) residue is replaced by another amino acid residue. More preferably, said isoleucine is replaced by an aromatic, basic or acidic amino acid residue. Even more preferred, said isoleucine is replaced by an aliphatic amino acid residue which is particularly preferred to be an asparagine (N). The resulting polypeptide is shown in SEQ ID NO: 10.

In exon 13 at position 578 in the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 an arginine residue (R) is replaced by another amino acid residue. Said amino acid residue is preferably an aromatic, acidic or basic amino acid residue. More preferably, it is an aliphatic amino acid residue and particularly preferred it is a glutamine (Q) residue. The resulting polypeptide is shown in SEQ ID NO: 12.

It is envisaged that the above-mentioned mutations in the exons of the P2X7R gene occur due to point mutations caused by, e.g. chemical and/or physical means or inaccuracy of the replication complex followed by a failure of the reparation machinery of a cell, a change of a single codon occur can be achieved. Possible types of point mutations are transitions, i.e. change of a purine or pyrimidine base for another purine or pyrimidine base, e.g. adenine to guanine or thymidine to cytosine or transversions, i.e. change of a purine or pyrimidine base for another pyrimide or purine base, e.g., adenine to thymidine or guanine to cytosine. Additionally a point mutation can also be caused by insertion or deletion of one or more nucleotides.

The mutations leading to the replacement of the amino acids as mentioned hereinabove and hereinbelow are indicated in Table 1 hereinbelow.

---

The third group of mutations in the P2X7R gene has been identified to be in exons 5 and 8 of the P2X7R gene depicted in SEQ ID NO: 1 and to be silent, i.e. they do not lead to amino acid changes. In particular, at position 32548 in exon 5 of the wild-type



genomic sequence P2X7R gene depicted in SEQ ID NO: 1 a cytidine residue is replaced by another nucleotide. Said nucleotide is preferably a pyrimidine base and particularly preferred a thymine. The exchange of the cytidine residue at position 32548 in exon 5 of the P2X7R gene by another nucleotide preferably does not lead to the replacement of the amino acid cysteine by another amino acid residue.

In exon 8 of the wild-type P2X7R gene depicted in SEQ ID NO: 1 at position 37633 a cytidine residue is replaced by another nucleotide residue. Said nucleotide residue is preferably a pyrimidine base and particularly preferred thymine. Due to this replacement the amino acid aspartate (D) encoded by the respective codon in which at position 37633 a replacement has taken place is preferably not replaced by another amino acid residue.

The above-mentioned mutations in exons 5 and 8 at positions 32548 and 37633, respectively, of the wild-type P2X7R gene depicted in SEQ ID NO:1 are mutations at the third position of a triplet codon, i.e. at the wobble base, which lead to so-called silent mutations. Silent mutations do normally not lead to a change of the amino acid due to the degeneracy of the genetic code, i.e. 64 triplets encode at all 20 naturally occurring amino acids. However, said silent mutations lead to a change in the codon encoding its respective amino acids insofar that the newly generated codon may not fit so well into the codon usage of an organism. Namely, the newly generated codon is not translated by the ribosome with the same efficiency as the "old" codon. This may lead to insufficient amounts of the corresponding polypeptide causing an distinct phenotype.

The fourth group of mutations in the P2X7R gene described hereinabove in item (d) is a deletion of 7 amino acids corresponding to positions 488 to 494 of the wild-type P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4. Thus, the present invention also relates to nucleic acid sequences encoding a P2X7R protein in which amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted. This means, according to the present invention, that a fragment encompassing amino acid positions 488 to 494 of the corresponding wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4 is deleted which results in a shortened polypeptide. An example

for such a shortened polypeptide is depicted in SEQ ID NO: 11. This type of mutation as described herein preferably encodes a non-functional ATP-gated ion channel P2X7R. In the present invention the deletion of a fragment encompassing amino acids 488 to 494 of the wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4 is the result of a deletion in exon 13. The resulting protein depicted in SEQ ID NO: 11 lacks amino acids 488 to 494 of the corresponding wild type amino acid sequence depicted in SEQ ID NO: 3 or 4 such that amino acid position 494 of the deleted polypeptide depicted in SEQ ID NO: 11 corresponds to amino acid position 502 of the wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4. Preferably, the nucleic acid sequence of the invention encodes a P2X7R polypeptide in which exactly amino acids corresponding to positions 488 to 494 of SEQ ID NO: 3 or 4 are deleted. However, also mutants are comprised in which either more or less amino acids within the P2X7R amino acid sequence set forth in SEQ ID NO: 3 or 4 may be deleted due to, for example, atypical splicing or deletion of nucleotides of the nucleic acid molecule encoding P2X7R or wrong posttranslational processes, as long as the P2X7R ATP-gated ion channel is non-functional. For example, it is also possible that further amino acids preceding amino acid position 488 or amino acids succeeding amino acid position 494 may be deleted or that less amino acids are deleted.

Preferably at least one, more preferably at least two, even more preferably at least three and most preferably at least 5 amino acid residues are further deleted upstream from the position corresponding to amino acid residue 488 and/or downstream of the position corresponding to amino acid residue 494 of SEQ ID NO: 3 or 4.

However, it is preferred that not more than 20, preferably not more than 15, even more preferably not more than 10 and most preferably not more than 7 amino acid residues are further deleted upstream of the position corresponding to amino acid residue 488 of SEQ ID NO: 3 or 4 or downstream of the position corresponding to amino acid residue 494 of SEQ ID NO: 3 or 4.

---

Another group of mutation (mentioned in item (e), supra) resides in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 of the wild-type genomic sequence of P2X7R depicted in SEQ ID NO: 1. Said mutations in said introns are point mutations as shown in Table B hereinabove and in Table 1, hereinbelow.

At the respective position indicated in the column "Position in wild-type" in Table B or indicated in the column "Polymorphism" in Table 1 the position of the nucleotide residue in the respective intron which is replaced by another nucleotide residue is shown. Accordingly, the term "a nucleotide as indicated in column "Intron" of the Table B corresponding to the position as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B is replaced by another nucleotide means that a nucleotide residue in a P2X7R encoding sequence would be located at position Y in SEQ ID NO: 1 when the P2X7R sequence is compared and aligned with the sequence of SEQ ID NO: 1.

If the nucleotide at the respective position is a purine base such as adenine or guanine it is preferred that due to a transition it is replaced by another purine base. For example, an adenine is replaced by a guanine or a guanine is replaced by an adenine. If the nucleotide at the respective position is a pyrimidine base it is preferred that due to a transition it is replaced by another pyrimidine base. For example, thymine is replaced by a cytidine and a cytidine is replaced by a thymine.

It is also preferred that due to a transversion a purine base is replaced by a pyrimidine base or vice versa. For example, an adenine is replaced by a thymine and a guanine is replaced by a cytidine. Particularly preferred, said nucleotide in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 of the P2X7R gene depicted in SEQ ID NO: 1 is replaced by the nucleotide depicted in column "Polymorphism" of Table 1, hereinbelow.

A last group of mutations that has been identified relates to mutations which reside in the 3'UTR of the wild-type P2X7R gene depicted in SEQ ID NO: 1. The mutations were found at positions 54925, 55169, 55170, 55171 or 55917 respectively, of the wild-type P2X7R gene depicted in SEQ ID NO: 1.

At position 54925 a guanine residue was found to be replaced by another nucleotide. Preferably, said guanine residue is replaced by a pyrimidine base, more preferably by a purine base and particularly preferred by an adenine.

At position 55169 a cytidine residue is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, it is replaced by a purine base and particularly preferred, it is replaced by an adenine.

At positions 55170 and 55171 an adenine residue is replaced by another nucleotide residue, preferably by a purine base. More preferably, said adenine residue is replaced by a pyrimidine base and particularly preferred said adenine residue is replaced by a cytidine residue. It was also found that at position 55917 a cytidine residue is replaced by another nucleotide. Preferably, said nucleotide residue is a purine base, more preferably a pyrimidine base and particularly preferable a thymine.

As is evident from the above, not all identified mutations are located in exons or lead to a change in the amino acid sequence. Some of the mutations are located in the 5'UTR, the 3'UTR or in introns.

It is known that polymorphisms in promoter and enhancer regions can affect gene function by modulating transcription, particularly if they are situated at recognition sites for DNA binding proteins (Fishman et al., J. Clin. Invest. 102 (1998), 1369-1376). The term "polymorphism" which is used in the present invention means single nucleotide substitution, nucleotide insertion and nucleotide deletion which in the case of insertion and deletion includes insertion or deletion of one or more nucleotides at a position of a gene and corresponding alterations in expressed proteins. Polymorphisms in the 5' untranslated region (5'UTR) of genes can affect the efficiency with which proteins are translated. A representative example of this is in the c-myc gene where a C-G SNP that creates an internal ribosome entry site is associated with increased efficiency of c-myc translation and myeloma (Chappell et al., Oncogene 19 (2000), 4437-4440). Polymorphisms in the 3'UTR can affect gene function by altering the secondary structure of RNA and efficiency of translation or by affecting motifs in the RNA that bind proteins which regulate RNA degradation. Polymorphisms within introns can affect gene function by affecting RNA splicing resulting in aberrant polypeptides. Another way in which intronic polymorphisms can affect gene function is when they affect regulatory motifs within introns. Examples are the Sp1 binding site polymorphism within intron 1 of the COLIA1 gene (Mann et al., J. Clin. Invest 107 (2001), 899-907) and a repeat polymorphisms within the IL-1Ra

gene (Keen et al., Bone 23 (1998), 367-371). Further examples between intronic SNPs and gene function are described in Caceres and Kornblihtt, Trends Genet. 4 (2002), 186-93. Example 4 on page 52, line 30 to page 53, line 51 of the text describes potential alternative splicing events and aberrant protein production associated with three SNPs disclosed in the application.

The nucleic acid sequences described hereinabove may comprise at least 56580 nucleotides, preferably at least 10000 nucleotides, at least 5000 nucleotides, at least 1000 nucleotides, at least 500 nucleotides, at least 100 nucleotides. More preferably, said nucleic acid sequences comprise at least 50 nucleotides and particularly preferred they comprise at least 20 or 21 nucleotides comprising the mutations or deletions as described hereinabove. Most preferably such a nucleic acid sequence has a sequence as depicted in any one of SEQ ID NOs: 13 to 51.

The nucleic acid sequences described hereinabove which comprise mutations in exons leading to a replacement of the corresponding amino acid sequence of the P2X7R wild-type polypeptide depicted in SEQ ID NO: 3 or 4 encode polypeptides shown in SEQ ID NOs: 5 to 10 and 12

Additionally, the nucleic acid sequences described hereinabove which comprise a deletion leading to a truncated polypeptide in comparison to the full-length polypeptide of the wild-type P2X7R polypeptide shown in SEQ ID NO: 3 or 4 is shown in SEQ ID NO: 11.

The present invention also relates to nucleic acid molecules which hybridize to one of the above described nucleic acid molecules and which shows a mutation as described hereinabove.

The term "hybridizes" as used in accordance with the present invention may relate to hybridizations under stringent or non-stringent conditions. If not further specified, the conditions are preferably non-stringent. Said hybridization conditions may be established according to conventional protocols described, for example, in Sambrook, Russell "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Laboratory, N.Y. (2001); Ausubel, "Current Protocols in Molecular Biology", Green Publishing Associates and Wiley Interscience, N.Y. (1989), or Higgins and Hames

(Eds.) "Nucleic acid hybridization, a practical approach" IRL Press Oxford, Washington DC, (1985). The setting of conditions is well within the skill of the artisan and can be determined according to protocols described in the art. Thus, the detection of only specifically hybridizing sequences will usually require stringent hybridization and washing conditions such as 0.1xSSC, 0.1% SDS at 65°C. Non-stringent hybridization conditions for the detection of homologous or not exactly complementary sequences may be set at 6xSSC, 1% SDS at 65°C. As is well known, the length of the probe and the composition of the nucleic acid to be determined constitute further parameters of the hybridization conditions. Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility. Hybridizing nucleic acid molecules also comprise fragments of the above described molecules. Such fragments may represent nucleic acid sequences which code for a non-functional ATP-gated ion channel P2X7R or a non-functional fragment thereof, and which have a length of at least 12 nucleotides, preferably at least 15, more preferably at least 18, more preferably of at least 21 nucleotides, more preferably at least 30 nucleotides, even more preferably at least 40 nucleotides and most preferably at least 60 nucleotides. Furthermore, nucleic acid molecules which hybridize with any of the aforementioned nucleic acid molecules also include complementary fragments, derivatives and allelic variants of these molecules. Additionally, a hybridization complex refers to a complex between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., Cot or Rot analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., membranes, filters, chips, pins or glass slides to which, e.g., cells have been fixed). The terms complementary or

complementarity refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, the sequence "A-G-T" binds to the complementary sequence "T-C-A". Complementarity between two single-stranded molecules may be "partial", in which only some of the nucleic acids bind, or it may be complete when total complementarity exists between single-stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands.

The term "hybridizing sequences" preferably refers to sequences which display a sequence identity of at least 40%, preferably at least 50%, more preferably at least 60%, even more preferably at least 70%, particularly preferred at least 80%, more particularly preferred at least 90%, even more particularly preferred at least 95% and most preferably at least 97% identity with a nucleic acid sequence as described above encoding a P2X7R protein having a described mutation. Moreover, the term "hybridizing sequences" preferably refers to sequences encoding a P2X7R protein having a sequence identity of at least 40%, preferably at least 50%, more preferably at least 60%, even more preferably at least 70%, particularly preferred at least 80%, more particularly preferred at least 90%, even more particularly preferred at least 95% and most preferably at least 97% identity with an amino acid sequence of a P2X7R mutant as described herein above.

In accordance with the present invention, the term "identical" or "percent identity" in the context of two or more nucleic acid or amino acid sequences, refers to two or more sequences or subsequences that are the same, or that have a specified percentage of amino acid residues or nucleotides that are the same (e.g., 60% or 65% identity, preferably, 70-95% identity, more preferably at least 95% identity), when compared and aligned for maximum correspondence over a window of comparison, or over a designated region as measured using a sequence comparison algorithm as known in the art, or by manual alignment and visual inspection. Sequences having, for example, 60% to 95% or greater sequence identity are considered to be substantially identical. Such a definition also applies to the complement of a test sequence. Preferably the described identity exists over a region that is at least about 15 to 25 amino acids or nucleotides in length, more preferably,

over a region that is about 50 to 100 amino acids or nucleotides in length. Those having skill in the art will know how to determine percent identity between/among sequences using, for example, algorithms such as those based on CLUSTALW computer program (Thompson Nucl. Acids Res. 2 (1994), 4673-4680) or FASTDB (Brutlag Comp. App. Biosci. 6 (1990), 237-245), as known in the art.

Although the FASTDB algorithm typically does not consider internal non-matching deletions or additions in sequences, i.e., gaps, in its calculation, this can be corrected manually to avoid an overestimation of the % identity. CLUSTALW, however, does take sequence gaps into account in its identity calculations. Also available to those having skill in this art are the BLAST and BLAST 2.0 algorithms (Altschul Nucl. Acids Res. 25 (1977), 3389-3402). The BLASTN program for nucleic acid sequences uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, and an expectation (E) of 10. The BLOSUM62 scoring matrix (Henikoff Proc. Natl. Acad. Sci., USA, 89, (1989), 10915) uses alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

Moreover, the present invention also relates to nucleic acid molecules the sequence of which is degenerate in comparison with the sequence of an above-described hybridizing molecule. When used in accordance with the present invention the term "being degenerate as a result of the genetic code" means that due to the redundancy of the genetic code different nucleotide sequences code for the same amino acid.

The present invention also related to nucleic acid molecules which comprise one or more of the above-described mutations or deletions.

The nucleic acid molecules according to the invention may be derived from any organism encoding corresponding P2X7R ATP-gated ion channels. For example, P2X7R ATP-gated ion channels have been reported in various organisms, for example, rat (see, Suprenant (1996), loc. cit.), mouse (Genbank Accession No. AJ 489297), xenopus (Genbank Accession No. AJ 345114), chicken (Genbank Accession No. BM 491404) or Bos Taurus (Genbank Accession No. AF 083073) . In



a preferred embodiment the nucleic acid molecule of the invention is derived from a vertebrate, preferably from a mammal, even more preferably the nucleic acid molecule is derived from rabbit or guinea pig, and most preferably the nucleic acid is derived from mouse, rat or human.

The nucleic acid molecule according to the invention may be any type of nucleic acid, e.g. DNA, RNA or PNA (peptide nucleic acid).

For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine, thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by Nielsen et al., Science 254:1497 (1991); and Egholm et al., Nature 365:666 (1993), PNAs bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point ( $T_{sub.m}$ ) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

The DNA may, for example, be cDNA. In a preferred embodiment it is a genomic DNA. The RNA may be, e.g., mRNA. The nucleic acid molecule may be natural, synthetic or semisynthetic or it may be a derivative, such as peptide nucleic acid (Nielsen, Science 254 (1991), 1497-1500) or phosphorothioates. Furthermore, the nucleic acid molecule may be a recombinantly produced chimeric nucleic acid molecule comprising any of the aforementioned nucleic acid molecules either alone or in combination.

Preferably, the nucleic acid molecule of the present invention is part of a vector. Therefore, the present invention relates in another embodiment to a vector comprising the nucleic acid molecule of this invention. Such a vector may be, e.g., a plasmid, cosmid, virus, bacteriophage or another vector used e.g. conventionally in genetic engineering, and may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions.

The nucleic acid molecules of the present invention may be inserted into several commercially available vectors. Nonlimiting examples include plasmid vectors compatible with mammalian cells, such as pUC, pBluescript (Stratagene), pET (Novagen), pREP (Invitrogen), pCRTopo (Invitrogen), pcDNA3 (Invitrogen), pCEP4 (Invitrogen), pMC1 neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), EBO-pSV2neo, pBPV-1, pdBPVMMTneo, pRSVgpt, pRSVneo, pSV2-dhfr, pUCTag, pIZD35, pLXIN and pSIR (Clontech) and pIRES-EGFP (Clontech). Baculovirus vectors such as pBlueBac, BacPacz Baculovirus Expression System (CLONTECH), and MaxBac<sup>TM</sup> Baculovirus Expression System, insect cells and protocols (Invitrogen) are available commercially and may also be used to produce high yields of biologically active protein. (see also, Miller (1993), Curr. Op. Genet. Dev., 3, 9; O'Reilly, Baculovirus Expression Vectors: A Laboratory Manual, p. 127). In addition, prokaryotic vectors such as pcDNA2; and yeast vectors such as pYes2 are nonlimiting examples of other vectors suitable for use with the present invention. For vector modification techniques, see Sambrook and Russel (2001), loc. cit. Vectors can contain one or more replication and inheritance systems for cloning or expression, one or more markers for selection in the host, e. g., antibiotic resistance, and one or more expression cassettes.

The coding sequences inserted in the vector can be synthesized by standard methods, isolated from natural sources, or prepared as hybrids. Ligation of the coding sequences to transcriptional regulatory elements (e. g., promoters, enhancers, and/or insulators) and/or to other amino acid encoding sequences can be carried out using established methods.

Furthermore, the vectors may, in addition to the nucleic acid sequences of the invention, comprise expression control elements, allowing proper expression of the coding regions in suitable hosts. Such control elements are known to the artisan and may include a promoter, translation initiation codon, translation and insertion site or internal ribosomal entry sites (IRES) (Owens, Proc. Natl. Acad. Sci. USA 98 (2001), 1471-1476) for introducing an insert into the vector. Preferably, the nucleic acid molecule of the invention is operatively linked to said expression control sequences allowing expression in eukaryotic or prokaryotic cells. Particularly preferred are in this context control sequences which allow for correct expression in neuronal cells and/or cells derived from nervous tissue.

Control elements ensuring expression in eukaryotic and prokaryotic cells are well known to those skilled in the art. As mentioned above, they usually comprise regulatory sequences ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers, and/or naturally-associated or heterologous promoter regions. Possible regulatory elements permitting expression in for example mammalian host cells comprise the CMV-HSV thymidine kinase promoter, SV40, RSV-promoter (Rous sarcome virus), human elongation factor 1 $\alpha$ -promoter, CMV enhancer, CaM-kinase promoter or SV40-enhancer.

For the expression for example in nervous tissue and/or cells derived therefrom, several regulatory sequences are well known in the art, like the minimal promoter sequence of human neurofilament L (Charron, J. Biol. Chem. 270 (1995), 25739-25745). For the expression in prokaryotic cells, a multitude of promoters including, for example, the tac-lac-promoter, the lacUV5 or the trp promoter, has been described. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pRc/CMV, pcDNA1, pcDNA3 (Invitrogen, as used, inter alia in the appended examples), pSPORT1 (GIBCO BRL) or pGEMHE (Promega), or prokaryotic expression vectors, such as lambda gt11.

An expression vector according to this invention is at least capable of directing the replication, and preferably the expression, of the nucleic acids and protein of this invention. Suitable origins of replication include, for example, the Col E1, the SV40 viral and the M 13 origins of replication. Suitable promoters include, for example, the cytomegalovirus (CMV) promoter, the iacZ promoter, the gai10 promoter and the Autographa californica multiple nuclear polyhedrosis virus (AcMNPV) polyhedral promoter. Suitable termination sequences include, for example, the bovine growth hormone, SV40, iacZ and AcMNPV polyhedral polyadenylation signals. Examples of selectable markers include neomycin, ampicillin, and hygromycin resistance and the like. Specifically-designed vectors allow the shuttling of DNA between different host cells, such as bacteria-yeast, or bacteria-animal cells, or bacteria-fungal cells, or bacteria invertebrate cells.

Beside the nucleic acid molecules of the present invention, the vector may further comprise nucleic acid sequences encoding for secretion signals. Such sequences are well known to the person skilled in the art. Furthermore, depending on the expression system used leader sequences capable of directing the expressed polypeptide to a cellular compartment may be added to the coding sequence of the nucleic acid molecules of the invention and are well known in the art. The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein, or a part thereof, into, inter alia, the extracellular membrane. Optionally, the heterologous sequence can encode a fusion protein including an C- or N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. Once the vector has been incorporated into the appropriate host, the host is maintained under conditions suitable for high level expression of the nucleotide sequences, and, as desired, the collection and purification of the proteins, antigenic fragments or fusion proteins of the invention may follow. Of course, the vector can also comprise regulatory regions from pathogenic organisms.

Furthermore, said vector may also be, besides an expression vector, a gene transfer and/or gene targeting vector. Gene therapy, which is based on introducing therapeutic genes (for example for vaccination) into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable

vectors, vector systems and methods for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, Nature Medicine 2 (1996), 534-539; Schaper, Circ. Res. 79 (1996), 911-919; Anderson, Science 256 (1992), 808-813, Isner, Lancet 348 (1996), 370-374; Muhlhauser, Circ. Res. 77 (1995), 1077-1086; Wang, Nature Medicine 2 (1996), 714-716; WO 94/29469; WO 97/00957; Schaper, Current Opinion in Biotechnology 7 (1996), 635-640 or Verma, Nature 389 (1997), 239-242 and references cited therein. The nucleic acid molecules of the invention and vectors as described herein above may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Additionally, baculoviral systems or systems based on vaccinia virus or Semliki Forest Virus can be used as eukaryotic expression system for the nucleic acid molecules of the invention. In addition to recombinant production, fragments of the protein, the fusion protein or antigenic fragments of the invention may be produced by direct peptide synthesis using solid-phase techniques (cf Stewart et al. (1969) Solid Phase Peptide Synthesis; Freeman Co, San Francisco; Merrifield, J. Am. Chem. Soc. 85 (1963), 2149-2154). In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Foster City CA) in accordance with the instructions provided by the manufacturer. Various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

The present invention in addition relates to a host transformed with a vector of the present invention or to a host comprising the nucleic acid molecule of the invention. Said host may be produced by introducing said vector or nucleotide sequence into a host cell which upon its presence in the cell mediates the expression of a protein encoded by the nucleotide sequence of the invention or comprising a nucleotide sequence or a vector according to the invention wherein the nucleotide sequence and/or the encoded polypeptide is foreign to the host cell.

By "foreign" it is meant that the nucleotide sequence and/or the encoded polypeptide is either heterologous with respect to the host, this means derived from a cell or organism with a different genomic background, or is homologous with respect to the

host but located in a different genomic environment than the naturally occurring counterpart of said nucleotide sequence. This means that, if the nucleotide sequence is homologous with respect to the host, it is not located in its natural location in the genome of said host, in particular it is surrounded by different genes. In this case the nucleotide sequence may be either under the control of its own promoter or under the control of a heterologous promoter. The location of the introduced nucleic acid molecule or the vector can be determined by the skilled person by using methods well-known to the person skilled in the art, e.g., Southern Blotting. The vector or nucleotide sequence according to the invention which is present in the host may either be integrated into the genome of the host or it may be maintained in some form extrachromosomally. In this respect, it is also to be understood that the nucleotide sequence of the invention can be used to restore or create a mutant gene via homologous recombination.

Said host may be any prokaryotic or eukaryotic cell. Suitable prokaryotic/bacterial cells are those generally used for cloning like *E. coli*, *Salmonella typhimurium*, *Serratia marcescens* or *Bacillus subtilis*. Said eukaryotic host may be a mammalian cell, an amphibian cell, a fish cell, an insect cell, a fungal cell, a plant cell or a bacterial cell (e.g., *E. coli* strains HB101, DH5a, XL1 Blue, Y1090 and JM101). Eukaryotic recombinant host cells are preferred. Examples of eukaryotic host cells include, but are not limited to, yeast, e.g., *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis* or *Pichia pastoris* cells, cell lines of human, bovine, porcine, monkey, and rodent origin, as well as insect cells, including but not limited to, *Spodoptera frugiperda* insect cells and *Drosophila*-derived insect cells as well as zebra fish cells. Mammalian species-derived cell lines suitable for use and commercially available include, but are not limited to, L cells, CV-1 cells, COS-1 cells (ATCC CRL 1650), COS-7 cells (ATCC CRL 1651), HeLa cells (ATCC CCL 2), C1271 (ATCC CRL 1616), BS-C-1 (ATCC CCL 26) and MRC-5 (ATCC CCL 171).

---

In a particularly preferred embodiment said mammalian cell is a neuronal cell and/or a cultured cell like, inter alia, a HEK 293 (human embryonic kidney) cell, a CHO,

HeLa, NIH3T3, BHK, PC12 cell or a neuronal stem cell preferably derived from a mammal and more preferably from a human.

In another more preferred embodiment said amphibian cell is an oocyte. In an even more preferred embodiment said oocyte is a frog oocyte, particularly preferred a *Xenopus laevis* oocyte.

In a more preferred embodiment, the host according to the invention is a non-human transgenic organism. Said non-human organism may be a mammal, amphibian, a fish, an insect, a fungus or a plant. Particularly preferred non-human transgenic animals are *Drosophila* species, *Caenorhabditis elegans*, *Xenopus* species, zebra fish, *Spodoptera frugiperda*, *Autographa californica*, mice and rats. Transgenic plants comprise, but are not limited to, wheat, tobacco, parsley and *Arabidopsis*. Transgenic fungi are also well known in the art and comprise, inter alia, yeasts, like *S. pombe* or *S. cerevisiae*, or *Aspergillus*, *Neurospora* or *Ustilago* species or *Pichia* species.

In another embodiment, the present invention relates to a method for producing the polypeptide encoded by a nucleic acid molecule of the invention comprising culturing/raising the host of the invention and isolating the produced polypeptide.

A large number of suitable methods exist in the art to produce polypeptides in appropriate hosts. If the host is a unicellular organism or a mammalian or insect cell, the person skilled in the art can revert to a variety of culture conditions that can be further optimized without an undue burden of work. Conveniently, the produced protein is harvested from the culture medium or from isolated (biological) membranes by established techniques. Furthermore, the produced polypeptide may be directly isolated from the host cell. Said host cell may be part of or derived from a part of a host organism, for example said host cell may be part of the CNS of an animal or the harvestable part of a plant. Additionally, the produced polypeptide may be isolated from fluids derived from said host, like blood, milk or cerebrospinal fluid.

Additionally the present invention relates to polypeptides depicted in SEQ ID NOs: 5 to 12 which are encoded by the nucleic acid molecules of the invention or produced by the method of the invention. The polypeptide of the invention may accordingly be produced by microbiological methods or by transgenic mammals. It is also envisaged

that the polypeptide of the invention is recovered from transgenic plants. Alternatively, the polypeptide of the invention may be produced synthetically or semi-synthetically.

For example, chemical synthesis, such as the solid phase procedure described by Houghton Proc. Natl. Acad. Sci. USA (82) (1985), 5131-5135, can be used. Another method is in vitro translation of mRNA. A preferred method involves the recombinant production of protein in host cells as described above. For example, nucleotide acid sequences comprising all or a portion of any one of the nucleotide sequences according to the invention can be synthesized by PCR, inserted into an expression vector, and a host cell transformed with the expression vector. Thereafter, the host cell is cultured to produce the desired polypeptide, which is isolated and purified. Protein isolation and purification can be achieved by any one of several known techniques; for example and without limitation, ion exchange chromatography, gel filtration chromatography and affinity chromatography, high pressure liquid chromatography (HPLC), reversed phase HPLC, preparative disc gel electrophoresis. In addition, cell-free translation systems can be used to produce the polypeptides of the present invention. Suitable cell-free expression systems for use in accordance with the present invention include rabbit reticulocyte lysate, wheat germ extract, canine pancreatic microsomal membranes, E. coli S30 extract, and coupled transcription/translation systems such as the TNT-system (Promega). These systems allow the expression of recombinant polypeptides or peptides upon the addition of cloning vectors, DNA fragments, or RNA sequences containing coding regions and appropriate promoter elements. As mentioned supra, protein isolation/purification techniques may require modification of the proteins of the present invention using conventional methods. For example, a histidine tag can be added to the protein to allow purification on a nickel column. Other modifications may cause higher or lower activity, permit higher levels of protein production, or simplify purification of the protein.

---

In a further embodiment, the present invention relates to an antibody specifically directed to a polypeptide of the invention, wherein said antibody specifically reacts with an epitope generated and/or formed by the mutation in the ATP-gated ion channel P2X7R selected from the group consisting of:



- (i) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein the R (Arg), G (Gly), E (Glu), L (Leu), R (Arg), I (Ile) or R (Arg) residue corresponding to position 117, 150, 186, 191, 270, 568 or 578 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue; and
- (ii) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted.

With respect to preferred embodiments of (i) and (ii) the same applies as described above in connection with the nucleic acid molecules. The term "specifically" in this context means that the antibody reacts with the mutant P2X7R protein but not with a wild-type P2X7R protein. Preferably this term also means that such an antibody does not bind to other mutant forms of the P2X7R protein, in particular those described herein. Whether the antibody specifically reacts as defined herein above can easily be tested, inter alia, by comparing the reaction of said antibody with a wild-type ATP-gated ion channel P2X7R (or a subunit or a fragment thereof) with the reaction of said antibody with a mutant P2X7R polypeptide of the invention.

The antibody of the present invention can be, for example, polyclonal or monoclonal. The term "antibody" also comprises derivatives or fragments thereof which still retain the binding specificity. Techniques for the production of antibodies are well known in the art and described, e.g. in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. These antibodies can be used, for example, for the immunoprecipitation and immunolocalization of the polypeptides of the invention as well as for the monitoring of the presence of such polypeptides, for example, in recombinant organisms or in diagnosis. They can also be used for the identification of compounds interacting with the proteins according to the invention (as mentioned herein below). For example, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of the polypeptide of the invention (Schier, Human

Antibodies Hybridomas 7 (1996), 97-105; Malmborg, J. Immunol. Methods 183 (1995), 7-13).

The present invention furthermore includes chimeric, single chain and humanized antibodies, as well as antibody fragments, like, inter alia, Fab fragments. Antibody fragments or derivatives further comprise F(ab')<sub>2</sub>, Fv or scFv fragments; see, for example, Harlow and Lane, loc. cit.. Various procedures are known in the art and may be used for the production of such antibodies and/or fragments. Thus, the (antibody) derivatives can be produced by peptidomimetics. Further, techniques described for the production of single chain antibodies (see, inter alia, US Patent 4,946,778) can be adapted to produce single chain antibodies to polypeptide(s) of this invention. Also, transgenic animals may be used to express humanized antibodies to polypeptides of this invention. Most preferably, the antibody of this invention is a monoclonal antibody. For the preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples for such techniques include the hybridoma technique (Köhler and Milstein Nature 256 (1975), 495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor, Immunology Today 4 (1983), 72) and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc. (1985), 77-96). Techniques describing the production of single chain antibodies (e.g., US Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptides as described above. Furthermore, transgenic mice may be used to express humanized antibodies directed against said immunogenic polypeptides. It is in particular preferred that the antibodies/antibody constructs as well as antibody fragments or derivatives to be employed in accordance with this invention or capable to be expressed in a cell. This may, inter alia, be achieved by direct injection of the corresponding proteineous molecules or by injection of nucleic acid molecules encoding the same. Furthermore, gene therapy approaches are envisaged. Accordingly, in context of the present invention, the term "antibody molecule" relates to full immunoglobulin molecules as well as to parts of such immunoglobulin molecules. Furthermore, the term relates, as discussed above, to modified and/or altered antibody molecules, like chimeric and humanized antibodies. The term also relates to monoclonal or polyclonal antibodies as well as to recombinantly or

synthetically generated/synthesized antibodies. The term also relates to intact antibodies as well as to antibody fragments thereof, like, separated light and heavy chains, Fab, Fab/c, Fv, Fab', F(ab')<sub>2</sub>. The term "antibody molecule" also comprises bifunctional antibodies and antibody constructs, like single chain Fvs (scFv) or antibody-fusion proteins. It is also envisaged in context of this invention that the term "antibody" comprises antibody constructs which may be expressed in cells, e.g. antibody constructs which may be transfected and/or transduced via, inter alia, viruses or vectors. It is in particular envisaged that such antibody constructs specifically recognize the polypeptides of the present invention. It is, furthermore, envisaged that said antibody construct is employed in gene therapy approaches.

The present invention relates also to an aptamer specifically binding to a polypeptide according to the invention wherein said aptamer reacts with an epitope of a polypeptide of the present invention. The present invention furthermore relates to an aptamer specifically directed to a corresponding nucleic acid molecule according to the invention.

In accordance with the present invention, the term "aptamer" means nucleic acid molecules that can bind to target molecules. Aptamers commonly comprise RNA, single stranded DNA, modified RNA or modified DNA molecules. The preparation of aptamers is well known in the art and may involve, inter alia, the use of combinatorial RNA libraries to identify binding sides (Gold, Ann. Rev. Biochem. 64 (1995), 763-797).

Furthermore, the present invention relates to a primer or pair of primers capable of specifically amplifying the nucleic acid molecules of the present invention. The term "primer" when used in the present invention means a single-stranded nucleic acid molecule capable of annealing the nucleic acid molecule of the present application and thereby being capable of serving as a starting point for amplification. Said term also comprises oligoribo- or desoxyribonucleotides which are complementary to a region of one of the strands of a nucleic acid molecule of the present invention. According to the present invention the term "pair of primers" means a pair of primers that are with respect to a complementary region of a nucleic acid molecule directed

in the opposite direction towards each other to enable, for example, amplification by polymerase chain reaction (PCR).

The term "amplifying" refers to repeated copying of a specified sequence of nucleotides resulting in an increase in the amount of said specified sequence of nucleotides. and allows the generation of a multitude of identical or essentially identical (i.e. at least 95% more preferred at least 98%, even more preferred at least 99% and most preferred at least 99.5% such as 99.9% identical) nucleic acid molecules or parts thereof. Such methods are well established in the art; see Sambrook et al. "Molecular Cloning, A Laboratory Manual", 2<sup>nd</sup> edition 1989, CSH Press, Cold Spring Harbor. They include polymerase chain reaction (PCR) and modifications thereof, ligase chain reaction (LCR) to name some preferred amplification methods.

When used in the context of primers the term "specifically" means that only the nucleic acid molecules as described herein above are amplified and nucleic acid molecules encoding the wild-type P2X7R ATP-gated receptor as depicted in SEQ ID NO: 1 are not amplified. Thus, a primer according to the invention is preferably a primer which binds to a region of a nucleic acid molecule of the invention which is unique for this molecule and which is not present in the wild-type P2X7R encoding sequence, i.e. the primer binds in a region in which one of the above described mutations occur. In connection with a pair of primers according to the invention it is possible that one of the primers of the pair is specific in the above described meaning or both of the primers of the pair are specific. In both cases, the use of such a pair of primers would allow to specifically amplify a mutant of the invention as described herein-above but not the wild-type P2X7R encoding sequence.

The 3'-OH end of a primer is used by a polymerase to be extended by successive incorporation of nucleotides. The primer or pair of primers of the present invention can be used, for example, in primer extension experiments on template RNA according to methods known by the person skilled in the art. Preferably, the primer or pair of primers of the present invention are used for amplification reactions on template RNA or template DNA, preferably cDNA or genomic DNA. The terms "template DNA" or "template RNA" refers to DNA or RNA molecules or fragments thereof of any source or nucleotide composition, that comprise a target nucleotide sequence as defined above. The primer or pair of primers can also be used for

hybridization experiments as known in the art. Preferably, the primer or pair of primers are used in polymerase chain reactions to amplify sequences corresponding to a sequence of the nucleic acid molecule of the present invention. It is known that the length of a primer results from different parameters (Gillam, Gene 8 (1979), 81-97; Innis, PCR Protocols: A guide to methods and applications, Academic Press, San Diego, USA (1990)). Preferably, the primer should only hybridize or bind to a specific region of a target nucleotide sequence. The length of a primer that statistically hybridizes only to one region of a target nucleotide sequence can be calculated by the following formula:  $(\frac{1}{4})^x$  (whereby x is the length of the primer). For example a hepta- or octanucleotide would be sufficient to bind statistically only once on a sequence of 37 kb. However, it is known that a primer exactly matching to a complementary template strand must be at least 9 base pairs in length, otherwise no stable-double strand can be generated (Goulian, Biochemistry 12 (1973), 2893-2901). It is also envisaged that computer-based algorithms can be used to design primers capable of amplifying the nucleic acid molecules of the invention. Preferably, the primers of the invention are at least 10 nucleotides in length, more preferred at least 12 nucleotides in length, even more preferred at least 15 nucleotides in length, particularly preferred at least 18 nucleotides in length, even more particularly preferred at least 20 nucleotides in length and most preferably at least 25 nucleotides in length. The invention, however, can also be carried out with primers which are shorter or longer.

It is also envisaged that the primer or pair of primers is labeled. The label may, for example, be a radioactive label, such as  $^{32}\text{P}$ ,  $^{33}\text{P}$  or  $^{35}\text{S}$ . In a preferred embodiment of the invention, the label is a non-radioactive label, for example, digoxigenin, biotin and fluorescence dye or a dye.

In another preferred embodiment said primers are selected from the group consisting of SEQ ID NOs: 52 to 111.

In yet another embodiment, the present invention relates to a composition comprising a nucleic acid molecule, a vector, a polypeptide, an antibody, an aptamer and/or a primer or pair of primers of the invention.

The term "composition", as used in accordance with the present invention, relates to compositions which comprise at least one nucleic acid molecule, vector, polypeptide,

an antibody and/or primer or pair of primers of this invention. It may, optionally, comprise further molecules capable of altering the characteristics of the component of the invention thereby, for example, suppressing, blocking, modulating and/or activating their function which have neuroprotective, nootropic and/or antidepressive properties. The composition may be in solid, liquid or gaseous form and may be, inter alia, in the form of (a) powder(s), (a) tablet(s), (a) solution(s) or (an) aerosol(s).

In a preferred embodiment the composition according to the invention is a diagnostic composition, optionally further comprising suitable means for detection. As described above, the present invention is based on the surprising finding that mutations in the P2X7R protein are connected with affective disorders. Thus, this knowledge now allows to diagnose affective disorders in an easy way. The diagnostic composition comprises at least one of the aforementioned compounds of the invention. The diagnostic composition may be used, inter alia, for methods for determining the presence and/or expression of the nucleic acids and/or polypeptides of the invention. This may be effected by detecting, e.g., the presence of a corresponding gene in the genetic material of an individual or the presence of the corresponding mRNA which comprises isolation of DNA or RNA from a cell derived from said individual, contacting the DNA or RNA so obtained with a nucleic acid probe as described above under hybridizing conditions, and detecting the presence of mRNAs hybridized to the probe. Alternatively, the diagnostic composition may also be used for detecting the presence of a nucleic acid molecule of the invention by PCR. Furthermore, polypeptides of the invention can be detected with methods known in the art, which comprise, inter alia, immunological methods, like, RIA, FIA, ELISA, FACS or Western blotting.

Furthermore, the diagnostic composition of the invention may be useful, inter alia, in detecting the prevalence, the onset or the progress of a disease related to the expression of a polypeptide of the invention. Accordingly, the diagnostic composition of the invention may be used, inter alia, for assessing the prevalence, the onset and/or the disease status of affective disorders, as defined herein above. It is also contemplated that the diagnostic composition of the invention may be useful in discriminating (the) stage(s) of a disease.

The diagnostic composition optionally comprises suitable means for detection. The nucleic acid molecule(s), vector(s), host(s), antibody(ies), aptamer(s), polypeptide(s) described above are, for example, suitable for use in immunoassays in which they can be utilized in liquid phase or bound to a solid phase carrier. Examples of well-known carriers include glass, polystyrene, polyvinyl ion, polypropylene, polyethylene, polycarbonate, dextran, nylon, amyloses, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the carrier can be either soluble or insoluble for the purposes of the invention.

Solid phase carriers are known to those in the art and may comprise polystyrene beads, latex beads, magnetic beads, colloid metal particles, glass and/or silicon chips and surfaces, nitrocellulose strips, membranes, sheets, duracytes and the walls of wells of a reaction tray, plastic tubes or other test tubes. Suitable methods of immobilizing nucleic acid molecule(s), vector(s), host(s), antibody(ies), aptamer(s), polypeptide(s), etc. on solid phases include but are not limited to ionic, hydrophobic, covalent interactions or (chemical) crosslinking and the like. Examples of immunoassays which can utilize said compounds of the invention are competitive and non-competitive immunoassays in either a direct or indirect format. Commonly used detection assays can comprise radioisotopic or non-radioisotopic methods. Examples of such immunoassays are the radioimmunoassay (RIA), the sandwich (immunometric assay) and the Northern or Southern blot assay. Furthermore, these detection methods comprise, inter alia, IRMA (Immune Radioimmunometric Assay), EIA (Enzyme Immuno Assay), ELISA (Enzyme Linked Immuno Assay), FIA (Fluorescent Immuno Assay), and CLIA (Chemiluminescent Immune Assay). Furthermore, the diagnostic compounds of the present invention may be employed in techniques like FRET (Fluorescence Resonance Energy Transfer) assays.

Appropriate labels and methods for labeling are known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include inter alia, fluorochromes (like fluorescein, rhodamine, Texas Red, etc.), enzymes (like horse radish peroxidase,  $\beta$ -galactosidase, alkaline phosphatase), radioactive isotopes (like  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{35}\text{S}$  or  $^{125}\text{I}$ ), biotin, digoxigenin, colloidal metals, chemi- or bioluminescent compounds (like dioxetanes, luminol or acridiniums).

A variety of techniques are available for labeling biomolecules, are well known to the person skilled in the art and are considered to be within the scope of the present invention and comprise, inter alia, covalent coupling of enzymes or biotinyl groups, phosphorylations, biotinylations, random priming, nick-translations, tailing (using terminal transferases). Such techniques are, e.g., described in Tijssen, "Practice and theory of enzyme immunoassays", Burden and von Knippenburg (Eds), Volume 15 (1985); "Basic methods in molecular biology", Davis LG, Dibmer MD, Battey Elsevier (1990); Mayer, (Eds) "Immunochemical methods in cell and molecular biology" Academic Press, London (1987); or in the series "Methods in Enzymology", Academic Press, Inc.

Detection methods comprise, but are not limited to, autoradiography, fluorescence microscopy, direct and indirect enzymatic reactions, etc.

Said diagnostic composition may be used for methods for detecting the presence and/or abundance of a nucleic acid molecule of the invention in a biological and/or medical sample and/or for detecting expression of such a nucleic acid molecule (e.g. by determining the mRNA or the expressed polypeptide). Furthermore, said diagnostic composition may also be used in methods of the present invention, inter alia, for the detection of specific antagonists or agonists for P2X7R ATP-gated ion channels (see herein below).

In a further embodiment the present invention provides a method of diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R protein expressed in the cells of said individual is non-functional, shows an altered ATP-gating in comparison to the wild-type P2X7R protein or is over- or under-expressed in comparison to the P2X7R protein level of an unaffected individual.

The term "over- or under-expressed in comparison to the P2X7R protein level" in the context of the present invention means that the P2X7R protein level is higher or lower than the P2X7R level of a healthy individual, i.e. an individual not affected with an affective disorder. The over-expression may result, e.g. from an increased amount of P2X7R mRNA caused by enhanced transcription rates due to increased activity of the RNA-polymerase II. The amount of mRNA may accordingly lead to an increased translation and, thus, to an higher protein level of P2X7R. It may also



be possible that a higher amount of P2X7R protein is caused by increased stability of the protein. An under-expression of P2X7R protein may be caused by low transcription rates of the P2X7R gene and, thus, insufficient amounts of P2X7R mRNA give only rise to a low P2X7R protein amount. Another reason may be that the P2X7R protein is unstable and, thus, is not present in amounts comparable to the wild-type protein level.

The under- or over-expression of P2X7R protein may be determined by methods well-known to the person skilled in the art. These include, but are not limited, to methods for determining the amount of mRNA or the amount and/or activity of the protein. Examples are Northern Blot analysis or immuno based techniques, such as Western Blotting.

"Non-functional" means that the P2X7R protein has lost at least one functional property displayed by the wild-type P2X7R protein as described herein above. Preferably, "non-functional" means that the P2X7R protein does no longer function as a channel. Non-functionality may, e.g., be caused by the fact that one allele occurring in an individual codes for a P2X7R protein which leads to non-functional dimers (dominant negative mutation). Whether a P2X7R protein in an individual is functional or non-functional can be determined by the methods described herein above and in the examples.

The term "altered ATP-gating" means that the respective P2X7R protein reacts in a different way to ATP than the wild-type P2X7R protein. This can be determined as described in the appended examples or as described hereinabove.

In the context of diagnosis, not only the activity of the P2X7R could be of diagnostic value but also the amount of expression. For example, if a polymorphism affects RNA stability or translation efficiency, this could lead to lower expression of the P2X7 protein not only in the hippocampus but also in the blood. Therefore, one could speculate that a lower amount of P2X7 detected by western blot in blood cells could be related to depression.

Another aspect of the present invention is a method for diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R gene

sequence or encoded protein thereof comprises a mutation in comparison to the wild-type P2X7R sequence.

A preferred embodiment of the present invention is a method, wherein a mutation is a mutation in a P2X7R sequence as defined hereinabove and/or a nucleotide replacement or deletion selected from the following Table C indicating in column "Region of P2X7R" the region of the P2X7R genomic nucleotide sequence in which the replacement or deletion occurs, in column "Nucleotide" of Table C the nucleotide which is replaced by another nucleotide or the nucleotides which are deleted and in column "Position in wild-type" of Table C the corresponding position in the nucleotide sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1

**Table C**

<b>Region of P2X7R</b>	<b>Nucleotide</b>	<b>Position in wild-type</b>
5'UTR	T	362
5'UTR	T	532
5'UTR	A	1100
5'UTR	A	1122
5'UTR	C	1171
5'UTR	T	1351
5'UTR	G	1702
5'UTR	T	1731
5'UTR	C	1860
5'UTR	C	2162
5'UTR	C	2238
5'UTR	A	2373
5'UTR	G	2569
5'UTR	G	2702
intron 1	G	3166
intron 1	C	24778
intron 1	C	24830
exon 2	T	24942
exon 3	C	26188
exon 3	A	26308
exon 3	G	26422
intron 4	G	32394
intron 4	T	32434
exon 5	G	32493
exon 5	G	32506

exon 5	C	32507
exon 5	C	32548
intron 5	A	32783
intron 5	T	35309
intron 5	C	35374
intron 5	A	35378
exon 6	G	35438
exon 6	T	35454
intron 6	T	35549
intron 6	G	35641
intron 6	A	35725
intron 6	T	36001
intron 6	A	36064
intron 6	deletion of GTTT	36091 to 36094
intron 6	C	36108
intron 7	C	36374
intron 7	G	36378
intron 7	T	36387
intron 7	G	36398
intron 7	C	37439
intron 7	T	37513
exon 8	C	37604
exon 8	G	37605
exon 8	G	37623
exon 8	C	37633
intron 9	C	47214
exon 11	G	47383
exon 11	C	47411
intron 11	T	47563
intron 12	C	54307
intron 12	G	54308
exon 13	C	54399
exon 13	A	54480
exon 13	C	54523
exon 13	deletion of CCCTGAGAGCCACAGGTGCCT	54562 to 54582
exon 13	A	54588
exon 13	C	54664
exon 13	G	54703
exon 13	A	54804
exon 13	G	54834
exon 13	G	54847
3'UTR	G	54925
3'UTR	C	55169
3'UTR	A	55170
3'UTR	A	55171
3'UTR	C	55917

As indicated hereinabove, if the respective nucleotide which is replaced by another nucleotide is a purine base, it is preferred to be replaced by another purine base. If it is a pyrimidine base, it is preferred to be replaced by another pyrimidine base. It is also preferred that a purine base is replaced by a pyrimidine base and that a pyrimidine base is replaced by a purine base. Most preferably, the nucleotides indicated in Table C are replaced by the nucleotides indicated at the respective position in Table 12 hereinbelow (see Example 3).

In a preferred embodiment the present invention relates to diagnostic composition designed for use in a method in which the occurrence of the mutation in the ATP-gated ion channel P2X7R gene is determined by PCR, immunological methods and/or electrophysiological methods as described herein below and in the appended Examples. Additionally, it is possible to determine the occurrence of a mutation in the ATP-gated ion channel P2X7R as described hereinabove.

In yet another aspect the present invention relates to the use of a nucleic acid molecule, a vector, a polypeptide, an antibody, aptamer and/or a primer or pair of primers of the present invention for the preparation of a diagnostic composition for the detection of an affective disorder.

It is also envisaged that the present invention relates to methods of diagnosing an affective disorder of an individual comprising:

- (a) isolating DNA from cells obtained from an individual;
- (b) determining all or part of the nucleotide composition of the P2X7R gene; and
- (c) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism, mutation or allelic variation.

The term "gene" means a nucleotide sequence associated with the production of a protein, including promoter sequences, enhancer sequences, intron sequences, exon sequences, coding regions, 5' untranslated region (5'UTR), 3' untranslated region (3'UTR), and splice variants.

In a preferred embodiment of the described method the individual is a mammal and more preferably human. Moreover, the cells are preferably derived from skin, blood, urine or cerebral spinal fluid.

The method of the present invention allows for the diagnosis of an affective disorder according to the composition of a genetic marker corresponding to the P2X7R gene. As is demonstrated by the appended examples, polymorphisms in the P2X7R are genetically linked to patients suffering from an affective disorder.

In accordance with this embodiment of the present invention, the diagnosis of an affective disorder can, e.g., be effected by isolating cells from an individual, and isolating the genomic DNA of said cells. Such cells can be collected from body fluids, skin, hair, biopsies and other sources. Collection and analysis of cells from bodily fluids such as blood, urine and cerebrospinal fluid is well known to the art; see for example, Rodak, "Haematology: Clinical Principles & Applications" second ed., WB Saunders Co, 2002; Brunzel, "Fundamentals of Urine and Body Fluids Analysis", WB Saunders Co, 1994; Herndon and Brumback (Ed.), "Cerebrospinal Fluid", Kluwer Academic Pub., 1989. In addition, methods for DNA isolation are well described in the art; see, for example, Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3<sup>rd</sup> edition, Cold Spring Harbor Laboratory, 2001.

Once DNA has been isolated, various oligonucleotide primers spanning the P2X7R locus may be designed in order to amplify the genetic material by Polymerase Chain Reaction (PCR). Conventional methods for designing, synthesizing, producing said oligonucleotide primers and performing PCR amplification may be found in standard textbooks, see, for example Agrawal (Ed.), "Protocols for Oligonucleotides and Analogs: Synthesis and Properties (Methods in Molecular Biology, 20)", Humana Press, 1993; Innis et al. (Ed.), "PCR Applications: Protocols for Functional Genomics", Academic Press, 1999; Chen and Janes (Ed.), "PCR Cloning Protocols: From Molecular Cloning to Genetic", 2<sup>nd</sup> edition, Humana Press, 2002. Primers for the detection of P2X7R polymorphisms are also given in, but not limited to, SEQ ID NO: 52 to SEQ ID NO: 111. Once DNA has been amplified, nucleotide structure can be analysed by sequencing methods and compared to normal P2X7R DNA.

Sequencing may be performed manually by any molecular biologist of ordinary skills or by an automated sequencing apparatus. These procedures are common in the art; see, for example, Adams et al. (Ed.), "Automated DNA Sequencing and Analysis", Academic Press, 1994; Alphey, "DNA Sequencing: From Experimental Methods to Bioinformatics", Springer Verlag Publishing, 1997.

Detection and analysis of polymorphisms in P2X7R can also be performed using amplification refractory mutation system (ARMSTM), amplification refractory mutation system linear extension (ALEXTM), single-strand conformation polymorphism (SSCP), heteroduplex analysis, PCR-SSCP, fluorescent SSCP in an automated DNA sequencer, denaturing gradient gel electrophoresis, RNase protection assays, detection of mutations by sequence specific oligonucleotide hybridization, chemical cleavage methods, enzyme mismatch cleavage methods, cleavage fragment length methods, allele-specific oligonucleotide hybridization on DNA chips, and other such methods known in the art, see, for example Nollau et al, Clin. Chem. 43 (1997), 1114-1128; Burczak and Mardis (Ed.), "Polymorphism Detection & Analysis Techniques", Eaton Pub Co, 2000; Cotton et al. (Ed.), "Mutation Detection: A Practical Approach", Irl Press, 1998; Taylor (Ed.), "Laboratory Methods for the Detection of Mutations and Polymorphisms in DNA", CRC Press, 1997.

The present invention also relates to a method of diagnosing an affective disorder in an individual comprising:

- (a) isolating RNA from cells obtained from an individual;
- (b) converting the RNA into cDNA;
- (c) determining all or part of the nucleotide composition of the cDNA so obtained; and
- (c) analyzing said nucleotide composition for the presence of one or more polymorphism(s) or allelic variation.

---

With respect to the preferred embodiments the same applies as already described above.

Detection and analysis of polymorphisms in the P2X7R RNA can be performed according to the methods described above.

The present invention also relates to a method for diagnosing an affective disorder in an individual comprising:

- (a) isolating RNA or proteins from cells obtained from an individual;
- (b) determining the levels of P2X7R RNA or protein; and
- (c) comparing the levels of P2X7R RNA or protein with the corresponding levels from a normal individual not afflicted with an affective disorder.

With respect to the preferred embodiments the same applies as already described above.

As is demonstrated by the appended examples, a relationship exists between the expression, or protein level of P2X7R and an affective disorder. This and other embodiments of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

According to another aspect on the invention, there is provided a polynucleotide comprising at least 20 bases of the human P2X7R gene and comprising a mutation or polymorphism selected from any of the following:

**Table 1: Novel polymorphisms in the human P2X7R**

<b>Region in P2X7</b>	<b>Polymorphism</b>	<b>Protein Modification</b>
5'UTR	362 T-C	
5'UTR	532 T-G	
5'UTR	1100 A-G	
5'UTR	1122 A-G	
5'UTR	1171 C-G	
5'UTR	1702 G-A	
Intron01	3166 G-C	
Intron01	24778 C-T	
Intron01	24830 6C-T	
Exon03	26188 C-T	Arg117Trp

Intron03	26308 A-G	
Intron03	26422 G-A	
Intron04	32394 G-A	
Intron04	32434 T-C	
Exon05	32493 G-A	Gly150Arg
Exon05	32548 C-T	Silent Cys168
Intron05	32783 A-C	
Exon06	35438 G-A	Glu186Lys
Exon06	35454 T-C	Leu191Pro
Intron06	35641 G-C	
Intron06	35725 A-C	
Intron06	36001 T-G	
Intron07	36378 G-A	
Intron07	36387 T-A	
Intron07	36398 G-C	
Exon08	37604 C-T	Arg270Cys
Exon08	37633 C-T	Silent Asp279
Intron09	47214 C-T	
Intron11	47563 T-C	
Intron12	54307 C-T	
Intron12	54308 G-A	
Exon13	54562-54582 deletion of CCCTGAGA GCCACAGGTGCCT	deletion of 7aa 488 to 494 (PESHRCL)
Exon13	54804 A-T	Ile568Asn
Exon13	54834 G-A	Arg578Gln
3'UTR	55169 C-A	
3'UTR	55170 A-C	
3'UTR	55171 A-C	
3'UTR	55917 C-T	
3'UTR	54925 G-A	

The polymorphism describes the position and the variation observed. The position and numbering of the polymorphism corresponds to the human P2X7R gene as defined in SEQ ID No 1. Primers used for SNP amplification and sequencing are shown in Table 1a and listed in SEQ ID NO: 52 to SEQ ID NO: 111.



**Table 1a. Primer sequences for SNP amplification and sequencing**

<b>Primer Name</b>	<b>Orientation</b>	<b>Sequence</b>	<b>Begin</b>	<b>End</b>
P2RX7_01.for	Sense	cgtaggacttggcgcttct	2785	2803
P2RX7_01.rev	Anti sense	gagcacgtctcagattcgaaa	3224	3244
P2RX7_02.for	Sense	ccatgaggcaggtatgactattc	24665	24687
P2RX7_02.rev	Antisense	ctcctggatctcaccagtt	25168	25187
P2RX7_03.for	Sense	ctcgtccagctttgatattaagc	25966	25988
P2RX7_03.rev	Antisense	ggccctagtgctagaaccaga	26426	26447
P2RX7_04.for	Sense	attcatccgtcagtgccc	30794	30811
P2RX7_04.rev	Antisense	gccatgtgaattttctaccgat	31277	31298
P2RX7_05.for	Sense	ttcgttggtggttaggatggg	32314	32333
P2RX7_05.rev	Antisense	caaggatgctcagggtagtagc	32805	32826
P2RX7_06.for	Sense	cactaggtttgctgtatccatttct	35277	35301
P2RX7_06.rev	Antisense	gcaactgtgtgagagcttgg	35731	35750
P2RX7_07.for	Sense	tcaaccttggtccagtgtg	35950	35968
P2RX7_07.rev	Antisense	caccaagtagctctcactcataagg	36424	36448
P2RX7_08.for	Sense	caataacacttggtgcgagttaggt	37380	37403
P2RX7_08.rev	Antisense	catcttggtgccttggaacc	37750	37770
P2RX7_09.for	Sense	gtgagtggtaaatcctgctactgc	45321	45343
P2RX7_09.rev	Antisense	aggccactcctgtactcg	45743	45761
P2RX7_10_11.for	Sense	ccaagtcacagcatgaggc	47119	47137
P2RX7_10_11.rev	Antisense	accagcgcagctatccac	47632	47649
P2RX7_12.for	Sense	aagcatgggggttccatttc	50252	50268
P2RX7_12.rev	Antisense	gcataaaagggaactcctgctagta	50691	50714
P2RX7_13a.for	Sense	gcttacagaaacacatgcatgg	54232	54252
P2RX7_13a.rev	Antisense	gcacctgtaggcacagtgc	54739	54757
P2RX7_13b.for	Sense	atcaccacctcagagctgttc	54620	54640
P2RX7_13b.rev	Antisense	gttaacatggctactgcagcc	55203	55223
P2XR7_13d.for	Sense	gcttagaaaggaggcgactcc	54484	54504
P2XR7_Pro13.for	Sense	ttgtgacatttgcaaggctgcc	2617	2638
P2XR7_Pro7.rev	Antisense	tctgaagctctgctcctgag	1955	1974
P2XR7_Pro8.rev	Antisense	ctcaccttctggcttccagt	1611	1630
P2XR7_Pro9.for	Sense	cttaccactcccaggactaa	1496	1515
P2XR7_Pro10.for	Sense	gtctgcctgttcaactgcat	1149	1168
P2XR7_Pro1.for	Sense	cagagaccttcagaaacttcg	1841	1861
P2XR7_Pro2.rev	Antisense	agatcaccagggacacagtgc	2261	2280
P2XR7_Pro3.for	Sense	ctcaactccactttcctcgg	2133	2152
P2XR7_Pro4.rev	Antisense	cctttcacttttttggtctcatg	2655	2677
P2XR7_Pro5.for	Sense	gggagaattctgaaaatgccc	2691	2711
P2XR7_Pro6.rev	Antisense	ggaccagagctctactcttc	2951	2970
P2XR7_Pro11.for	Sense	aggcatagatcgacctgcc	2296	2315
P2XR7_Pro12.rev	Antisense	aagaagcgccaagtcctacg	2785	2804
P2XR7_Pro14.for	Sense	gcaatccagactgaagttgac	2051	2071
P2XR7_Pro15.rev	Antisense	actctgggtctgcagttgggtg	2428	2447
P2XR7_Pro21.for	Sense	cctttaaaatcagagaccttcaga	1831	1854
P2XR7_Pro22.for	Sense	gcccatacctctgaacaccat	2708	2727
P2XR7_3UTR10.for	Sense	cccttggaactcttgctatcg	55804	55824
P2XR7_3UTR1.for	Sense	ggcagtacagtggttcaaga	54858	54878
P2XR7_3UTR2.rev	Antisense	gtgggacagtttgctgtgcct	55150	55170

P2XR7_3UTR3.for	Sense	gagtccttaccatagcagg	55183	55202
P2XR7_3UTR4.rev	Antisense	gtcaaagaatttgtggccacc	55643	55663
P2XR7_3UTR5.for	Sense	catgaactgtcttttaatgtgtaaag	55515	55540
P2XR7_3UTR6.rev	Antisense	gagatacggtttcaccatgttg	55955	55976
P2XR7_3UTR7.for	Sense	aattagctgggcatggtgcg	55992	56011
P2XR7_3UTR8.rev	Antisense	ttgagatggagtctcgctctg	56122	56140
P2XR7_3UTR9.rev	Antisense	cactgtccacgtgactgctt	56208	56227
P2RX7_11.For	Sense	tcctacttcggtctggttaagagatt	47281	47305
P2RX7_11.Rev	Antisense	gggcctaattttcgtgcat	47591	47609
P2RX7_13G.For	Sense	aagaacctagaacctgagggctt	54333	54355
P2RX7_13G.Rev	Antisense	ttgagatgggaggcagctt	54541	54559
P2RX7_13H.For	Sense	ttcggctcccaggacat	54773	54789
P2RX7_13H.Rev	Antisense	cacagagctttgcaggtgaa	55248	55267

Another aspect of the present invention is in the form of a diagnostic kit for affective disorders comprising a specific oligonucleotide probe, or primer corresponding to P2XR7 polymorphisms. The diagnostic kit may comprise appropriate packaging and instructions for the use in the method of the invention. Said kit may further comprise appropriate buffer, and enzymes such as reverse transcriptase, and thermostable polymerases.

In a preferred embodiment of the invention, diagnosis can be performed on a mouse, rat or human. The invention is generally applied in vitro, e.g. using cells or other material obtained from an individual. However, it can also be applied on a living individual, or post mortem.

In accordance with the embodiments of the present invention, diagnosis of an affective disorder may be followed by prescription, or administration of an antidepressant drug. Administration and dosage of antidepressive drugs can vary between patients and are well known in the medical art, see, for example Benkert and Hippus, "Kompendium der Psychiatrischen Pharmakotherapie", Springer Verlag Publishing, 2000; Albers, "Handbook of Psychiatric Drugs: 2001-2002 Edition", Current Clinical Strategies Publishing, 2000. Preferred examples include between 5 mg and 80 mg per day, preferably 20 mg, fluoxetine; between 5 mg and 50 mg per day, preferably 20 mg, paroxetine; between 5 mg and 200 mg per day, preferably 50 mg, sertraline; between 5 mg and 300 mg per day, preferably 100 mg, fluvoxamine; between 5 mg and 100 mg per day, preferably 30 mg, mirtazapine; between 4 mg

and 50 mg, preferably 8 mg, reboxetine; between 5 mg and 600 mg per day, preferably 200 mg, nefazodone; between 450 mg and 1800 mg per day, preferably 900 mg, lithium carbonate.

The P2X7R protein is also useful for monitoring the efficacy and/or dosing of a drug or the likelihood of a patient to respond to a drug. Thus, in yet another embodiment the invention relates to a method for, monitoring the efficacy and/or dosing of a drug, e.g. an antidepressive drug, and/or the likelihood of a patient to respond to said drug which comprises determining the level of expression and/or activity of the P2X7R protein in a patient before and after administration of the respective drug. As presented in the examples below, treatment with an antidepressive drug results in an upregulation in P2X7R activity. In humans, P2X7R activity can be monitored by Positron Emission Tomography (PET) or Single Photon Emission Computerised Tomography (SPECT) using a radiolabelled ligand tracer for P2X7R. Examples of P2X7R ligands can be, but are not limited to, ATP, an antagonist binding P2X7R, an agonist binding P2X7R, or a small polynucleotide comprising at least 20 bases of the human P2X7R gene. A modulation of P2X7R activity, membrane distribution or expression levels would reflect the activity and potency of the antidepressive drug. Methods and techniques required for PET analysis are well known in the art, see, for example Paans and Vaalburg, Curr. Pharmac. Design 6 (2000), 1583-1591; van Waarde, Curr. Pharmac. Design. 6 (2000), 1593-1610; Paans et al, Methods 27 (2002), 195-207; Passchier et al., Methods 27 (2002), 278-286; Laruelle et al., Methods 27 (2002), 287-299.

In accordance with the present invention by the term "sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source containing polynucleotides or polypeptides or portions thereof. As indicated, biological samples include body fluids (such as blood, sera, plasma, urine, synovial fluid and spinal fluid) and tissue sources found to express the polynucleotides of the present invention. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. A biological sample which includes genomic DNA, mRNA or proteins is preferred as a source.

As described herein above, mutations of the P2X7R encoding gene can occur on DNA level or on mRNA level and may result in an altered expression of P2X7R or in the expression of P2X7R ATP-gated ion channels which show either an altered function or no function when compared to the wild-type P2X7R ATP-gated ion channel as described herein. Thus, various methods on DNA level, RNA level or protein level exist for determining whether the ATP-gated ion channel P2X7R gene shows a mutation as described herein above. Consequently, mRNA, cDNA, DNA and genomic DNA are the preferred nucleic acid molecules to be used in the below mentioned methods. Also polypeptides or fragments thereof are preferred if a mutation in the P2X7R ATP-gated ion channel protein as described herein is to be determined.

Preferably, a point mutation leading to the replacement of an amino acid residue at the positions as indicated in Table 1 of the corresponding wild-type P2X7R amino acid sequence depicted in SEQ ID NO: 3 or 4 by another amino acid can be determined by PCR. Said PCR is followed by a restriction fragment length polymorphism (RFLP) analysis if due to the point mutation a recognition site for a restriction endonuclease is generated which is not present in the wild-type nucleotide sequence or a recognition site for a restriction enzyme is created which does not occur in the wild-type P2X7R. More preferably said mutation can be determined by PCR using primers and conditions that allow only an amplification of the wild-type nucleotide sequence encoding the corresponding wild-type amino acid at the respective position, but not of the nucleotide sequence of a nucleic acid molecule encoding a different amino acid residue at the corresponding position. It is even more preferred that PCR is performed to determine a mutation using primers and conditions that allow no amplification if the wild-type nucleotide sequence is present, but only if another amino acid residue is encoded at the respective position. Particularly preferred is a method using PCR and primers under conditions that allow amplification of a fragment comprising at least the nucleotide residues encoding the amino acid residue corresponding to positions of SEQ ID NO: 1.

Said PCR is followed by e.g., sequencing and/or single strand conformation analysis (SSCA). Said fragment is preferably of at least 25 nucleotides in length, more preferred of at least 50 nucleotide in length, even more preferred of at least 75

nucleotides in length, particularly preferred of at least 100 nucleotides in length, more particularly preferred of at least 200 nucleotides in length, also more particularly preferred at least 250 nucleotides in length, even more particularly preferred at least 300 nucleotides in length and most particularly preferred at least 600 nucleotides in length. Said primers are preferably of at least 12 nucleotides in length, more preferred of at least 15 nucleotides in length, even more preferred of at least 18 nucleotides in length and most preferred of at least 21 nucleotides in length as depicted in SEQ ID NOs: 52 to 111. The temperature for annealing said primers is preferably at least 50°C, more preferred at least 55°C and most preferred at least 58°C. The temperature for denaturation is preferably at least 95°C for preferably at least 10 sec, more preferably at least 20 sec, even more preferred at least 30 sec and most preferred at least 60 sec. However, depending on the length and the G-C content of the nucleic acid sequence to be amplified the temperature for denaturation may be shorter or longer. The temperature for extension of the annealed primers is preferably at least 10 sec, more preferably at least 20 sec, even more preferred at least 30 sec and most preferred at least 60 sec. A PCR reaction comprising the aforementioned conditions is exemplified in the Examples herein below. The subsequent sequencing and/or SSCA is carried out as known in the art. Preferably, the PCR fragments are separated on a 10% polyacrylamide gel at 4°C or also preferred at room temperature. PCR fragments showing a SSCA band shift are amplified with the primers under conditions as mentioned above and are subsequently sequenced. Alternatively, it is also possible to directly sequence genomic DNA in order to determine whether a mutation in the CLCN2 gene has occurred. A direct genomic sequencing approach is, for example, demonstrated for baker's yeast in Horecka, Yeast 16 (2000), 967-970.

Preferably, a deletion is determined by using hybridization techniques as known in the art. In particular, a primer is designed as mentioned herein above that is capable to only hybridize to wild-type genomic DNA as depicted in SEQ ID NO: 1 but not to a nucleotide sequence comprising a deletion of a fragment between nucleotides 54562 and 54582 of SEQ ID NO: 1. Also preferred is the method of fluorescent in situ hybridization (FISH) for determining on whole chromosomes, in particular on chromosome 12q23-q24 that said chromosome has the above mentioned deletion. Even more preferred is that a deletion of nucleotide residues as described herein

may be determined by using PCR, wherein one primer of a pair of primers is located within the region of genomic DNA comprising said deletion. Preferably, said deletion is between nucleotide positions 54562 and 54582 as depicted in SEQ ID NO: 1. Thus, under the appropriate conditions no PCR fragment will result if the genomic DNA comprises said deletion. It is particularly preferred that PCR using primers which are located upstream or downstream of the deletion is performed to determine said deletion. Under appropriate conditions as mentioned herein above, both a fragment of genomic DNA of the wild-type nucleotide sequence as set forth in SEQ ID NO: 1 and a fragment of the nucleotide sequence comprising a deletion of preferably the nucleotides between positions 54562 and 54582 as depicted in SEQ ID NO: 1 will be amplified.

It is also possible to determine the above-described P2X7R mutations on the protein level. Some of the mutations described above lead to shortened versions of the P2X7R protein. Thus, it is conceivable to determine the occurrence of these mutations by determining the length or molecular weight of the P2X7R protein expressed in an individual, e.g. by SDS PAGE.

It is also possible to determine the mutations of the P2X7R ATP-gated channel as described herein by using the antibodies of the present invention. Said antibodies specific for said mutations of P2X7R proteins will be determined by assay techniques such as radioimmunoassays, competitive-binding assays, Western blot analysis and ELISA assay. Also preferred are classical immunohistological methods.

The finding, described in the present invention, that certain mutations in the P2X7R encoding gene and/or the corresponding protein are connected with affective disorder is indicative that the non- or dysfunction of the P2X7R protein is responsible for various forms of affective disorders. Thus, the finding of these mutations not only allows the diagnosis of affective disorders by determining whether the above-described mutations occur in an individual. It also allows to develop a treatment of ~~affective disorders which has been diagnosed to be the result of a mutation in the~~ P2X7R encoding gene. Such a treatment can, e.g., comprise the introduction of a nucleic acid molecule encoding a non-functional or functional wild-type P2X7R protein thereby restoring in said individual the P2X7R activity or the activation or

repression of (a) P2X7R gene(s) in vivo. The term "activation or repression" in this context means that the expression of the gene is either enhanced (activation) or reduced (repression). An enhancement of expression can, e.g., be achieved by increasing the efficiency of transcription initiation, for example, by using suitable compounds which have an activating effect on transcription. Alternatively, an enhancement can be achieved by replacing the naturally occurring promoter by a more efficient promoter.

A repression may be achieved by suppressing expression of the gene, e.g., by specifically suppressing transcription from the respective promoter by suitable compounds or by rendering the promoter less efficient or non-functional.

In another embodiment the present invention also relates to a pharmaceutical composition. In accordance with the present invention the term "pharmaceutical composition" relates to a composition comprising a nucleic acid molecule comprising a nucleotide sequence which encodes an ATP-gated ion channel P2X7R and which is selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as depicted in SEQ ID NO: 3 or 4;
- (b) a nucleotide sequence comprising the nucleotide sequence as depicted in SEQ ID NO: 1 or SEQ ID NO: 2;
- (c) a nucleotide sequence which hybridizes to the nucleotide sequence of (a) or (b); and
- (d) a nucleotide sequence which is degenerated as a result of the genetic code to the nucleotide sequence of (c).

Such pharmaceutical compositions comprise a therapeutically effective amount of a nucleic acid molecule encoding a functional P2X7R protein and, optionally, a pharmaceutically acceptable carrier. The pharmaceutical composition may be administered with a physiologically acceptable carrier to a patient, as described herein. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency or other generally recognized pharmacopoeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered.

Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium ion, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the aforementioned compounds, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In another preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilised powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition



is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The pharmaceutical composition of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

In vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems. Preferably, the pharmaceutical composition is administered directly or in combination with an adjuvant.

The pharmaceutical composition is preferably designed for the application in gene therapy. The technique of gene therapy has already been described above in connection with the nucleic acid molecules of the invention and all what has been said there also applies in connection with the pharmaceutical composition. For example, the nucleic acid molecule in the pharmaceutical composition is preferably in a form which allows its introduction, expression and/or stable integration into cells of an individual to be treated.

For gene therapy, various viral vectors which can be utilized, for example, adenovirus, herpes virus, vaccinia, or, preferably, an RNA virus such as a retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can also incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a P2X7R sequence of interest encoding a functional P2X7R protein into the

viral vector, along with another gene which encodes, for example, the ligand for a receptor on a specific target cell, for example, the vector is now target specific.

Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a sugar, a glycolipid, or a protein. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector containing the inserted polynucleotide sequence.

Since recombinant retroviruses are preferably defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include, but are not limited to w2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced. Alternatively, NIH 3T3 or other tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes gag, pol and env, by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral vector into the culture medium. Another targeted delivery system for P2X7R polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0  $\mu\text{m}$  can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraleigh, et al., Trends Biochem. Sci.,

6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, et al., Biotechniques, 6:682, 1988). The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations. Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebrosides, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine. The targeting of liposomes can be classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries.

In another aspect the present invention relates to a method of treating an affective disorder comprising administering a therapeutically effective amount of the pharmaceutical composition comprising a nucleotide sequence encoding a functional ATP-gated ion channel as described herein above to a subject suffering from said disorder.

In the context of the present invention the term "subject" means an individual in need of a treatment of an affective disorder. Preferably, the subject is a vertebrate, even more preferred a mammal, particularly preferred a human.

The term "administered" means administration of a therapeutically effective dose of the aforementioned nucleic acid molecule encoding a functional P2X7R protein to an individual. By "therapeutically effective amount" is meant a dose that produces the effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is known in the art and described above, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art.

The methods are applicable to both human therapy and veterinary applications. The compounds described herein having the desired therapeutic activity may be administered in a physiologically acceptable carrier to a patient, as described herein. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways as discussed below. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt %. The agents maybe administered alone or in combination with other treatments.

The administration of the pharmaceutical composition can be done in a variety of ways as discussed above, including, but not limited to, orally, subcutaneously, intravenously, intra-arterial, intranodal, intramedullary, intrathecal, intraventricular, intranasally, intrabronchial, transdermally, intranodally, intrarectally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the candidate agents may be directly applied as a solution dry spray.

The attending physician and clinical factors will determine the dosage regimen. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. A typical dose can be, for example, in the range of 0.001 to 1000  $\mu$ g; however, doses below or above this exemplary range are envisioned, especially considering the aforementioned factors.

The dosages are preferably given once a week, however, during progression of the treatment the dosages can be given in much longer time intervals and in need can be given in much shorter time intervals, e.g., daily. In a preferred case the immune response is monitored using herein described methods and further methods known to those skilled in the art and dosages are optimized, e.g., in time, amount and/or composition. Dosages will vary but a preferred dosage for intravenous administration of DNA is from approximately  $10^6$  to  $10^{12}$  copies of the DNA molecule. If the regimen is a continuous infusion, it should also be in the range of 1  $\mu$ g to 10 mg units per kilogram of body weight per minute, respectively. Progress can be monitored by periodic assessment. The pharmaceutical composition of the invention may be administered locally or systemically. Administration will preferably be parenterally, e.g., intravenously. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium ion solution, Ringer's dextrose, dextrose and sodium ion, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

It is also envisaged that the pharmaceutical compositions are employed in co-therapy approaches, i.e. in co-administration with other medicaments or drugs, for example other drugs for preventing, treating or ameliorating affective disorders.

Another aspect of the present invention is a pharmaceutical composition comprising a compound, the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells or comprising a nucleic acid molecule the expression of which in cells or the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells. Said pharmaceutical

composition may be useful for treating individuals having an increased amount of the P2X7R protein or expression level as described hereinabove.

Preferably, the above-mentioned pharmaceutical composition is an antisense nucleic acid, a ribozyme, a co-suppressive nucleic acid, iRNA or siRNA.

An siRNA approach is, for example, disclosed in Elbashir ((2001), Nature 411, 494-498)). It is also envisaged in accordance with this invention that for example short hairpin RNAs (shRNAs) are employed in accordance with this invention as pharmaceutical composition. The shRNA approach for gene silencing is well known in the art and may comprise the use of st (small temporal) RNAs; see, inter alia, Paddison (2002) Genes Dev. 16, 948-958.

As mentioned above, approaches for gene silencing are known in the art and comprise "RNA"-approaches like RNAi or siRNA. Successful use of such approaches has been shown in Paddison (2002) loc. cit., Elbashir (2002) Methods 26, 199-213; Novina (2002) Mat. Med. June 3, 2002; Donze (2002) Nucl. Acids Res. 30, e46; Paul (2002) Nat. Biotech 20, 505-508; Lee (2002) Nat. Biotech. 20, 500-505; Miyagishi (2002) Nat. Biotech. 20, 497-500; Yu (2002) PNAS 99, 6047-6052 or Brummelkamp (2002), Science 296, 550-553. These approaches may be vector-based, e.g. the pSUPER vector, or RNA polIII vectors may be employed as illustrated, inter alia, in Yu (2002) loc. cit.; Miyagishi (2002) loc. cit. or Brummelkamp (2002) loc. cit.

A compound which leads to a reduction of the expression of the P2X7R gene may, e.g., be a compound which acts on the regulatory region of the gene and thereby reduces the level of transcription. Such compounds can be identified by methods as described herein below.

The invention also relates to the use of a nucleic acid molecule encoding a functional P2X7R protein as described herein above in connection with the pharmaceutical composition for the preparation of a pharmaceutical composition for treating an affective disorder.

Furthermore, the present invention relates to a method of treating an affective disorder comprising administering a therapeutically effective amount of the nucleic

acid molecule according to the invention or a therapeutically effective amount of the corresponding encoded polypeptide to a subject suffering from said disorder.

In another preferred embodiment the present invention relates to a pharmaceutical composition comprising, inter alia, the polynucleotides according to the present invention, i.e. polynucleotides having mutations and/or deletions as described hereinabove. Such pharmaceutical compositions may, e.g., be useful for treating individuals having an increased or amount of the P2X7R protein or having a P2X7R protein showing an increased activity which can be determined as described hereinabove. It is envisaged that, e.g. a non-functional P2X7R protein comprised by said pharmaceutical composition is incorporated in a P2X7R complex which naturally exists in cells as described hereinabove. It is also envisaged that the above-described techniques for gene therapy can be used for treating an individual with the nucleic acid molecules of the present invention, *mutatis mutandis*.

With respect to the possible modes of administration and preferred embodiments the same applies as has been set forth above.

Additionally, the present invention also envisages the use of the nucleic acid molecules, the vectors, the polypeptides, the antibody and/or the aptamer according to the invention for the preparation of a pharmaceutical composition for the treatment of an affective disorder.

A further aspect of the present invention is the use of a modulator of P2X7R activity or expression for the preparation of a pharmaceutical composition for treating an affective disorder. In the context of the present invention the term "modulator" means (a) compound(s), a complex of compounds, (a) substance(s) or complex of substances which can modify, i.e. modulate the activity of P2X7R or the expression of P2X7R. The modulation can, for example, occur at the protein level. Particularly, the P2X7R protein may interfere with the modulator such that it is either more active or less active. The modulation can also occur on nucleic acid level. Namely, the gene is transcribed more frequently or less frequently giving rise to more or less protein. Modulation can also influence RNA or protein stability. Since it was


surprisingly found that agonists of P2X7R improve the symptoms of mice selected for anxiety and depressive behaviour, the modulator of P2X7R activity used for the preparation of a pharmaceutical composition for treating an affective disorder is preferably an agonist. The term "agonist" means an agent or a compound that can interact with a receptor and initiate a physiological or a pharmacological response characteristic of that receptor. Examples of P2X7R agonist include but are not restricted to ATP, ATP-4, and BzATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate ( $C_{24}H_{24}N_5O_{15}P_3$ )). Particularly preferred, said agonist used to treat depression or anxiety is BzATP as demonstrated in Example 9 hereinbelow.

Although it was reported that activation of P2X7R could induce apoptosis and cell death in vitro (Di Virgilio et al., Cell Death Differ. 5 (1988), 191-199; Virginio et al., J. Physiol. 519 (1999), 335-346) the present application demonstrates in Example 9 hereinbelow that treatment of the brain of mice selected for anxiety and depressive behaviour with BzATP revealed no significant difference in the numbers of apoptotic cells between control mice and mice treated with BzATP. This result indicates that activation of P2X7R did not result in cerebral cell death in vivo which, thus, renders BzATP to be a candidate drug for treatment of affective disorders.

It is furthermore envisaged that said modulator is selected from the group consisting of piperidine and piperazine derivatives, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate and 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate as, for example, described in WO 99/29660, WO 99/29661; WO 99/296896; WO 00/61569; WO 01/42194; WO 01/44170; WO 01/44213; WO 00/71529; WO 01/46200. The following compounds illustrate compounds which are also preferred to be used as modulators of P2X7R activity.


A compound of general formula:






where A is phenyl or a 5- or 6-membered heterocyclic ring containing one or two heteroatoms selected from O, N or S; and optionally substituted by  $C_{1-6}$ alkyl, halogen, nitro, amino, alkylamino,  $CF_3$ ,  $SO_2Me$ ,  $NHSO_2Me$  or cyano; B is  $C=O$ ,  $NH$  or  $SO_2$ ; X is  $C=O$ ,  $CH(Me)$ , O or  $(CH_2)_p$  where p is 0 or 1; Y is O,  $CH_2$ ,  $NH$  or S; Z is  $C=O$  or  $SO_2$ , provided that when Z is  $C=O$ , then Y is O,  $CH_2$  or S; R is hydrogen or  $C_{1-6}$ alkyl;  $R^1$  is hydrogen, halogen;  $R^2$  is phenyl optionally substituted by  $CO_2H$ ,  $CO_2$ alkyl,  $CONH_2$  or  $R^2$  is OH,  $NHR^3$ ,  $NHCH(R^4)(CHR^5)_nR^6$ ,  $NH-R^7-R^8$ ,  $SO_2NH$ alkyl,  $NHCO$ alkyl,  $NHSO_2$ alkyl, morpholine,  $NR^9R^{10}$ , piperazine substituted by phenyl, alkoxyphenyl, pyridyl or fluorophenyl; n is 0, 1 or 2;  $R^3$  is hydrogen, a bi- or tricyclic saturated ring system optionally containing a nitrogen atom, piperidinyl, alkylpyrrolidine, ethynylcyclohexyl, a 5-membered aromatic ring containing 2 or 3 heteroatoms,  $C_{4-6}$  cycloalkyl optionally substituted by alkyl, cyano or hydroxy, or  $C_{1-8}$  alkyl optionally containing an oxygen atom in the alkyl chain and being optionally substituted by one or more substituents selected from ethynyl, cyano, fluoro, dialkylamino, hydroxy, thioalkyl,  $CO_2R^{11}$  or  $CONH_2$ ;  $R^4$  is hydrogen or alkyl optionally substituted by hydroxy or alkoxy;  $R^5$  is hydrogen or hydroxy;  $R^6$  is  $CO_2R^{11}$ ,  $NHCO_2R^{12}$ ,  $CONH_2$  or a 5 or 6-membered saturated ring containing an oxygen atom, a 5-membered heterocyclic ring containing one or two heteroatoms selected from O, N or S, or phenyl optionally substituted by one or more groups selected from alkyl, hydroxy, amino, alkoxy, or nitro;  $R^6$  is alkyl;  $R^7$  is a cyclopentane ring;  $R^8$  is phenyl;  $R^9$  and  $R^{10}$  are independently hydrogen, benzyl, alkenyl, cycloalkyl, alkyl optionally substituted by hydroxy, alkoxy, cyano, dialkylamino, phenyl, pyridyl or  $CO_2R^{11}$  or  $R^9$  and  $R^{10}$  together form a 5- to 7-membered saturated or partially saturated ring optionally containing a further heteroatom and optionally substituted by one or more groups selected from alkyl (optionally containing an oxygen atom in the chain and optionally substituted 14 by hydroxy),  $CO$ alkyl,  $CO_2R^{11}$ ,  $COR^{13}R^{14}$ ,  $CHO$  or piperidine,  $R^{11}$  is hydrogen or alkyl;  $R^{12}$  is alkyl; and  $R^{13}$  and  $R^{14}$  are independently hydrogen or alkyl, is or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:






wherein A represents a group  $\text{CH}_2$  or an oxygen atom; B represents a hydrogen or halogen atom; D represents a group  $\text{CH}_2$ ,  $\text{OCH}_2$ ,  $\text{NHCH}_2$  or  $\text{CH}_2\text{CH}_2$ ; R represents a phenyl, benzothiazolyl, indolyl, indazolyl, purinyl, pyridyl, pyrimidinyl or thiophenyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or a cyano, carboxyl, hydroxyl, nitro, halo- $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $-\text{N}(\text{R}^1)\text{-C(=O)-R}^2$ ,  $-\text{C(O)-NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{R}^6$ ,  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$ , 3- to 8-membered heterocyclyl,  $\text{C}_3\text{-C}_8\text{-cycloalkyloxy}$ ,  $\text{C}_1\text{-C}_6\text{-alkylcarbonyl}$ , phenoxy, benzyl,  $\text{C}_1\text{-C}_6\text{-alkylthio}$ , phenylthio,  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkylsulphinyl}$  or  $\text{C}_1\text{-C}_6\text{-alkylsulphonyl}$  group, or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_1\text{-C}_6\text{-alkoxy}$  group optionally substituted by one or more substituents independently selected from a halogen atom or an amino, carboxyl, hydroxyl,  $\text{C}_1\text{-C}_6\text{-alkoxy}$ , (di) $\text{C}_1\text{-C}_6\text{-alkylamino}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$ , imidazolyl, morpholinyl, piperidinyl or pyrrolidinyl group;  $\text{R}^1$  represents a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^2$  represents a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group; and  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$  and  $\text{R}^6$  each independently represent a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group; with the provisos that when A is  $\text{CH}_2$ , B is H and D is  $\text{CH}_2$ , then R does not represent a phenyl, ortho-carboxyphenyl, methylphenyl or para-phenoxyphenyl group, and that when A is  $\text{CH}_2$ , D is  $\text{CH}_2$  or  $\text{CH}_2\text{CH}_2$  and R represents a substituted phenyl group, the substituent or substituents present do not comprise, in an ortho position, a  $\text{C}_1\text{-C}_6\text{-alkoxy}$  group substituted by an amino, (di) $\text{C}_1\text{-C}_6\text{-alkylamino}$ , imidazolyl, morpholinyl, piperidinyl or pyrrolidinyl group; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:



(I)


wherein x represents 1 or 2; A represents a group  $\text{CH}_2$  or an oxygen atom; B represents a hydrogen or halogen atom; R represents a phenyl, pyridyl, indolyl, indazolyl, pyrimidinyl or thiophenyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or an amino, cyano, carboxyl, hydroxyl, nitro,  $\text{C}_1\text{-C}_6\text{-alkyl}$ , halo- $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $-\text{N}(\text{R}^1)\text{-C}(=\text{O})\text{-R}^2$ ,  $-\text{C}(\text{O})\text{NR}^3\text{R}^4$ ,  $-\text{NR}^5\text{R}^6$ ,  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$ , 3- to 8-membered heterocyclyl,  $\text{C}_3\text{-C}_8\text{-cycloalkyloxy}$ ,  $\text{C}_1\text{-C}_6\text{-alkylcarbonyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkylsulphinyl}$  or  $\text{C}_1\text{-C}_6\text{-alkylsulphonyl}$  group, or a  $\text{C}_1\text{-C}_6\text{-alkoxy}$ ,  $\text{C}_1\text{-C}_6\text{-alkylamino}$ , phenoxy, benzyl,  $\text{C}_1\text{-C}_6\text{-alkylthio}$  or phenylthio group optionally substituted by one or more substituents independently selected from a 15 halogen atom or an amino, cyano, carboxyl, hydroxyl, nitro, 1-pyrrolidinyl, 1-piperidinyl,  $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxy}$ , (di) $\text{C}_1\text{-C}_6\text{-alkylamino}$ , halo- $\text{C}_1\text{-C}_6\text{-alkyl}$ ,  $\text{C}_1\text{-C}_6\text{-alkoxycarbonyl}$  or one of the following groups:




$\text{R}^1$  represents a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^2$  represents a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^3$  and  $\text{R}^4$  each independently represent a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^5$  represents a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^6$  represents a  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group and, additionally, a  $\text{C}_1\text{-C}_6\text{-alkyl}$  group when  $\text{R}^5$  is not a hydrogen atom;  $\text{R}^7$  represents a hydrogen atom or a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^8$  represents a  $\text{C}_1\text{-C}_6\text{-alkyl}$  or  $\text{C}_3\text{-C}_8\text{-cycloalkyl}$  group;  $\text{R}^9$  represents a hydrogen atom or a hydroxyl group; and  $\text{R}^{10}$  represents a hydrogen atom or a phenyl or imidazolyl

group; with the provisos that R does not represent an unsubstituted pyridyl group when A represents a group  $\text{CH}_2$  and B represents a hydrogen atom, and that when R represents a substituted phenyl, indolyl or indazolyl group, the substituent or substituents present do not comprise an amido, carboxyl, (di) $\text{C}_1\text{-C}_6$ -alkylamido or  $\text{C}_1\text{-C}_6$ -alkoxycarbonyl group in an ortho position; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:




wherein D represents  $\text{CH}_2$  or  $\text{CH}_2\text{CH}_2$ ; E represents  $\text{C}(\text{O})\text{NH}$  or  $\text{NHC}(\text{O})$ ;  $\text{R}^1$  and  $\text{R}^2$  each independently represent a hydrogen or halogen atom, or an amino, nitro,  $\text{C}_1\text{-C}_6$ -alkyl or trifluoromethyl group;  $\text{R}^3$  represents a group of formula:




X represents an oxygen or sulphur atom or a group  $\text{NH}$ ,  $\text{SO}$  or  $\text{SO}_2$ ; Y represents an oxygen or sulphur atom or a group  $\text{NR}^{11}$ ,  $\text{SO}$  or  $\text{SO}_2$ ; Z represents a group  $-\text{OH}$ ,  $-\text{SH}$ ,  $-\text{CO}_2\text{H}$ ,  $\text{C}_1\text{-C}_6$ -alkoxy,  $\text{C}_1\text{-C}_6$ -alkylthio,  $\text{C}_1\text{-C}_6$ -alkylsulphinyl,  $\text{C}_1\text{-C}_6$ -alkylsulphonyl,  $-\text{NR}^6\text{R}^7$ ,  $-\text{C}(\text{O})\text{NR}^8\text{R}^9$ , imidazolyl, 1-methylimidazolyl,  $-\text{N}(\text{R}^{10})\text{C}(\text{O})\text{-C}_1\text{-C}_6\text{-alkyl}$ ,  $\text{C}_1\text{-C}_6$ -alkylcarbonyloxy,  $\text{C}_1\text{-C}_6$ -alkoxycarbonyloxy,  $-\text{OC}(\text{O})\text{NR}^{12}\text{R}^{13}$ ,  $-\text{OCH}_2\text{OC}(\text{O})\text{R}^{14}$ ,  $\text{OCHOC}(\text{O})\text{OR}^{15}$  or  $-\text{OC}(\text{O})\text{OCH}_2\text{OR}^{16}$ ,  $\text{R}^4$  represents a  $\text{C}_2\text{-C}_6$ -alkyl group;  $\text{R}^5$  represents a  $\text{C}_1\text{-C}_6$ -alkyl group;  $\text{R}^6$ ,  $\text{R}^7$ ,  $\text{R}^8$ ,  $\text{R}^9$ ,  $\text{R}^{10}$ ,  $\text{R}^{12}$ , and  $\text{R}^{13}$  each independently represent a hydrogen atom, or a  $\text{C}_1\text{-C}_6$ -alkyl group optionally substituted by at least one hydroxyl group;  $\text{R}^{11}$  represents a hydrogen atom, or a  $\text{C}_1\text{-C}_6$ -alkyl group optionally substituted by at least one substituent independently selected from hydroxyl and  $\text{C}_1\text{-C}_6$ -alkoxy; and  $\text{R}^{14}$ ,  $\text{R}^{15}$  and  $\text{R}^{16}$  each independently represent a  $\text{C}_1\text{-C}_6$ -alkyl group; with the provisos that (i) when E represents  $\text{NEC}(\text{O})$ , X represents O, S or NH and Y represents O, then Z represents  $-\text{NR}^6\text{R}^7$  where  $\text{R}^6$  represents a

hydrogen atom and  $R^7$  represents either a hydrogen atom or a  $C_1$ - $C_6$ -alkyl group substituted by at least one hydroxyl group, and (ii) when E represents  $NHC(O)$ , X represents O, S or NH, Y represents NH, and  $R^5$  represents  $CH_2CH_2$ , then Z is not -OH or imidazolyl; or a pharmaceutically acceptable salt or solvate thereof.


A compound of general formula:



wherein D represents  $CH_2$  or  $CH_2CH_2$ ; E represents  $C(O)NH$  or  $NHC(O)$ ;  $R^1$  and  $R^2$  each independently represent hydrogen, halogen, amino, nitro,  $C_1$ - $C_6$ -alkyl or trifluoromethyl, but  $R^1$  and  $R^2$  may not both simultaneously represent hydrogen;  $R^3$  represents a group of formula




$R^4$  represents a  $C_1$ - $C_6$ -alkyl group; X represents an oxygen or sulphur atom or a group  $NR^{13}$ , SO or  $SO_2$ ;  $R^5$  represents hydrogen, or  $R^5$  represents  $C_1$ - $C_6$ -alkyl or  $C_2$ - $C_6$ -alkenyl, each of which may be optionally substituted by at least one substituent selected from halogen, hydroxyl, (di) $C_1$ - $C_6$ -alkylamino,  $-Y-R^6$ ,




$NH_2$ , and a 5- or 6-membered heteroaromatic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur which heteroaromatic ring may itself be optionally substituted by at least one substituent selected from halogen, hydroxyl and  $C_1$ - $C_6$ -alkyl; Y represents an oxygen or sulphur atom or a group NH, SO or  $SO_2$ ;  $R^6$  represents a group  $-R^7Z$  where  $R^7$  represents a  $C_2$ - $C_6$ -alkyl group and Z represents an  $-OH$ ,  $-CO_2H$ ,  $-NR^8R^9$ ,  $-C(O)NR^{10}R^{11}$  or -

$N(R^{12})C(O)-C_1-C_6$ -alkyl-group, and, in the case where Y represents an oxygen or sulphur atom or a group NH,  $R^6$  additionally represents hydrogen,  $C_1-C_6$ -alkyl,  $C_1-C_6$ -alkylcarbonyl,  $C_1-C_6$ -alkoxycarbonyl,  $-C(O)NR^{14}R^{15}$ ,  $-CH_2OC(O)R^{16}$ ,  $-CH_2OC(O)OR^{17}$  or  $-C(O)OCH_2OR^{18}$ ;  $R^8$ ,  $R^9$ ,  $R^{10}$ ,  $R^{11}$  and  $R^{12}$  each independently represent a hydrogen atom or a  $C_1-C_6$ -alkyl group;  $R^{13}$  represents hydrogen,  $C_3-C_8$ -cycloalkyl,  $C_3-C_8$ -cycloalkylmethyl, or  $R^{13}$  represents a  $C_1-C_6$ -alkyl group optionally substituted by at least one substituent selected from hydroxyl to and  $C_1-C_6$ -alkoxy; and  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ , and  $R^{18}$  each independently represent a  $C_1-C_6$ -alkyl group; with the proviso that when E is  $C(O)NH$ , X is O, NH or  $N(C_1-C_6\text{-alkyl})$ , then  $R^5$  is other than a hydrogen atom or an unsubstituted  $C_1-C_6$ -alkyl group; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:




wherein m represents 1, 2 or 3; each  $R^1$  independently represents a hydrogen or halogen atom; A represents  $C(O)NH$  or  $NHC(O)$ ; Ar represents a group



X represents a bond, an oxygen atom or a group CO,  $(CH_2)_{1-6}$ ,  $CH=$ ,  $(CH_2)_{1-6}O$ ,  $O(CH_2)_{1-6}$ ,  $O(CH_2)_{2-8}O$ ,  $O(CH_2)_{2-3}O(CH_2)_{1-3}$ ,  $CR'(OH)$ ,  $(CH_2)_{1-3}O(CH_2)_{1-3}$ ,  $(CH_2)_{1-3}O(CH_2)_{2-3}O$ ,  $NR^5$ ,  $(CH_2)_{1-6}NR^5$ ,  $NR^5(CH_2)_{1-6}$ ,  $(CH_2)_{1-3}NR^5(CH_2)_{1-3}$ ,  $O(CH_2)_{2-6}NR^5$ ,  $O(CH_2)_{2-3}NR^5(CH_2)_{1-3}$ ,  $(CH_2)_{1-3}NR^5(CH_2)_{2-3}O$ ,  $NR^5(CH_2)_{2-6}O$ ,  $NR^5(CH_2)_{2-3}O(CH_2)_{1-3}$ ,  $CONR^5$ ,  $NR^5CO$ ,  $S(O)_n$ ,  $S(O)_nCH_2$ ,  $CH_2S(O)_n$ ,  $SO_2NR^5$  or  $NR^5SO_2$ ; n is 0, 1 or 2;  $R^1$  represents a hydrogen atom or a  $C_1-C_6$ -alkyl group; one of  $R^2$  and  $R^3$  represents a halogen, cyano, nitro, amino, hydroxyl, or a group selected from (i)  $C_1-C_6$ -alkyl

optionally substituted by at least one C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, (ii) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, (iii) C<sub>1</sub>-C<sub>6</sub>-alkyloxy optionally substituted by at least one C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, and (iv) C<sub>3</sub>-C<sub>8</sub>-cycloalkyloxy, each of these groups being optionally substituted by one or more fluorine atoms, and the other of R<sup>2</sup> and R<sup>3</sup> represents a hydrogen or halogen atom; either R<sup>4</sup> represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl, -NR<sup>6</sup>R<sup>7</sup>, -(CH<sub>2</sub>)<sub>r</sub>NR<sup>6</sup>R<sup>7</sup> and CONR<sup>6</sup>R<sup>7</sup>, r is 1, 2, 3, 4, 5 or 6; R<sup>5</sup> represents a hydrogen atom or a C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>3</sub>-C<sub>8</sub>-cycloalkyl group; R<sup>6</sup> and R<sup>7</sup> each independently represent a hydrogen atom or a C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-hydroxyalkyl or C<sub>3</sub>-C<sub>8</sub>-cycloalkyl group, or R<sup>6</sup> and R<sup>7</sup> together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring; with the provisos that, (a) when A represents C(O)NH and R<sup>4</sup> represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and (b) when A represents C(O)NH and X represents a group (CH<sub>2</sub>)<sub>1-6</sub> or O(CH<sub>2</sub>)<sub>1-6</sub>, then R<sup>4</sup> does not represent an unsubstituted imidazolyl, unsubstituted morpholinyl, unsubstituted piperidinyl or unsubstituted pyrrolidinyl group, and (c) when A represents NHC(O) and R<sup>4</sup> represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and (d) when A represents NHC(O) and X represents O(CH<sub>2</sub>)<sub>1-6</sub>, NH(CH<sub>2</sub>)<sub>1-6</sub> or SCH<sub>2</sub>, then R<sup>4</sup> does not represent an unsubstituted 1-piperidinyl or unsubstituted 1-pyrrolidinyl group, and (e) when A represents NHC(O) and X represents O(CH<sub>2</sub>)<sub>2-3</sub>NH(CH<sub>2</sub>)<sub>2</sub>, then R<sup>4</sup> does not represent an imidazolyl group; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:




X represents a nitrogen atom or a group  $C(R^5)$ ; Y represents an oxygen or sulphur atom or a group  $NR^6$ ; either  $R^1$  and  $R^2$  each independently represent a hydrogen atom or a  $C_1$ - $C_4$ -alkyl group but do not both simultaneously represent a hydrogen atom, or  $R^1$  and  $R^2$  together represent a group  $-CH_2ZCH_2-$ ; Z represents a bond, an oxygen or sulphur atom or a group  $CH_2$  or  $NR^7$ ; m is 0 or 1;  $R^3$  represents a 5- to 10-membered unsaturated ring system which may comprise from 1 to 4 ring heteroatoms independently selected from nitrogen, oxygen and sulphur, the ring system being optionally substituted by one or more substituents independently selected from halogen, nitro, cyano,  $NR^8R^9$ ,  $C_1$ - $C_4$ -alkyl- $C(O)NH-$ ,  $NHR^{12}C(O)-$ ,  $C_1$ - $C_4$ -alkyl- $SO_2-$ ,  $C_1$ - $C_4$ -alkyl- $SO_2NH-$ ,  $C_1$ - $C_4$ -alkyl- $NHSO_2-$ ,  $C_1$ - $C_4$ -alkoxy, and  $C_1$ - $C_4$ -alkyl optionally substituted by one or more fluorine atoms;  $R^4$  represents a phenyl or pyridinyl group, each of which is substituted in an ortho position with a substituent selected from halogen,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio, and  $C_1$ - $C_4$ -alkyl optionally substituted by one or more fluorine atoms, the phenyl or pyridinyl group being optionally further substituted by one or more substituents independently selected from halogen, cyano, hydroxyl,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkyl- $NH-$ ,  $NHR^{13}$ - $C_1$ - $C_4$ -alkyl-,  $C_1$ - $C_4$ -alkyl- $SO_2-$ ,  $C_1$ - $C_4$ -alkyl- $SO_2NH-$ ,  $C_1$ - $C_4$ -alkyl- $NHSO_2-$ ,  $C_1$ - $C_4$ -alkyl- $C(O)NH-$ ,  $C_1$ - $C_4$ -alkyl- $NHC(O)-$ ,  $-D-G$ ,  $C_1$ - $C_4$ -alkoxy optionally substituted by  $-NR^{14}R^{15}$  or by  $R^{16}$ , and  $C_1$ - $C_4$ -alkyl optionally substituted by one or more fluorine atoms or by one or more hydroxyl groups, or  $R^4$  represents a 9- or 10-membered unsaturated bicyclic ring system which may comprise from 1 to 4 ring heteroatoms independently selected from nitrogen, oxygen and sulphur, the bicyclic ring system being optionally substituted by one or more substituents independently selected from halogen, oxo,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkylthio and  $-NR^{10}R^{11}$ ; D represents an oxygen atom or a group  $(CH_2)_{sub.n}$  or  $CH_2NH$ ; n is 1, 2 or 3; G represents a piperazinyl, morpholinyl or 2,5-diazabicyclo[2.2.1]heptyl group, or G represents a piperidinyl group optionally substituted by amino;  $R^5$  represents a hydrogen atom, or a hydroxyl or  $C_1$ - $C_4$ -alkoxy group;  $R^6$  represents a hydrogen atom, or a cyano, nitro, hydroxyl,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -alkoxy group;  $R^7$ ,  $R^8$  and  $R^9$  each independently represent a hydrogen atom or a  $C_1$ - $C_4$ -alkyl group;  $R^{10}$  and  $R^{11}$  each independently represent a hydrogen atom or a  $C_1$ - $C_4$ -alkyl group, or  $R^{10}$  and  $R^{11}$  together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated heterocyclic ring comprising one or two ring nitrogen atoms;  $R^{12}$  represents a hydrogen atom, or a  $C_1$ -




C<sub>4</sub>-alkyl group optionally substituted by amino; R<sup>13</sup> represents a hydrogen atom, or a C<sub>1</sub>-C<sub>4</sub>-alkyl group optionally substituted by hydroxyl; R<sup>14</sup> and R<sup>15</sup> each independently represent a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub>-alkyl group optionally substituted by hydroxyl, or R<sup>14</sup> and R<sup>15</sup> together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated heterocyclic ring comprising one or two ring nitrogen atoms; and R<sup>16</sup> represents a 1-(C<sub>1</sub>-C<sub>4</sub>-alkyl)-piperidinyl group; with the proviso that when m is 0, X is N and Y is O, then R<sup>4</sup> does not represent 2-benzothiazolyl; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:




wherein: each R<sup>1</sup> independently represents a hydrogen or halogen atom, or a trifluoromethyl, cyano, nitro, C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>1</sub>-C<sub>6</sub>-alkoxy group; T represents an oxygen atom or a group NH; U represents an oxygen or sulphur atom or a group NH; Ar represents a group:



X represents a bond, an oxygen atom or a group CO, CH<sub>2</sub>, CH<sub>2</sub>O, O(CH<sub>2</sub>)<sub>m</sub>, CH<sub>2</sub>OCH<sub>2</sub>, NR<sup>5</sup>, CH<sub>2</sub>NR<sup>5</sup>, NR<sup>5</sup>CH<sub>2</sub>, CH<sub>2</sub>NR<sup>5</sup>CH<sub>2</sub>, CONR<sup>5</sup>, S(O)<sub>n</sub> or SO<sub>2</sub>NR<sup>5</sup>, m is 1, 2 or 3; n is 0, 1 or 2; one of R<sup>2</sup> and R<sup>3</sup> represents a halogen, cyano, nitro, amino, hydroxyl, or a group selected from C<sub>1</sub>-C<sub>6</sub>-alkyl optionally substituted by at least one C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-alkyloxy optionally substituted by at least one C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyloxy, S(O)<sub>p</sub>C<sub>1</sub>-C<sub>6</sub>-alkyl or S(O)<sub>q</sub>C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, each of these groups being optionally substituted by one or more fluorine atoms, and the other of R<sup>2</sup> and R<sup>3</sup> represents a hydrogen or halogen atom or a methyl group; p is 0, 1 or 2; q is 0, 1 or 2; R<sup>4</sup> represents di(C<sub>1-2</sub>alkyl)N(CH<sub>2</sub>)<sub>t</sub> where t is 0, 1 or 2 or

imidazolyl, or  $R^4$  represents a 3- to 9-membered saturated heterocyclic ring system containing one or two nitrogen atoms, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl,  $C_1$ - $C_6$ -alkyl, acetyl, hydroxy $C_1$ - $C_6$ -alkyl,  $-NR^6R^7$ ,  $-(CH_2)_rNR^6R^7$ ,  $-CONR^6R^7$  and pyrimidinyl, or  $R^4$  represents a 3- to 8-membered saturated carbocyclic ring system substituted by one or more substituents independently selected from  $-NR^6R^7$ ,  $-(CH_2)_rNR^6R^7$ ,  $-CONR^6R^7$  the ring system being optionally further substituted by one or more substituents independently selected from fluorine atoms, hydroxyl and  $C_1$ - $C_6$ -alkyl;  $r$  is 1, 2, 3, 4, 5 or 6;  $R^5$  represents a hydrogen atom or a  $C_1$ - $C_6$ -alkyl or  $C_3$ - $C_8$ -cycloalkyl group; and  $R^6$  and  $R^7$  each independently represent a hydrogen atom or a  $C_1$ - $C_6$ -alkyl or  $C_3$ - $C_8$ -cycloalkyl group, or  $R^6$  and  $R^7$  together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, provided that when  $R^3$  represents a cyano group, then  $X$  is other than a bond; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:



wherein  $X$  represents an oxygen or sulphur atom or a group  $NH$ ,  $CH_2$ ,  $CH_2CH_2$  or  $OCH_2$ ;  $Y$  represents a group  $CH_2$  or  $C=O$ ;  $R^1$  represents a pyridyl or pyrimidinyl group;  $R^2$  represents a phenyl, pyridyl or pyrimidinyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or an amino, cyano, hydroxyl, nitro,  $C_1$ - $C_6$ -alkyl, halo- $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -alkylthio, (di) $C_1$ - $C_6$ -alkylamino,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl,  $C_1$ - $C_6$ -alkylsulphinyl,  $C_1$ - $C_6$ -alkylsulphonyl,  $-NR^3SO_2R^4$  or  $-SO_2NR^5R^6$  group, or a group  $-Z-(CH_2)_p-Z-(CH_2)_q-H$  wherein each  $Z$  independently represents a nitrogen or oxygen atom,  $p$  is an integer from 2 to 5 and  $q$  is 0 or an integer from 1 to 5;  $R^3$  and  $R^4$  each independently represent a hydrogen atom or a  $C_1$ - $C_6$ -alkyl group; and  $R^5$  and  $R^6$  each independently represent a hydrogen atom or a  $C_1$ - $C_6$ -alkyl group,

or together with the nitrogen atom to which they are attached form a pyrrolidinyl or piperidinyl group; or a pharmaceutically acceptable salt or solvate thereof.

An additional embodiment of the invention provides a method for treating affective disorders by administering an agent modulating the activity of P2X7R, such as an antagonist of the P2X7R. The term "antagonist" means an agent or drug or a compound that opposes the physiological effects of another. Examples of P2X7R antagonists include, but are not restricted to, adamantane derivatives, isoquinolines and their derivatives, substituted phenyl compounds, piperidine derivatives, piperazine derivatives. P2X7R antagonists are described in the art and include the compounds found in Chen et al., Bioconjugate Chem. 13 (2002), 1100-1111; WO 99/29660; WO 99/29661; WO 99/296896; WO 00/61569; WO 01/42194; WO 01/44170; WO 01/44213; WO 00/71529; WO 01/46200. P2X7R activity can also be modulated by RNA-based interference mechanisms and methods such as, but not limited to, small interference RNA (siRNA) molecules, and long double-stranded RNA (dsRNA).

Since it was unexpectedly found that agonists of P2X7R improve the symptoms of mice selected for anxiety and depressive behaviour, the present invention relates in a further embodiment, to a method for treating affective disorders such as anxiety and depressive behaviour by administering an agent modulating the activity of P2X7R such as an agonist of the P2X7R.. Examples of P2X7R agonists include but are not restricted to ATP, ATP-4, and BzATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate ( $C_{24}H_{24}N_5O_{15}P_3$ )). Preferably, the P2X7R agonist to be used for treating affective disorders is BzATP. More preferably, BzATP is used to treat depression or anxiety as demonstrated in Example 9 hereinbelow.

Dosage, pharmaceutical preparation and delivery of P2X7R modulating agent for use in accordance with the present invention may be formulated in conventional manner according to methods found in the art, using one or more physiological carriers or excipient, see, for example Ansel et al., "Pharmaceutical Dosage Forms and Drug Delivery Systems", 7<sup>th</sup> edition, Lippincott Williams & Wilkins Publishers, 1999. Thus, the P2X7R modulating agent and its physiologically acceptable salts and solvates

may be formulated for administration by inhalation, insufflation (either through the mouth, or nose), oral, buccal, parenteral, or rectal administration.

For oral administration, the pharmaceutical composition of the P2X7 modulating agent may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutical acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone, hydroxypropyl methylcellulose), fillers (e.g., lactose, microcrystalline cellulose, calcium hydrogen phosphate), lubricants (e.g., magnesium stearate, talc, silica), disintegrants (e.g., potato starch, sodium starch glycolate), or wetting agents (e.g., sodium lauryl sulphate). Liquid preparations for oral administration may take the form of, for example, solutions, syrups, or suspensions, or may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparation may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol, syrup, cellulose derivatives, hydrogenated edible fats), emulsifying agents (e.g., lecithin, acacia), non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol, fractionated vegetable oils), preservatives (e.g., methyl or propyl-p-hydroxycarbonates, sorbic acids). The preparations may also contain buffer salts, flavouring, coloring and sweetening agents as deemed appropriate. Preparations for oral administration may be suitably formulated to give controlled release of the agent modulating P2X7R activity.

For administration by inhalation, the agent modulating P2X7R activity for use according to the present invention is conveniently delivered in the form of an aerosol spray presentation from a pressurised pack or a nebulizer, with the use of a suitable propellant (e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas). In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatine, for use in an inhaler or insufflator may be formulated containing a powder mix of the P2X7R activity modulating agent and a suitable powder base such as lactose or starch.

An agent modulating P2X7R activity may be formulated for parenteral administration by injection, for example, by bolus injection or continuous infusion. Site of injections include intra-venous, intra-peritoneal or sub-cutaneous. Formulations for injection may be presented in unit dosage form (e.g., in phial, in multi-dose container), and with an added preservative. The agent modulating P2X7R activity may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing, or dispersing agents. Alternatively, the agent may be in powder form for constitution with a suitable vehicle (e.g., sterile pyrogen-free water) before use.

An agent modulating P2X7R activity may, if desired, be presented in a pack, or dispenser device which may contain one or more unit dosage forms containing the said agent. The pack may for example comprise metal or plastic foil, such as blister pack. The pack or dispenser device may be accompanied with instruction for administration.

In a more preferred embodiment the aforementioned methods or uses are envisaged to treat affective disorders selected from the group consisting of major depression, generalized anxiety disorder and bipolar disorder.

In a particularly preferred embodiment said major depression is selected from the group consisting of major depression, dysthymia, atypical depression, premenstrual dysphoric disorder and seasonal affective disorder.

In another particularly preferred embodiment said generalized anxiety disorder is selected from the group consisting of panic disorder, phobias, agoraphobia, social phobia, specific phobia, obsessive-compulsive disorder, post-traumatic stress disorder, separation anxiety disorder, mania, hypomania and cyclothymic disorder.

A still also particularly preferred embodiment is that said bipolar disorder is bipolar disorder type I or bipolar disorder type II.

Additionally, the present invention relates to a kit comprising the nucleic acid molecule, the vector, the host, the polypeptide, the antibody or the aptamer, the primer or pair of primers of the invention or the molecule as identified or characterized in a method herein below of the present invention.

Advantageously, the kit of the present invention further comprises, optionally (a) reaction buffer(s), storage solutions and/or remaining reagents or materials required for the conduct of scientific or diagnostic assays or the like. Furthermore, parts of the kit of the invention can be packaged individually in vials or bottles or in combination in containers or multicontainer units.

The kit of the present invention may be advantageously used, inter alia, for carrying out the method of producing a polypeptide of the invention, the method(s) of identification and/or characterization of molecules specifically interacting with P2X7R ATP-gated ion channels as described herein below and/or it could be employed in a variety of applications referred herein, e.g., as diagnostic kits, as research tools or therapeutic tools. Additionally, the kit of the invention may contain means for detection suitable for scientific, medical and/or diagnostic purposes. The manufacture of the kits follows preferably standard procedures which are known to the person skilled in the art.

Furthermore, the present invention relates to a method for identifying compounds or mixtures of compounds which are capable of specifically interacting with a polypeptide of the present invention, comprising the steps of (a) contacting a polypeptide of the present invention with a candidate compound or mixture of compounds to be tested; and (b) determining whether said is capable of specifically interacting with said polypeptide. The polypeptide may be provided directly or by expression of a corresponding nucleic acid molecule or vector of the invention, e.g., in vitro or in a suitable host cell.

Additionally, the present invention relates to a method for the characterization of compounds which are capable of altering characteristics of the polypeptides of the present invention, comprising the steps of (a) contacting a polypeptide of the invention with said compound; and (b) determining whether the compound alters a characteristic of said polypeptide.

The term "altering characteristic of the polypeptide of the present invention" means that the functional characteristics to the polypeptides of the present invention in comparison to functional characteristics which they had before being contacted with the compounds identified by the above-described method: as described hereinabove are altered; i.e. changed.

Said identification and/or characterization of which are capable of interacting with or altering characteristics of the polypeptide of this invention, may be, inter alia, achieved by transfecting an appropriate host with a nucleic acid molecule of invention. Said hosts comprise, but are not limited to, HEK 293 cells or are injected into frog oocytes, preferably a *Xenopus* oocyte for functional expression (Goldin, *Methods Enzymol.* 207 (1992), 266). Expressed P2X7R ATP-gated channels can be examined using standard two-electrode voltage clamp techniques (Stuhmer, *Methods Enzymol.* 207 (1992), 319; Kohler, *Science* 273 (1996), 1709). After expression of a P2X7R ATP-gated ion channel as defined herein, membrane currents may be deduced in the absence and/or presence of the molecule to be identified and/or characterized. Methods for the deduction of membrane currents are well known in the art and comprise, e.g., patch clamp methods as described in Hamill, *Pflüger's Arch.* 391 (1981), 85-100 or two-electrode voltage clamp in oocytes, as described in Methfessel, *Pflügers Archive* 407 (1986), 577-588. In accordance with the present invention the term "interacting with the polypeptides of the present invention" means that the polypeptides of the present invention interact directly and/or indirectly with compounds identified by the method described above. Furthermore, the present invention relates to a method of screening for molecules which are capable of interacting with the polypeptide of this invention, comprising the steps of (a) contacting a polypeptide of the invention with a molecule; and (b) measuring and/or detecting a response; and (c) comparing said response to a standard response as measured in the absence of said candidate molecule.

The present invention also relates to a method for identifying a compound which is capable of enhancing or reducing the expression of the P2X7R gene comprising the steps of contacting a cell which expresses the P2X7R gene from its natural

promoter or a reporter gene driven by the P2X7R promoter and determining whether the expression of the gene is increased or reduced when compared to conditions in which the compound is not present.

Potential candidate molecules or candidate mixtures of molecules may be, inter alia, substances, compounds or compositions which are of chemical or biological origin, which are naturally occurring and/or which are synthetically, recombinantly and/or chemically produced or compounds or compositions described hereinabove. Thus, candidate molecules may be proteins, protein-fragments, peptides, amino acids and/or derivatives thereof or other compounds, such as ions, which bind to and/or interact with wild-type P2X7R ATP-gated ion channels. Such binding and/or interacting candidate compounds may be found employing, inter alia, yeast two-hybrid systems or modified yeast two-hybrid systems as described, for example in Fields, Nature 340 (1989), 245-246; Gyuris, Cell 75 (1993), 791-801; or Zervos, Cell 72 (1993), 223-232.

Furthermore, potential candidate molecules may be contacted with a cell, such as an oocyte or a HEK 293 cell, which expresses a polypeptide of the invention or with a membrane patch comprising a polypeptide of the invention and a corresponding response (inter alia, a dose-response response, a current-response, or single current channel response) may be measured in order to elucidate any effect said candidate molecule causes.

Within the scope of the present invention are also methods for identifying, characterizing and for screening of molecules which are capable of interacting with the P2X7R ATP-gated ion channels according to the invention which comprise so-called high-throughput screening methods and similar approaches which are known in the art (Spencer, Biotechnol. Bioeng. 61 (1998), 61-67; Oldenburg, Annu. Rev. Med. Chem. 33 (1998), 301-311) carried out using 96-well, 384-well, 1536-well (and other) commercially available plates. Further methods to be employed in accordance with the present invention comprise, but are not limited to, homogenous fluorescence readouts in high-throughput screenings (as described, inter alia, in Pope, Drug Discovery Today 4 (1999), 350-362). The method of the present invention for identification, characterization and/or screening of molecules capable of interacting



with P2X7R ATP-gated ion channels can, inter alia, employ hosts as defined herein which express the polypeptide of the present invention. Cell-based assays, instrumentation for said assays and/or measurements are well-known in the art and described, inter alia, in Gonzalez, Drug Discovery Today 4 (1999), 431-439 or Ramm, Drug Discovery Today 4 (1999), 401-410. It is also envisaged that the high through put screens described herein are conducted by using, for example cRNA, i.e. synthetic RNA from a cDNA construct) that can be introduced in host cells, such as Xenopus oocytes using routine methods in the art. As an example, direct nucleic acid injection can be employed, such as the Eppendorf microinjection system (Micromnipulator 5171 and Transjector 5242). The injected/transformed cells can be analyzed for ion currents about 4 hours later using patch-clamp techniques which are commonly practiced in the art.

Additionally, the present invention relates to a method for the production of a pharmaceutical composition comprising the steps of a method of the invention for identifying, characterizing and/or screening of molecules which are capable of interacting with and/or altering the characteristics of a P2X7R ATP-gated ion channel of the invention and further comprising a step, wherein a derivative of said identified, characterized and/or screened molecule is generated. Such a derivative may be generated by, inter alia, peptidomimetics.

The invention furthermore relates to a method for the production of a pharmaceutical composition comprising the steps of a method of the invention for identifying, characterizing, screening and/or derivatizing of molecules which are capable of interacting with and/or altering the characteristics of a P2X7R ATP-gated ion channel and formulating the molecules identified, characterized, screened and/or derivatized in pharmaceutically acceptable form.

In a more preferred embodiment the present invention relates to a method wherein said molecule(s) comprise(s) (a) neuroprotective, (a) nootropic and/or (a) antiepileptic molecule(s).

Yet another embodiment of the invention is the use of a P2X7R polypeptide, in particular those according to the present invention, to identify biological, chemical, or

pharmacological agents that can have an antidepressive effect. The term 'agent' refers to a chemical compound or composition capable of inducing a desired therapeutic or prophylactic effect when properly administered to a subject or cell. For example, the present invention allows the generation of cells expressing P2X7R for the identification and characterization of agents which modulate ionic influx and efflux. For example, HEK293 cells, or other cell lines (e.g., HCN-1A, HCN-2, HIT-T15, RIN-m5F, betaTC3, PC12, HT22, SH-SY5Y, Neuro2A or CA77), can be stably transfected with cDNA encoding the human P2X7R and plated in 12, 96 and 384 well plates. Said cells are cultured in appropriate medium. Examples of such medium are well known in the art, see, for example Freshney, "Culture of Animal Cells: A Manual of Basic Technique, 4th edition, Wiley-Liss Publishing, 2000.

Said cells can then be pre-incubated with said agents for 15 min prior to stimulation with 3 mM ATP for 10 minutes. Reactions are then terminated by rapid aspiration of the extracellular medium in each well. The cells in each well are subsequently extracted overnight with 1 ml 10% HNO<sub>3</sub>. Potassium (K<sup>+</sup>) content in the extracts can be determined by atomic absorbance spectrophotometry. Agent function is then measured by the percent inhibition or stimulation of the K<sup>+</sup> release triggered by 3 mM ATP and compared to K<sup>+</sup> release in the absence of the agents. P2X7R activity can also be monitored according to the movement of calcium (Ca<sup>2+</sup>; see Denyer et al., Drug Discov. Today 7 (1998), 323-332; González et al., Drug Discov. Today 9 (1999), 431-439; Helmchen and Waters, Eur. J. Pharmacol. 447 (2002), 119-129). Agents can also be verified in the absence of ATP.

P2X7R activity can also be monitored according to secretion of neurotransmitters such as glutamate and GABA. Neurotransmitter levels in treated cells can be quantified by suitable methods, e.g., Enzyme Linked Immunoabsorbent Assay (ELISA), Radio Immuno Assay (RIA), High Performance Liquid Chromatography (HPLC). Using these methods, a large number of compounds can be screened for increase in neurotransmitter (for example, glutamate) secretion. The release of glutamate can be measured for example by Fluorometric glutamate release assays (e.g., Amplex Red Glutamic Acid/Glutamate Oxidase Assay Kit, Molecular Probes) or High-Throughput ElectroPhysiology.

In a further aspect the present invention uses the P2X7R polypeptides disclosed herein or polypeptides of the present invention in a method for identifying compounds or agents having agonist activity to said P2X7R polypeptides or to the polypeptides of the present invention. Agents and compounds are defined and described hereinabove and hereinbelow. In particular, cells that express the P2X7 gene are contacted with candidate agents, molecules or compounds as described hereinabove and either calcium influx or ethidium bromid entry is measured by methods known in the art, described hereinabove and in particular described in Example 8 hereinbelow. The cells used in the method for identifying agonists to P2X7R are preferably cells of a hippocampal cell line. Hippocampal cell lines are prepared by methods known in the art, for example, decribed in EP 0 773 287 or EP 0 773 292. Non limiting examples of hippocampal cell lines are rat H19-7 hippocampal cells (ATCC-2526) described in Eves et al. Proc. Natl. Acad. Sci. USA 89 (1992), 4373-4377, mouse HN9.10 hippocampal cells described in Lee et al. J. Neurosci. 10 (1992), 1779-1787 or rat Hi5B hippocampal cells described in Renfranz et al., Cell 66 (1991), 713-729. Preferably, the hippocampal cells used in accordance with the aforementioned method are cells of the HT-series (see Davis and Maher (1994), Brain Res. 652, 169-173), Morimoto and Koshland (1990), Neuron 5, 875-880). It is also preferred that the hippocampal cells express the endogenous P2X7R gene. However, it is also envisaged that such cells may be genetically modified by introducing an exogenous P2X7R gene using methods commonly known in the art. More preferably, HT-22 cells are used for identifying agonists to the P2X7R polypeptides described herein or to the polypeptides of the present invention and HT-39 cells are used as a negative control as described in Example 8 hereinbelow.

In another embodiment, cells are transfected with nucleic acid constructs encoding a reporter gene regulated by the P2X7R promoter (see above), an increase or decrease in the expression of the reporter gene in response to biological or pharmaceutical agents can be analyzed using methods that detect levels or status of protein or mRNA present in the corresponding cell or detect biological activities of the reporter gene. Suitable reporter molecules or labels, which may be used, include radionucleotides, enzymes, fluorescent, chemiluminescent or chromogenic agents as

well as substrates, co-factors, inhibitors, magnetic particles, and the like. Designing such drug screening assays are well known in the art; see Harvey ed., 'Advances in drug discovery techniques', John Wiley and Sons, 1998; Vogel and Vogel eds., 'Drug discovery and evaluation: Pharmaceutical assays', Springer-Verlag Berlin, 1997). For example, drug screening in animal models, in vitro tests using animal cells, or in vivo tests involving toxicology tests in animals. An in vitro model can be used for screening libraries of compounds in any of a variety of drug screening techniques.

Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 Daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise carbocyclic or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups.

Candidate agents are also found among biomolecules including peptides, amino acids, saccharides, fatty acids, steroids, purines, pyrimidines, nucleic acids and derivatives, structural analogs or combinations thereof. Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs.

Another technique for drug screening, which may be used, provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO 84/03564. In this method, as applied to the proteins of the invention large numbers of different small test compounds, e.g. aptamers, peptides, low-molecular weight compounds etc., are provided or synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with the proteins or fragments thereof, and washed. Bound proteins are then detected by methods well known in the art. Purified proteins can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support. In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding the protein specifically compete with a test compound for binding the protein. In this manner, the antibodies can be used to detect the presence of any peptide, which shares one or more antigenic determinants with the protein.

The present invention further particularly provides a method, wherein the pharmaceutical composition to be produced further comprises neuroprotective substances, nootropic substances, brilliant blue, piperidine or derivatives thereof, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate or 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate. It is also envisaged that the pharmaceutical compositions to be produced further comprise antidepressants such as fluoxetine, paroxetine, sertraline, fluvoxamine, mirtazapine, reboxetine, nefazodone or lithium carbonate.

In a preferred embodiment of the present invention, the compounds of the aforementioned methods comprise antagonist(s), partial antagonist(s), partial agonist(s) and/or agonist(s) for an altered ATP-gated ion channel P2X7R.

In accordance with the present invention, the term "antagonist" denotes molecules/substances, which are capable of inhibiting and/or reducing an agonistic effect. The term "antagonist" comprises competitive, non-competitive, functional and chemical antagonists as described, inter alia, in Mutschler,

"Arzneimittelwirkungen" (1986), Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, Germany. The term "partial antagonist" in accordance with the present invention means a molecule/substance that is capable of incompletely blocking the action of agonists through, inter alia, a non-competitive mechanism.

In the context of the present invention, an antagonist is preferably a drug that does not provoke a response itself, but blocks agonist-mediated responses. It is a chemical entity that opposes the receptor-associated responses normally induced by another bioactive agent. For the P2X7R, the antagonists have an IC50 between 10 nanomolar and 300 micromolar.

As "agonist", in accordance with this invention, molecules/substances are denoted which have an affinity as well as an intrinsic activity. Mostly, said intrinsic activity ( $\alpha$ ) is defined as being proportional to the quotient of the effect, triggered by said agonist (EA) and the effect which can be maximally obtained in a given biological system (Emax): therefore, the intrinsic activity can be defined as

$$\alpha \sim \frac{E_A}{E_{\max}}$$

The highest relative intrinsic activity results from  $EA/Emax=1$ . Agonists with an intrinsic activity of 1 are full agonists, whereas substances/molecules with an intrinsic activity of  $>0$  and  $<1$  are partial agonists. Partial agonists show a dualistic effect, i.e. they comprise agonistic as well as antagonistic effects.

Preferably, in the context of the present invention, an agonist (or full agonist) is an endogenous substance or a drug that can interact with a receptor and initiate a maximal or complete physiological or a pharmacological response characteristic of that receptor. ATP, the natural ligand for the P2X7R, is an agonist with an EC50 of 300 micromolar while the synthetic P2X7R agonist Bz-ATP has an EC50 of 8 micromolar. Thus, agonists of P2X7R have an EC50 equal or below 300 micromolar. The EC50 is defined as the concentration of agonist that provokes a response half way between the baseline response and maximum response on a dose response curve where the X-axis plots concentration of an agonist and the Y-axis plots ion current. An inverse agonist (also called negative antagonist) is a drug which acts at

the same receptor as that of an agonist, yet produces an opposite effect. A partial agonist is an endogenous substance or a drug that also provokes physiological or a pharmacological response but, the maximum response is less than the maximum response to a full agonist, regardless of the amount of drug applied. In the case of P2X7R, partial agonists have EC50s higher than 300 micromolar.

The person skilled in the art can, therefore, easily employ the compounds and the methods of this invention in order to elucidate the agonistic and/or antagonistic effects and/or characteristics of a compound/molecule/substance to be identified and/or characterized in accordance with any of the above described methods. Preferably, an identified antagonist of the ATP-gated ion channel P2X7R comprising the mutation(s) and/or deletion(s) described hereinabove may be useful to reestablish the properties normally shown by wild-type P2X7R ATP-gated ion channels. An identified agonist of the ATP-gated ion channel P2X7R comprising the mutation(s) and/or deletion(s) described hereinabove may be useful to reestablish the lost functionality of the P2X7R ATP-gated ion channel.

The Figures show:

**Figure 1a.** Genomic map of the region on the human chromosome 12 associated to bipolar affective disorder. Genes found between markers NBG11 and NBG2 are depicted.

**Figure 1b.** Graphic illustrating the multipoint analysis using ASPEX on independent sib-pairs.

**Figure 1c.** Graphic illustrating the multipoint analysis using ASPEX on all sib-pairs

**Figure 1d.** Graphic illustrating the ASPEX sib\_phase by considering only independent sib-pairs

**Figure 1e.** Graphic illustrating the ASPEX sib\_phase by considering all sib-pairs

**Figure 2.** RT-PCR analysis of the complete coding sequence of P2X7R in different tissues

**Figure 3.** P2X7R expression in the olfactory bulb, hypothalamus and ependymal cells in the brain of a stress-free mouse. Magnification 100X.

**Figure 4.** P2X7R expression in the hippocampus/dentate gyrus and subcommisural organ in the brain of a stress-free mouse. Magnification 100X.

**Figure 5.** Floating behaviour in the forced swim test. Passive stress coping behaviour decreased after long-term treatment with the antidepressant paroxetine (Par28: treated with paroxetine for 28 days, per os). Basal n=8; vehicle n=8; Par28 n=8.

**Figure 6.** Comparative analysis of P2X7R expression in the olfactory bulb of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 100X.



**Figure 7.** Comparative analysis of P2X7R expression in the hypothalamus of stress-free, treated-treated and antidepressant-treated mice. Magnification 100X.

**Figure 8.** Comparative analysis of P2X7R expression in ependymal cells of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 100X.

**Figure 9.** Comparative analysis of P2X7R expression in the hippocampus of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 25X.

**Figure 10.** P2X7R expression in the hippocampus of a vehicle treated mouse. Magnification 25X.

**Figure 11.** P2X7R expression in the hippocampus of a mouse treated with the antidepressant paroxetine. Magnification 25X.

**Figure 12.** Detailed expression of P2X7R in the dentate gyrus of a mouse treated with the antidepressant paroxetine. Magnification 400x.

**Figure 13.** Comparative analysis of P2X7R expression and apoptotic cells in the hippocampus of a mouse treated with the antidepressant paroxetine. Magnification 100X.

**Figure 14.** Floating behaviour in the forced swim test. Passive stress coping behaviour increased after acute intrahippocampal (bilateral, dentate gyrus) of siRNA targeting P2X7R. Vehicle n=7; control RNA n=10; P2X7R siRNA n=9.

**Figure 15.** Comparative analysis of P2X7R expression in the hippocampus of mice treated with vehicle, control RNA and of siRNA targeting P2X7R. Magnification 100X upper row, 25X lower row.

**Figure 16a, b, c, d, e.** Three splicing variants caused by polymorphisms in the introns of P2X7R.

**Figure 17.** Expression of P2X7R in immortalized hippocampal cell lines.

**Figure 18.** Increase calcium influx in hippocampal cells treated with a P2X7R agonist compound (BzATP).

**Figure 19a, b.** Entry of ethidium bromide dye into hippocampal cells (a) treated with P2X7R agonist compound (BzATP) or (b) pre-treated with a P2X7R antagonist compound.

**Figure 20.** Effect of intrahippocampal injection of a P2X7R agonist compound (BzATP) on behaviour in the forced swim test.

**Figure 21.** Open field test measuring locomotor activity of mice treated with a P2X7R agonist compound (BzATP).

**Figure 22.** Comparative analysis of apoptotic cells in the hippocampus of a mouse treated with control vehicle solution or a P2X7R agonist compound (BzATP).

A better understanding of the present invention and of its many advantages will be had from the following examples, offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

### **EXAMPLE 1**

#### **Linkage analysis of Bipolar Affective Disorder in a homogeneous human population**

41 families of different sizes containing a total of 485 sampled individuals from the region of Saguenay/Lac St-Jean were used in the linkage analysis. Individuals were distributed according to their diagnoses as follows: 105 individuals afflicted with Bipolar Disorder type I (BPI) or schizoaffective disorder bipolar type; 42 individuals diagnosed with Bipolar Disorder type II (BP II); 54 individuals with recurrent major depression; and 57 individuals with single episode major depression. The remaining 227 individual unaffected and normal. For the purpose of the calculation, the following classification was used: individuals diagnosed with either BPI, schizoaffective disorder, bipolar type, BP II and recurrent major depression were considered as affected (n=201); individuals with a single major depression episode were scored as unknown phenotype (n=57); and all others diagnoses as unaffected (n=227).

Blood samples from each individual were collected in 10-ml K3 EDTA Vacutainer tube (Becton-Dickinson) and genomic DNA was isolated by Puregene DNA Isolation kit (Gentra Systems). Blood was poured into 50 ml conical tube and diluted with four volume of Red Blood Cell Lysis Solution. After an incubation of 10 minutes at room temperature, the tube was centrifuged for 10 minutes at 2,000 g and supernatant was removed leaving behind cell pellet and 200-400 µl of the residual liquid. Cells were resuspended by vortexing the tube and 9 ml of Cell Lysis Solution were added with up and down pipetting. 40 µl of RNase A Solution (20 mg/ml) were added and the sample was mixed by inverting the tube several times. Sample was incubated at 37°C for 15 minutes and cooled to room temperature. 3 ml of Protein Precipitation Solution were added to cell lysate. Tube was vigorously vortexed for 30 seconds and

centrifuged at 2,000 g for 10 minutes. Supernatant was poured into a new tube containing 9 ml of 100% isopropanol. Sample was mixed by inverting gently several times. Tube was centrifuged at 2,000 g for 5 minutes. The DNA white pellet was washed with 10 ml 70% ethanol and the tube was centrifuged at 2,000 g for 3 minutes. Ethanol was poured off and pellet allowed to partially air dry. DNA was solubilized in 500 µl of DNA Hydration Solution. Final concentration was adjusted to 300-400 µg/ml.

A fluorescent-based method was used for the genotyping of microsatellite markers. Briefly, the region encompassing each repeated sequence was amplified by PCR using an unlabeled primer and a fluorescent-labeled primer (Applied biosystems inc, CA, USA). The marker-associated dyes and the corresponding PCR product length are listed in table 2. The PCR reaction was performed using 10ng of DNA sample, 0,2 unit of Taq platinum DNA polymerase (Invitrogene, CA, USA), 20 mM Tris-Cl (pH 8.4), 50 mM KCl, 1.5mM MgCl<sub>2</sub>, 100µM of dNTP, and 1,5 µM of each primer in a final volume of 7 µl. The samples were incubated at 95°C for 3 minutes to activate the Taq platinum DNA polymerase, then 10 cycles of PCR amplification were performed as follows: 95°C for 15 seconds; 58°C for 15 seconds; 72°C for 30 seconds; after that 15 cycles were performed as follow: 89°C for 15 seconds; 58°C for 15 seconds; 72°C for 30 seconds. Finally, the samples were incubated at 72°C for 30 min. Following the PCR amplification samples were pooled according to their dye-labeled primer and their PCR product length (pool of four samples). Pooled sample were separated on an ABI 3100 DNA analyzer (Applied Biosystems inc, CA, USA). The resulting data were analysed using Genemapper2 (Applied Biosystems inc, CA, USA), and compiled in a 4D database (ACIUS) designed in a Macintosh environment as previously described (Morissette et al., Am. J. Med. Genet. (Neuropsychiatr. Genet.) 88 (1999), 567-587)

Markers used in the following linkage analysis are shown in table 2. Recombination fraction (q) between successive markers was computed according to the analyzed families.

**Table 2. Genomic markers used for the linkage analysis**

Locus	Associated dye	Allele length (bp)	Distance (q)	Cumulative distance (cM)	Heterozygosity (%)
D12S1619	VIC	170-210	0.0135	0.00	74.5
NBG11	VIC	204-218	0.006	1.37	65.5
D12S1666	FAM	241-281	0.001	1.97	66.9
NBG5	VIC	253-261	0.001	2.07	38.3
D12S1721	VIC	263-299	0.005	2.17	72.1
NBG8	VIC	166-188	0.011	2.67	73.3
NBG6	NED	182-218	0.0115	3.79	73.9
NBG9	VIC	156-180	0.0035	4.95	68.9
NBG10	FAM	174-186	0.001	5.30	49.7
NBG12	NED	165-207	0.009	5.40	64.2
NBG4	NED	171-199	0.001	6.31	66.4
NBG3	VIC	182-206	0.006	6.41	64.8
NBG2	VIC	171-199		7.01	54.2

-Haldane's map function was used for cumulative distance in cMorgans.

For bipoint parametric analysis, MOD score analysis were used where parametric LOD score were maximized over genetic models.

The following results were obtained under MOD score analysis for recessive models.

**Table 3. MOD score analysis for recessive models**

Locus	Distance (q)	Cumulative distance (cM)	LOD score ( $q_{\max}$ )
D12S1619	0.0135	0.00	3.46 (0.10)
NBG11	0.006	1.37	4.06 (0.04)
D12S1666	0.001	1.97	1.22 (0.14)
NBG5	0.001	2.07	0.66 (0.16)
D12S1721	0.005	2.17	2.82 (0.10)
NBG8	0.011	2.67	1.51 (0.00)
NBG6	0.0115	3.79	4.77 (0.06)
NBG9	0.0035	4.95	0.75 (0.22)
NBG10	0.001	5.30	0.74 (0.00)
NBG12	0.009	5.40	1.41 (0.16)
NBG4	0.001	6.31	3.56 (0.08)
NBG3	0.006	6.41	3.96 (0.08)
NBG2		7.01	2.59 (0.10)

Model-free LOD score studies using ANALYZE, sib\_phase from the ASPEX V1.85 package (David Hinds and Neil Risch 1999; <ftp://lahmed.stanford.edu/pub/aspepx>, see also <http://watson.hgen.pitt.edu/docs/usage.html>) and SIMWALK2 (Sobel and Lange, Am J Hum Genet 58 (1996), 1323-1337) were performed to analyze the allele sharing among affected sib-pairs. The ANALYZE program weights sibships according to their size. The ASPEX sib\_phase program uses allele frequencies to reconstruct missing information, and is tailored for data sets where parents are missing, but additional typed children may be used to reconstruct and phase the parents. SimWalk2 is a statistical genetics computer application for haplotype, parametric linkage, non-parametric linkage (NPL), identity by descent (IBD) and mistyping analyses on any size of pedigree. SimWalk2 uses Markov chain Monte Carlo (MCMC) and simulated annealing algorithms to perform these multipoint analyses.

ASPEX sib\_phase was used with two computational strategies: First, by using strictly independent sib pairs; secondly, by using all affected sib-pair combinations. ASPEX was performed for bi-point and multipoint calculations.

The bi-point results observed with ANALYZE and ASPEX are shown in Table 4.

**Table 4. Bi-point results observed with ANALYZE and ASPEX**

Locus	Distance (q)	Cumulative distance (cM)	Sib-pair from ANALYZE LOD score	sib_phase LOD score indep. sib-pairs	sib_phase LOD score all sib-pairs
D12S1619	0.0135	0.00	2.31	2.55	3.14
NBG11	0.006	1.37	2.83	2.72	3.27
D12S1666	0.001	1.97	1.01	2.52	3.14
NBG5	0.001	2.07	0.50	2.52	3.13
D12S1721	0.005	2.17	1.57	2.51	3.12
NBG8	0.011	2.67	0.51	2.24	2.75
NBG6	0.0115	3.79	2.55	2.11	2.64
NBG9	0.0035	4.95	0.49	1.65	1.97
NBG10	0.001	5.30	0.77	1.45	2.10
NBG12	0.009	5.40	0.47	1.44	2.17
NBG4	0.001	6.31	1.21	1.29	3.07
NBG3	0.006	6.41	1.84	1.29	3.07
NBG2		7.01	1.24	1.22	3.00

SIMWALK2 computed four different statistics based on descent trees. These statistics measure the degree of clustering among the marker alleles descending from the founders.

Statistic A is the number of different founder-alleles contributing alleles to the affected it is most powerful at detecting linkage to a recessive trait. Statistic B is the maximum number of alleles among the affected descended from any one founder-allele and most powerful at detecting linkage to a dominant trait. Statistic C is the 'entropy' of the marker alleles among the affected. Statistic D is the extent of allele sharing among all affected pairs as measured by their IBD kinship coefficient. Statistics C and D are more general statistics indicating whether a few founder-alleles are overly represented among the affected.

Table 5 shows the results observed with SIMWALK2. The authors signal that p-values should be generally conservative. They are expressed as  $-\text{Log}(\text{p-values})$ . For correspondence purpose,  $-\text{Log}(0.05)=1.30$ ,  $-\text{Log}(0.01)=2$ ,  $-\text{Log}(0.001)=3$  etc.

**Table 5. SIMWALK2 analysis**

Locus	Distance (q)	Cumul. Distance (cM)	STAT(A) -Log(p-value)	STAT(B) -Log(p-value)	STAT(C) -Log(p-value)	STAT(D) -Log(p-value)
D12S1619	0.0135	0.00	1.4550	0.4103	1.1306	1.1310
NBG11	0.006	1.37	2.0157	1.4375	1.5955	1.9845
D12S1666	0.001	1.97	2.0236	0.9765	1.4727	1.4614
NBG5	0.001	2.07	1.7596	0.8558	1.3866	1.3602
D12S1721	0.005	2.17	1.6628	1.1692	1.4235	1.6384
NBG8	0.011	2.67	1.5374	0.6940	1.0623	1.1552
NBG6	0.0115	3.79	1.5896	0.4452	1.0935	1.1786
NBG9	0.0035	4.95	1.2677	0.3815	0.8412	0.9133
NBG10	0.001	5.30	1.1117	0.3642	0.6987	0.7554
NBG12	0.009	5.40	1.0809	0.3485	0.6694	0.7179
NBG4	0.001	6.31	1.1024	0.4148	0.6368	0.8544
NBG3	0.006	6.41	1.1040	0.4146	0.6373	0.8559
NBG2		7.01	1.0963	0.5380	0.6587	0.9356

Multipoint result observed with ASPEX when only independent sib-pairs were used (Figure 1b). The maximum LOD score value was observed at NBG11.

Multipoint result observed with ASPEX when all sib-pairs were considered (Figure 1c). The maximum LOD score value was observed at NBG11 but a second peak appeared at NBG4 and NBG3.

Multipoint and bi-point LOD score values computed by ASPEX were similar. The second peak, observed when all sib-pairs are used, may be explained by the presence of a recombinant affected individual, with many affected sibs, sharing the chromosomal region telomeric to NBG12. This kind of individuals has a large impact on LOD score values when all sib-pairs are used instead of one sib-pair. This situation was observed in two sibships.

Strata analysis was subsequently performed. Although HOMOG did not detect evidence for heterogeneity, a homogeneity test was constructed based on allele sharing found in selected chromosomal regions. Only 20 of the 41 families were used for this analysis since the others were not genotyped in all these regions. For each marker within the selected regions, the proportion of alleles shared IBD by affected sib-pairs was estimated with ASPEX (sib\_phase). For each region retained, the proportion of shared alleles was used as variable for a Principal Component Analysis



and the first principal component as an index of linkage. Correlation analysis was done on these indexes to detect heterogeneity (correlation < 0) or epistasis (correlation > 0). Fisher algorithm was used to classify into two groups of families as linked or unlinked to a particular locus. A negative correlation was observed between the chromosome 12 region and the chromosome 15 area ( $r=-0.51$ ;  $p=0.023$ ). Cluster analysis suggested that 11 families out of 20 were linked to chromosome 12. This sub-sample was called the strata.

This strata included 11 families (266 sampled individuals) that include 52 BPI or schizoaffective disorder, bipolar type, 20 BP II and 28 recurrent major depression

The following MOD score values illustrated in Table 6 were obtained under recessive models.

**Table 6. MOD scores under recessive models**

Locus	Distance (q)	Cumulative distance (cM)	LOD score ( $q_{\max}$ )
D12S1619	0.0135	0.00	4.03 (0.08)
NBG11	0.006	1.37	4.98 (0.00)
D12S1666	0.001	1.97	1.49 (0.12)
NBG5	0.001	2.07	0.79 (0.14)
D12S1721	0.005	2.17	4.23 (0.06)
NBG8	0.011	2.67	2.79 (0.00)
NBG6	0.0115	3.79	5.06 (0.06)
NBG9	0.0035	4.95	1.57 (0.14)
NBG10	0.001	5.30	1.73 (0.00)
NBG12	0.009	5.40	1.65 (0.12)
NBG4	0.001	6.31	4.60 (0.08)
NBG3	0.006	6.41	4.84 (0.06)
NBG2		7.01	2.80 (0.06)

Model-free LOD score results obtained with ANALYZE and ASPEX applied to the strata are shown in Table 7.

**Table 7. Model-free LOD score obtained with ANALYZE and ASPEX**

Locus	Distance (q)	Cumulative distance (cM)	ANALYZE LOD score	sib_phase LOD score independent sib-pairs	sib_phase LOD score all sib-pairs
D12S1619	0.0135	0.00	4.54	5.29	7.65
NBG11	0.006	1.37	4.29	5.34	7.70
D12S1666	0.001	1.97	2.77	5.36	7.74
NBG5	0.001	2.07	0.67	5.36	7.74
D12S1721	0.005	2.17	4.48	5.35	7.74
NBG8	0.011	2.67	2.97	4.87	7.00
NBG6	0.0115	3.79	4.05	4.59	6.76
NBG9	0.0035	4.95	2.03	3.72	5.41
NBG10	0.001	5.30	2.00	3.42	5.89
NBG12	0.009	5.40	0.89	3.44	6.11
NBG4	0.001	6.31	2.84	3.71	9.00
NBG3	0.006	6.41	3.89	3.71	9.01
NBG2		7.01	1.91	3.52	8.73

Model-free results observed with SIMWALK2 are illustrated in Table 8.

**Table 8. Model-free LOD score obtained with SIMWALK2**

Locus	Distance (q)	Cumulative Distance (cM)	STAT(A) -Log(p-value)	STAT(B) -Log(p-value)	STAT(C) -Log(p-value)	STAT(D) -Log(p-value)
D12S1619	0.0135	0.00	2,5963	0,9565	3,2156	2,3584
NBG11	0.006	1,37	3,0698	1,7400	3,7747	3,0103
D12S1666	0.001	1,97	2,9340	1,6546	3,5812	2,7223
NBG5	0.001	2,07	2,9781	1,2722	3,6505	2,7846
D12S1721	0.005	2,17	2,9680	1,2630	3,6844	2,7752
NBG8	0.011	2,67	3,0954	1,0804	3,4399	2,5654
NBG6	0.0115	3,79	3,1632	1,0672	3,2670	2,5956
NBG9	0.0035	4,95	2,2106	1,0137	2,7765	2,4456
NBG10	0.001	5,30	2,5513	1,0251	2,7625	2,1914
NBG12	0.009	5,40	2,4893	0,9868	2,6841	2,0920
NBG4	0.001	6,31	2,9028	1,1312	3,4063	2,8156
NBG3	0.006	6,41	2,9070	1,1326	3,4637	2,8300
NBG2		7,01	2,8430	1,1108	3,3135	2,7978

Multipoint results on the strata with ASPEX sib\_phase by considering only independent sib-pairs (Figure 1b) or all sib-pairs (Figure 1c) are shown in Figures 1d and 1e. As previously reported a second peak appeared when all sib-pairs were observed.

A confidence interval was calculated. GENEFINDER (Liang et al., Am. J. Hum. Genet. 66 (2000), 1631-1641) was used to estimate the location of the susceptibility gene (say  $t$ ). The method is based on the IBD (Identity by Descent) sharing of affected sib-pairs for multiple markers. For the purpose of our analysis, pedigrees were divided into sibship. 56 nuclear families and 183 sib-pairs were used. Liang KY, Huang CY, Beaty TH (2000) A unified sampling approach for multipoint analysis of qualitative and quantitative traits in sib pairs. Am J Hum Genet 66:1631-1641

The GENEFINDER results points to localization of a susceptibility gene for affective disorders at  $3.19 \pm 0.446$  cM telomeric to the marker D12S1721 (D12S1721 is approximately located at 136.82 cM on the sex-averaged Marshfield chromosome 12 map).

95% C.I.: [2.32, 4.06];

99% C. I.: [2.03, 4.35]

99.9% C. I.: [1.71, 4.67]

From the strata, 24 nuclei, and 107 sib-pairs were obtained, and the location of the susceptibility gene was estimated at  $3.07 \pm 0.57$  (see map above). The following confidence interval (C.I.) was obtained:

95% C.I.: [1.95, 4.19];

99% C. I.: [1.59, 4.55]

99.9% C. I.: [1.18, 4.96]

An association study using the NBS microsatellite markers was done with CLUMP (Sham & Curtis, Ann. Hum. Genet. 59 (1995), 97-105). Samples were distributed as follow: 83 male/case; 124 female/case; 95 male/control; and 101 female/control. One thousand simulations were used to estimate p-values. The observed results are summarized in Table 9.

**Table 9. Association study using the NBG microsatellite**

<b>Locus</b>	<b>Sample</b>		<b>T1statistic (p-value)</b>	<b>T2statistic (p-value)</b>	<b>T3statistic (p-value)</b>	<b>T4statistic (p-value)</b>
	<b>Case</b>	<b>Control</b>				
NBG11	204	129	0.226	0.562	0.410	0.421
NBG5	206	194	0.972	0.980	0.948	0.971
NBG8	206	194	0.983	1.000	0.994	0.978
NBG6	206	194	0.147	0.074	0.759	0.485
NBG9	206	190	0.512	0.940	0.786	0.583
NBG10	206	190	0.594	0.480	0.403	0.709
NBG12	206	190	0.002	0.019	0.003	0.117

T1 statistic is the usual chi-squared statistic on the raw contingency table

T2 statistic is the usual chi-squared statistic apply on the contingency table obtained after collapsing columns with small expected values together

T3 statistic is the largest chi-squared statistic got by comparing one column of the original table against the total of the others columns

T4 statistic is the largest chi-squared statistic got by comparing any combination of alleles against the rest.

Only the NBG12 marker gave significant association at the 1% level. For the others markers, there was no single alleles that seems to be associated with bipolar disorder. It seems that no founder-alleles was overly represented among the affected. There is no significant result for association of genotypes with the NBG markers.

In conclusion, the parametric and model-free multipoint results suggest to investigate genes located between D12S1619 and D12S1666. Moreover, according to GENEFINDER results, genes situated centromeric to NBG9 should be considered for association and linkage disequilibrium analysis.

## EXAMPLE 2

### Physical mapping and Mutation analysis of chromosome 12 associating the P2X7R to Bipolar Affective Disorders

The most conservative prediction for the disease-associated region is included between markers NBG11 and NBG2 (see Figure 1a). This region was delimited according to linkage and association analysis described in Example 1, using genethon markers and NBG markers. The approximate length of this region is 5,2 Mb. Two major gaps (between FLJ10701 and FLJ32372, and between FLJ1466 and

MONDOA) were included in this region. At least 73 genes were listed in this area, where 48 are known genes and 25 are unknown but associated to mRNA and/or EST clusters based on the last genome assembly available at UCSC (November 2002). Predicted genes were not listed. However, the estimation of CI 99% (confidence interval) using GENEFINDER has limited the most interesting region between markers D12S1666 and NBG9. This genomic region covers 1,6 Mb and includes at least 28 genes, and has no major gap. Thus, the term fBAD (familial Bipolar Affective Disorders) region was used to describe the genomic segment between D12S1666 and NBG9. Genes found within this region include CaMKK2, CABP, P2X7, P2X4, PIN, PLA2, G1B, CIT, PXN, Rab35, and APC5. However, given the present art, it would not have been obvious to an ordinary person skilled in the art to select P2X7R as the gene associated with affective diseases. Other genes from the ones listed above would be obvious.

For example, the CaMKK2 gene (also known as Ca<sup>2+</sup>/Calmodulin-dependent protein kinase kinase beta, or CaMKKb) is a serine/threonine protein kinase involved in Ca<sup>2+</sup> dependent signalling pathways. CaMKK2 can activate in vitro the downstream kinases CaMKIV and CaMKI, which modulate gene transcription through phosphorylation of transcription factors (e.g., CREB, SRF, MEF2; Corcoran and Means, J. Biol. Chem. 276 (2001), 2975-2978; Soderling, Trends Biochem. Sci. 24 (1999), 232-236). Its role in the Ca<sup>2+</sup> cascade is not critical. Some studies suggest that CaMKs could be activated without the CaMKKs phosphorylation (Matsushita and Nairn, J. Biol. Chem. 274 (1998), 10086-10093). However, CaMKK phosphorylation step would contribute to amplification of the Ca<sup>2+</sup> signal since CaMKK is more sensitive to activation by Ca<sup>2+</sup>/Calmodulin, therefore CaMKK would be an important mediator when the levels of intracellular Ca<sup>2+</sup> are low (Anderson et al., J. Biol. Chem. 273 (1998), 31880-31889).

CaMKK2 is an obvious target for depression since prior art suggest that cAMP-dependent signaling pathways (mediated by PKA activation) is affected in brain from patients with Bipolar Affective Disorders (Field et al., J. Neurochem. 73 (1997), 1704-1710; Rahman et al., J. Neurochem. 68 (1997), 297-304; Takahashi et al., J. Neurosci. 19 (1999), 610-618). According to a study using lymphoblastic cell lines,

Bipolar disorder could be related to elevated intracellular calcium levels (Yoon et al., *Mol. Psychiatry* 6 (2001), 678-683). Moreover, some groups found relations between antidepressant drugs and CaMK activation (Budziszewska et al., *Br. J. Pharmacol.* 130 (2000), 1385-1393; Consogno et al., *Neuropsychopharmacology* 24 (2001), 21-30; Mori et al., *Neuropharmacology* 40 (2001), 448-456; Zanotti et al., *Neuropharmacology* 37 (1998), 1081-1089). Furthermore, inhibition of CaMKK by PKA-mediated phosphorylation suggest a close relationship between both pathways (Matsushita et al., *J. Biol. Chem.* 273 (1999), 21473-21481). These observations would suggest to a person skilled in the art that CaMKK2 is the gene responsible for bipolar affective disease.

Another obvious candidate for affective disorders would have been the CABP1 gene which generates four neuronal  $\text{Ca}^{2+}$ -binding protein by alternative usage of the 9 coding exons, which are L-CABP, S-CABP, calbrain, and caldendrin (Haeseleer et al., *J. Biol. Chem.* 275 (2000), 1247-1260). Their expression is almost totally restricted to brain tissues. A functional study on calbrain reveals its negative effect on  $\text{Ca}^{2+}$ /Calmodulin-dependent CaMKII activity by competitively interacts with the CaM-binding domain of CaMKII (Yamagushi et al., *J. Biol. Chem.* 274 (1999), 3610-3616). One would expect similar roles in  $\text{Ca}^{2+}$  signaling for other CABP1 alternative products. Participation of CABP1 gene in  $\text{Ca}^{2+}$ -dependent signaling pathways would make it obvious to one skilled in the art to select this gene as a candidate for bipolar affective disorder. However, all CABP1 exons were analyzed for the presence of mutations, and surprisingly only two mutations were detected in noncoding regions.

The PIN gene (Protein inhibitor of NOS (Nitric oxide synthase)) is another obvious candidate responsible for bipolar affective disorder. Nitric oxide (NO) in the brain, may be involved in apoptosis, synaptogenesis, and neuronal development. Because NO cannot be stored in vesicles like other neurotransmitters, its release is regulated by the activity of NOS (Nitric oxide synthase). PIN is a direct inhibitor of NOS by binding and destabilizing the active homodimer complex of NOS (Jaffrey et al., *Science* 274 (1996), 774-777). PIN is highly conserved throughout the evolution and is expressed in many cell types. A recent clinical study evaluating plasma nitrate levels in depressive states suggests that NO production is increased in depression

(Suzuki et al., J. Affect. Disord. 63 (2001), 221-224) and may result from a deficiency in NOS inhibition. Moreover in a mouse model, NO synthase antagonists have been linked to antidepressant properties (Harkin et al., 1999; Karolewicz et al., Eur. J. Pharmacol. 372 (1999), 215-220). Thus, PIN would be an obvious candidate. However, due to the pleiotropic action of NO, a deficiency in PIN function would generate many unrelated disorders throughout the body. Thus, without the information presented in the disclosure herein, a person of ordinary skills in the art would have predicted PIN and not P2X7R as the gene associated with affective disorders.

The human phospholipase A2 group IB (PLA2G1B) catalyses the release of fatty acids from glycerol-3-phosphocholines. Phospholipase A2 genes (PLA2) are expressed in many tissues. Some studies have demonstrated associations between excessive PLA2 activity in brain and affective disorders (Chang et al., Neurochem. Res. 23 (1998), 887-892; Hibbeln et al., Biol. Psychiatry 25 (1989), 945-961). Moreover, other genetic studies have found associations between PLA2G1B gene and bipolar affective disorder (Dawson et al., Psychiatr. Genet. 5 (1995), 177-180). Thus, PLA2G1B represent a likely candidate for affective disorders. However in the present example, only a single silent mutation was found within exon 3 of the PLA2G1B gene.

The human citron kinase gene, Rho-associated protein (CIT) is a 183 kDa protein which associates to the GTPase Rho. CIT shares strong similarity with ROCK and ROK proteins which are other Rho-associated kinases (Madaule et al., Nature 394 (1998), 491-494). Rho GTPases are involved in many processes such as cytoskeletal organization, membrane trafficking, cell growth, and transcriptional activation (Van Aelst and D'Souza-Schorey, Genes Dev. 11 (1997), 2295-2322). Studies on brain variants of Citron-K (without the kinase domain) reveal the association with postsynaptic density proteins (PSD-95), suggesting a role in either synapse organization or function (Zhang et al., J. Neurosci. 19 (1999), 96-108; Furuyashiki et al., J. Neurosci. 19 (1999), 109-118).

The human paxillin (PXN) gene encodes for a 68 kDa protein found in focal adhesions. It is within focal adhesions where adhesion molecules dynamically

interact with the cytoskeleton (Salgia et al., J. Biol. Chem. 270 (1995), 5039-5047). The signaling pathways that regulate these dynamic interactions begin to be elucidated. Many observations suggest that paxillin is involved in transducing signals from growth factor receptors to focal adhesions. The paxillin is expressed in many tissues including brain.

However as set forth below, the gene causative for affective diseases is identified as being the P2X7 receptor (P2X7R).

Mutations were searched in coding sequences and exon-intron boundaries of the above mentioned genes since such mutations are more likely to give a functionally significant Single Nucleotide Polymorphisms (SNP). The starting sample was composed of 16 unrelated affected individuals from the Saguenay/Lac St-Jean region, which gives an 80% power to detect polymorphisms with a frequency of 0.05. To identify polymorphisms, targeted sequences were first amplified by PCR. Then, PCR products are purified on Whatman GF/C membranes (VWR, Montreal, Canada), and quantified using the PicoGreen dsDNA quantitation assay (Molecular probes, Oregon, USA). 4 ng of purified PCR products are sequenced using the DYEnamic ET terminator cycle sequencing kit (Amersham Biosciences, Baie D'Urfé, Canada). The sequencing products are resolved on an ABI PRISM 3730XL DNA analyzer, and an ABI PRISM 3700 DNA analyzer. The PCR products are sequenced in both directions. The SNPs identified in studied genes are listed in Table 10.

**Table 10. Mutation analysis between markers D12S1666 and NBG9**

Genes	Positions	Variations	Alleles	Modifications
Rab35	Exon06	RABE06A	486G-A	Silent Asn162
Rab35	Intron04	RABI04A	51C-T	unknown
Rab35	Intron03	RABI03A	33G-A	unknown
Rab35	Intron02	RABI02B	85G-A	unknown
Rab35	Intron02	RABI02A	76C-G	unknown
PXN	Exon11	PXNE11A	1527C-T	Silent Thr509
PXN	Exon06	PXNE06A	750C-T	Silent Ser250
PXN	Exon02	PXNE02A	217G-A	Gly73Ser
PLA2G1B	Exon03	PLA2G1BE03A	294C-T	Silent Ser98
PIN	5'UTR01	PINUTR01A	-49T-G	unknown
PIN	5'UTR01	PINUTR01B	-80T-C	unknown
PIN	Intron02	PINI02A	26C-T	unknown



PIN	Intron02	PIN102B	50C-T	unknown
CaBP	Intron04	CaBPI04A	35C-T	unknown
CaBP	exon01	CaBPE01A	-23A-G	unknown
OASL	Exon02	OASLE02A	213G-T	Silent Gly72
OASL	Exon02	OASLE02B	408C-T	Silent Leu136
OASL	Exon05	OASLE05A	1042G-A	Val348Met
OASL	Exon06	OASLE06A	1509G-A	Silent Ser503
P2X7R	5'UTR	P2XR7UTR5L	362T-C	unknown
P2X7R	5'UTR	P2XR7UTR5M	532T-G	unknown
P2X7R	5'UTR	P2XR7UTR5K	1100A-G	unknown
P2X7R	5'UTR	P2XR7UTR5J	1122A-G	unknown
P2X7R	5'UTR	P2XR7UTR5I	1171C-G	unknown
P2X7R	5'UTR	P2XR7UTR5F	1351T-C	unknown
P2X7R	5'UTR	P2XR7UTR5N	1702G-A	unknown
P2X7R	5'UTR	P2XR7UTR5G	1731T-G	unknown
P2X7R	5'UTR	P2XR7UTR5H	1860C-T	unknown
P2X7R	5'UTR	P2XR7UTR5A	2162C-A	unknown
P2X7R	5'UTR	P2XR7UTR5B	2238C-T	unknown
P2X7R	5'UTR	P2XR7UTR5D	2373A-G	unknown
P2X7R	5'UTR	P2XR7UTR5E	2569G-A	unknown
P2X7R	5'UTR	P2XR7UTR5C	2702G-A	unknown
P2X7R	Intron01	P2XR7I01C	3166G-C	unknown
P2X7R	Intron01	P2XR7I01A	24778C-T	unknown
P2X7R	Intron01	P2XR7I01B	24830C-T	unknown
P2X7R	Exon02	P2XR7v02A	24942T-C	Val176Ala
P2X7R	Exon03	P2XR7E03A	26188C-T	Arg117Trp
P2X7R	Intron03	P2XR7I03A	26308A-G	unknown
P2X7R	Intron03	P2XR7I03B	26422G-A	unknown
P2X7R	Intron04	P2XR7I04A	32394G-A	unknown
P2X7R	Intron04	P2XR7v05B	32434T-C	unknown
P2X7R	Exon05	P2XR7E05D	32493G-A	Gly150Arg
P2X7R	Exon05	P2XR7v05A	32507C-T	Tyr155His
P2X7R	Exon05	P2XR7E05C	32783C-T	Silent Cys168
P2X7R	Intron05	P2XR7I05C	32783A-C	unknown
P2X7R	Intron05	P2XR7I05D	35309T-C	unknown
P2X7R	Intron05	P2XR7I05B	35374C-T	unknown
P2X7R	Intron05	P2XR7I05A	35378A-C	unknown
P2X7R	Exon06	P2XR7E06A	35438G-A	Glu186Lys
P2X7R	Exon06	P2XR7E06B	35454T-C	Leu191Pro
P2X7R	Intron06	P2XR7I06C	35549T-C	unknown
P2X7R	Intron06	P2XR7I06G	35641G-C	unknown
P2X7R	Intron06	P2XR7I06D	35725A-C	unknown
P2X7R	Intron06	P2XR7I06F	36001T-G	unknown
P2X7R	Intron06	P2XR7I06E	36064A-T	unknown
P2X7R	Intron06	P2XR7I06A	36091DelGTTT	unknown
P2X7R	Intron06	P2XR7I06B	36108C-G	unknown
P2X7R	Intron07	P2XR7I07A	36374C-T	unknown
P2X7R	Intron07	P2XR7I07B	36378G-A	unknown
P2X7R	Intron07	P2XR7I07C	36387T-A	unknown
P2X7R	Intron07	P2XR7I07D	36398G-C	unknown
P2X7R	Intron07	P2XR7I07E	37439C-T	unknown
P2X7R	Intron07	P2XR7I07F	37513T-C	unknown

P2X7R	Exon08	P2XR7E08C	37604C-T	Arg270Cys
P2X7R	Exon08	P2XR7v08A	37605G-A	Arg270His
P2X7R	Exon08	P2XR7v08B	37623G-A	Arg276His
P2X7R	Exon08	P2XR7E08D	37633C-T	Silent Asp279
P2X7R	Intron09	P2XR7v11A	47214C-T	unknown
P2X7R	Exon11	P2XR7v11B	47383G-A	Ala348Thr
P2X7R	Exon11	P2XR7v11C	47411C-G	Thr357Ser
P2X7R	Intron11	P2XR7I11D	47563T-C	unknown
P2X7R	Intron12	P2XR7I12A	54307C-T	unknown
P2X7R	Intron12	P2XR7I12B	54308G-A	unknown
P2X7R	Exon13	P2XR7v13F	54399C-T	Ala433Val
P2X7R	Exon13	P2XR7v13A	54480A-G	Gln460Arg
P2X7R	Exon13	P2XR7v13B	54523C-T	Silent Pro474
P2X7R	Exon13	P2XR7v13G	54562DelCCCTGAGAG CCACAGGTGCCT	Del of 7aa 488 to 494 PESHRCL
P2X7R	Exon13	P2XR7v13C	54588A-C	Glu496Ala
P2X7R	Exon13	P2XR7v13H	54664C-G	Silent His521
P2X7R	Exon13	P2XR7E13D	54703G-T	Silen Leu534
P2X7R	Exon13	P2XR7E13J	54804A-T	Ile568Asn
P2X7R	Exon13	P2XR7v13I	54834G-A	Arg578Gln
P2X7R	Exon13	P2XR7v13E	54847G-A	Silent Pro582
P2X7R	3'UTR	P2XR7UTR3A	55169C-A	unknown
P2X7R	3'UTR	P2XR7UTR3B	55170A-C	unknown
P2X7R	3'UTR	P2XR7UTR3C	55171A-C	unknown
P2X7R	3'UTR	P2XR7UTR3D	55917C-T	unknown
P2X7R	3'UTR	P2XR7UTR3E	54925G-A	unknown
P2X4R	5'UTR	P2XR4UTR5I	-1956G-A	unknown
P2X4R	5'UTR	P2XR4UTR5H	-1649G-A	unknown
P2X4R	5'UTR	P2XR4UTR5G	-800G-A	unknown
P2X4R	5'UTR	P2XR4UTR5A	-648C-A	unknown
P2X4R	5'UTR	P2XR4UTR5B	-537A-G	unknown
P2X4R	5'UTR	P2XR4UTR5C	-437A-G	unknown
P2X4R	5'UTR	P2XR4UTR5J	-206VNRG	unknown
P2X4R	5'UTR	P2XR4UTR5D	-211C-G	unknown
P2X4R	5'UTR	P2XR4UTR5F	-150VNRGGGGCCCC	unknown
P2X4R	5'UTR	P2XR4UTR5E	-98G-T	unknown
P2X4R	Intron01	P2XR4I01A	31G-T	unknown
P2X4R	Exon02	P2XR4E02A	262G-A	Silent mutation Ala87
P2X4R	Intron02	P2XR4I02A	4600C-T	unknown
P2X4R	Intron03	P2XR4I03A	15G-A	unknown
P2X4R	Intron03	P2XR4I03B	72G-A	unknown
P2X4R	Exon04	P2XR4E04A	355G-A	Ile119Val
P2X4R	Exon04	P2XR4E04A	375G-A	Silent Val125
P2X4R	Intron04	P2XR4I04B	17T-C	unknown
P2X4R	Intron04	P2XR4I04A	32G-A	unknown
P2X4R	Exon05	P2XR4E05A	465T-C	Silent Ser155
P2X4R	Exon07	P2XR4E07A	724A-G	Ser242Gly
P2X4R	Intron08	P2XR4I08A	DelT	unknown
P2X4R	Exon09	P2XR4E09A	944A-G	Tyr315Cys
P2X4R	Intron10	P2XR4I10A	11G-T	unknown
P2X4R	Intron10	P2XR4I10B	G-C	unknown
P2X4R	Intron10	P2XR4I10C	A-G	unknown

P2X4R	Intron11	P2XR4I11B	C-G	unknown
P2X4R	Intron11	P2XR4I11C	T-A	unknown
P2X4R	Intron11	P2XR4I11A	374C-T	unknown
CaMKK2	3'UTR	CaMKK2UTR3bA	733C-T	unknown
CaMKK2	3'UTR	CaMKK2UTR3aB	390G-A	unknown
CaMKK2	3'UTR	CaMKK2UTR3aA	239G-A	unknown
CaMKK2	Intron15	CaMKK2I15B	325T-C	unknown
CaMKK2	Intron15	CaMKK2I15A	169G-A	unknown
CaMKK2	Intron14	CaMKK2I14A	224A-G	unknown
CaMKK2	Intron10	CaMKK2I10A	156DelGTGATCCGCCT G	unknown
CaMKK2	intron09	CaMKK2I09B	528A-G	unknown
CaMKK2	intron09	CaMKK2I09A	521A-G	unknown
CaMKK2	Exon09	SNP6f18v5	1095C-A	Silent Ile365
CaMKK2	Exon09	SNP6f18v4	1087C-T	Arg363Cys
CaMKK2	Exon05	CaMKKE05A	687C-T	Silent Pro229
CaMKK2	Intron03	CaMKK2I03A	10C-T	unknown
CaMKK2	Intron02	CaMKK2I02A	39C-T	unknown
CaMKK2	Intron01	CaMKK2I01B	2911G-C	unknown
CaMKK2	Intron01	CaMKK2I01A	89C-A	unknown
CaMKK2	Exon01	SNP6f18v2	253A-T	Thr85Ser
CaMKK2	Exon01	SNP6f18v1	29G-A	Ser10Asn
CaMKK2	5'UTR01	CaMKK2UTR01B	253T-C	unknown
CaMKK2	5'UTR01	CaMKK2UTR01A	63C-A	unknown
APC5	Intron01	APC5I01A	10G-T	unknown
APC5	Intron01	APC5I01B	50A-T	unknown
APC5	Intron05	APC5I05A	73T-C	unknown
APC5	Intron06	APC5I06A	73T-G	unknown
APC5	Exon11	APC5E11A	1416C-T	Silent His472

Each SNP in genes Rab35, PXN, PLA2G1B, PIN, CaBP, OASL, P2X4R, CaMKK2 and APC5 was designated according to the gene where it was found, and its location in that gene (intronic or exonic regions). Each SNP in the P2X7R gene was designated according to their position on SEQ ID NO: 1. The allele describes the position and the variation observed. In coding regions, the position is relative to the start codon, whereas the intronic SNPs are positioned relative to the beginning of the corresponding intron (when known). Primers used for identifying the SNPs in the P2X7R and the location of each SNPs included in tables 2 and 12 are defined in table 1a and SEQ ID NOs 52 to 111.

Association studies using missense SNPs were performed. Missense SNPs or SNPs that could be close to the splice sites were used, because it is more likely that

diseases would be associated to an improper function in proteins. Case group was composed by bipolar I individuals, schizoaffective bipolar type (182 subjects) and bipolar II diagnosed persons (31 subjects). Many controls from the Saguenay/Lac-St-Jean region, were sampled from Steinert, Glaucoma and Paget DNA banks. The control individuals were not diagnosed for affective disorders. According to the lifetime risks of bipolar disorders (1%), there is no need to screen controls for psychiatric disorders.

Direct sequencing of PCR products is by far the most accurate method of analysis and is the method of choice in view of our sequencing platform capacity. PCR products were analyzed by direct sequencing as described above. After sequencing analysis, individuals are automatically typed for the corresponding SNP using a home-developed program, GENO.pl. The results of SNP genotyping are compiled in a 4D database.

The association hypothesis was tested with CLUMP (Sham & Curtis 1995, Ann. Hum. Genet. 59:97-105). One thousand simulations were used to estimate p-values. Results are illustrated in table 11. The T1 statistic, which is the usual chi-squared statistic on the raw contingency table, was used to test for allelic association. Moreover, the largest chi-squared statistic got by comparing one column of the original table against the total of the other columns, called T3 statistic, was added to the previous one to test for potential genotype association since T1 statistic results may be biased when the contingency table contains cells with low values.

**Table 11. Association hypothesis using CLUMP**

gene	SNPs	Effective		Allele Analysis p-value (T1)	Genotype Analysis	
		Cases	Controls		p-value (T2)	p-value (T3)
P2X7R	P2XR7v11B	208	211	0.795	0.036	0.028
	P2XR7v13A	212	214	0.344	0.250	0.186
	P2XR7v13E	212	211	0.780	0.017	0.017
CAMKK2	SNP6f18v5	206	135	1.00	1.00	1.00
	SNP6f18v4	206	135	0.816	0.962	0.841
	SNP6f18v2	205	135	0.057	0.110	0.095
	SNP6f18v1	206	135	0.512	0.532	0.385

The association studies using SNPs in P2X7, P2X4, and CaMKK2 reveal associations significant at level of about 5% or less. Three genotype associations in P2X7 were observed. However, SNPs P2XR7v11B and P2XR7v13E are closely linked together based on a contingency table. There is also an allele association at level of 5,7% for SNP6f18v2 in CaMKK2. The information associated to each relevant SNP can be found in Tables 10 and 12.

### EXAMPLE 3

#### Polymorphisms found in the P2X7R in individuals suffering from depression

Association studies using SNPs in the P2X7R gene was performed in a case/control sample (535 individuals) from a German population. The case group was composed of 36 individuals diagnosed with bipolar type I or type II, and 279 individuals diagnosed with unipolar disorders (i.e. depression) representing 133 affected males and 182 affected females. Among controls, we count The remaining 220 control individuals were normal (i.e. diagnosed as non depressive), and comprising 81 males, 182 females and 14 of unknown gender. The same sexual distribution was noted in both groups.

SNPs were identified in this sample by using a subgroup of 24 affected individuals. SNPs in the P2X7R gene detected in the German population were similar if not identical to the SNPs seen in the Saguenay/Lac-St-Jean population (see table 12).

Other rare missense SNPs were also noted in the German population, such as Arg117Trp (P2XR7E03A), Glu186Lys (P2XR7E06A), Leu191Pro (P2XR7E06B), Ile568Asn (P2XR7E13J). These amino acids are quite conserved between ortholog P2X7 genes. It is possible that the Ile568Asn (P2XR7E13J) mutation may be involved in the surface expression of P2X7.

**Table 12. Comparison between polymorphisms in the Saguenay/Lac-St-Jean population and the German population in the human P2X7R gene**

Associated exons or introns	Variation (SNP or others)	Allele	Position*	Modification	Frequency (Canada)	Frequency (Germany)
5'UTR	P2XR7UTR5L	T-C	362	unknown	0,13	0,08
5'UTR	P2XR7UTR5M	T-G	532	unknown	0,16	0,1
5'UTR	P2XR7UTR5K	A-G	1100	unknown	0,13	0,13
5'UTR	P2XR7UTR5J	A-G	1122	unknown	0,13	0,13
5'UTR	P2XR7UTR5I	C-G	1171	unknown	0,06	0,02
5'UTR	P2XR7UTR5F	T-C	1351	unknown	0,3	0,12
5'UTR	P2XR7UTR5N	G-A	1702	unknown	-	0,02
5'UTR	P2XR7UTR5G	T-G	1731	unknown	0,17	0,15
5'UTR	P2XR7UTR5H	C-T	1860	unknown	0,07	0,15
5'UTR	P2XR7UTR5A	C-A	2162	unknown	0,07	0,12
5'UTR	P2XR7UTR5B	C-T	2238	unknown	0,3	0,27
5'UTR	P2XR7UTR5D	A-G	2373	unknown	0,07	0,12
5'UTR	P2XR7UTR5E	G-A	2569	unknown	0,1	0,02
5'UTR	P2XR7UTR5C	G-A	2702	unknown	0,31	0,27
Intron01	P2XR7I01C	G-C	3166	unknown	0,03	-
Intron01	P2XR7I01A	C-T	24778	unknown	0,03	-
Intron01	P2XR7I01B	C-T	24830	unknown	0,03	RARE
Exon02	P2XR7v02A	T-C	24942	Val76Ala	0,06	0,08
Exon03	P2XR7E03A	C-T	26188	Arg117Trp	-	RARE
Intron03	P2XR7I03A	A-G	26308	unknown	0,7	0,44
Intron03	P2XR7I03B	G-A	26422	unknown	0,18	0,12
Intron04	P2XR7I04A	G-A	32394	unknown	0,03	0,01
Intron04	P2XR7v05B	T-C	32434	unknown	0,33	0,29
Exon05	P2XR7E05D	G-A	32493	Gly150Arg	RARE	0,02
Exon05	P2XR7E05E	G-A	32506	Silent Val154	-	RARE
Exon05	P2XR7v05A	C-T	32507	Tyr155His	0,33	0,44
Exon05	P2XR7E05C	C-T	32548	Silent Cys168	RARE	0,02
Intron05	P2XR7I05C	A-C	32783	unknown	0,25	-
Intron05	P2XR7I05D	T-C	35309	unknown	ND	0,35
Intron05	P2XR7I05B	C-T	35374	unknown	0,7	0,67

Intron05	P2XR7I05A	A-C	35378	unknown	0,7	0,65
Exon06	P2XR7E06A	G-A	35438	Glu186Lys	-	0,02
Exon06	P2XR7E06B	T-C	35454	Leu191Pro	-	0,02
Intron06	P2XR7I06C	T-C	35549	unknown	0,04	0,08
Intron06	P2XR7I06G	G-C	35641	unknown	-	0,02
Intron06	P2XR7I06D	A-C	35725	unknown	0,21	0,27
Intron06	P2XR7I06F	T-G	36001	unknown	0,17	0,3
Intron06	P2XR7I06E	A-T	36064	unknown	0,11	0,1
Intron06	P2XR7I06A	DelGTTT	36091-36094	unknown	0,14	0,3
Intron06	P2XR7I06B	C-G	36108	unknown	0,14	0,29
Intron07	P2XR7I07A	C-T	36374	unknown	0,07	-
Intron07	P2XR7I07B	G-A	36378	unknown	0,21	0,28
Intron07	P2XR7I07C	T-A	36387	unknown	0,21	0,28
Intron07	P2XR7I07D	G-C	36398	unknown	0,42	0,4
Intron07	P2XR7I07E	C-T	37439	unknown	0,41	-
Intron07	P2XR7I07F	T-C	37513	unknown	-	RARE
Exon08	P2XR7E08C	C-T	37604	Arg270Cys	RARE	-
Exon08	P2XR7v08A	G-A	37605	Arg270His	0,46	0,24
Exon08	P2XR7v08B	G-A	37623	Arg276His	0,03	0,02
Exon08	P2XR7E08D	C-T	37633	Silent Asp279	RARE	-
Intron09	P2XR7v11A	C-T	47214	unknown	0,08	0,03
Exon11	P2XR7v11B	G-A	47383	Ala348Thr	0,5	0,44
Exon11	P2XR7v11C	C-G	47411	Thr357Ser	0,08	0,07
Intron11	P2XR7I11D	T-C	47563	unknown	0,43	0,44
Intron12	P2XR7I12A	C-T	54307	unknown	0,32	-
Intron12	P2XR7I12B	G-A	54308	unknown	0,03	-
Exon13	P2XR7v13F	C-T	54399	Ala433Val	0,13	-
Exon13	P2XR7v13A	A-G	54480	Gln460Arg	0,13	0,17
Exon13	P2XR7v13B	C-T	54523	Silent Pro474	0,1	0,07
Exon13	P2XR7v13G	DelCCCTGAGA GCCACAGG TGCCT	54562-54582	Del of 7aa 488 to 494 (PESHRCL)	RARE	-
Exon13	P2XR7v13C	A-C	54588	Glu496Ala	0,13	0,06
Exon13	P2XR7v13H	C-G	54664	His521Gln	0,03	-
Exon13	P2XR7E13D	G-T	54703	Silent Leu534	0,1	0,02
Exon13	P2XR7E13J	A-T	54804	Ile568Asn	-	0,01
Exon13	P2XR7v13I	G-A	54834	Arg578Gln	-	RARE
Exon13	P2XR7v13E	G-A	54847	Silent Pro582	0,4	0,45
3'UTR	P2XR7UTR3A	C-A	55169	unknown	0,48	0,37
3'UTR	P2XR7UTR3B	A-C	55170	unknown	0,09	0,1
3'UTR	P2XR7UTR3C	A-C	55171	unknown	0,05	0,06
3'UTR	P2XR7UTR3D	C-T	55917	unknown	0,001	-
3'UTR	P2XR7UTR3E	G-A	54925	unknown	-	0,01

The position and numbering of the polymorphism corresponds to the human P2X7R gene as defined in SEQ ID NO: 1. To identify the genomic organization of the P2X7R gene, BAC clones were firstly organized using known polymorphic markers, sequence tag sites (STSs), BAC-end sequences and expressed sequence tags (ESTs). Unorientated and unordered DNA regions were reassembled into a sequences using Phrap and reordered the pieces using P2X7R exons as scaffolds. No complete gene organization for P2X7R has been done. There is only a partial gene structure from exon6 to 13, NT\_037809. Therefore, this genomic sequence encompassing the P2X7R gene as depicted in SEQ ID NO: 1 could contain some sequence errors, specifically in intronic regions. Primers used for SNP amplification and sequencing are shown in Table 1a and depicted in SEQ ID NOs: 52 to 111.

Statistical analysis was performed according to the CLUMP method (Sham & Curtis 1995, Ann. Hum. Genet. 59:97-105). Table 13 resumes the allelic and genotypic association studies for SNPs in P2X7 gene.

**Table 13. Allelic and genotypic association studies using CLUMP**

Locus	Allele Frequencies*	Effective		Allele Analysis p-value (T1)	Genotype Analysis	
		Cases	Controls		p-value (T1)	p-value (T3)
P2XR7UTR5F	2(0.23); 4(0.77)	311	217	0.109	0.319	0.339
P2XR7UTR5N	1(0.001); 3(0.999)**	314	218	0.038	0.048	0.048
P2XR7UTR5G	2(0.001); 3(0.105); 4(0.894)	314	218	0.993	0.714	0.761
P2XR7UTR5H	2(0.92); 4(0.08)	312	215	0.743	0.884	0.754
P2XR7UTR5A	1(0.08); 2(0.92)	312	219	0.557	0.786	0.678
P2XR7UTR5B	2(0.73); 4(0.27)	310	218	0.485	0.761	0.814
P2XR7UTR5D	1(0.92); 3(0.08)	311	217	0.555	0.787	0.691
P2XR7v02A	2(0.09); 4(0.91)	313	218	0.501	0.729	0.591
P2XR7I04A	1(0.04); 3(0.96)	314	220	0.604	0.433	0.348
P2XR7v05B	2(0.69); 4(0.31)	314	220	0.133	0.270	0.325
P2XR7E05D	1(0.03); 3(0.97)	314	220	0.842	0.827	0.827
P2XR7E05E	1(0.006); 3(0.994)**	314	220	0.048	0.045	0.045
P2XR7v05A	2(0.60); 4(0.40)	314	220	0.038	0.144	0.219
P2XR7E05C	2(0.98); 4(0.02)	314	220	1.000	1.000	1.000
P2XR7I07F	2(0.002); 4(0.98)	315	219	1.000	1.000	1.000
P2XR7v08A	1(0.23); 3(0.77)	315	219	0.454	0.673	0.634
P2XR7v08B	1(0.02); 3(0.98)	315	219	0.636	0.638	0.638
P2XR7v11A	2(0.95); 4(0.05)	311	218	0.348	0.391	0.436
P2XR7v11B	1(0.45); 3(0.55)	312	218	0.605	0.803	0.790
P2XR7v11C	2(0.93); 3(0.07)	312	218	0.793	0.256	0.924
P2XR7I11D	2(0.45); 4(0.55)	312	219	0.665	0.735	0.740
P2XR7v13A	1(0.87); 3(0.13)	305	215	0.017	<0.001	<0.001



P2XR7v13B	2(0.93); 4(0.07)	305	216	1.000	0.228	0.677
P2XR7V13C	1(0.91); 2(0.09)	305	216	0.151	0.006	0.008
P2XR7E13D	3(0.94); 4(0.06)	315	219	0.402	0.429	0.474
P2XR7E13J	1(0.01); 4(0.99)	315	219	0.618	0.603	0.603
P2XR7E13I	1(0.004); 3(0.996)	315	219	0.999	1.000	1.000
P2XR7v13E	1(0.46); 3(0.54)	314	219	0.699	0.866	0.845
P2XR7UTR3A	1(0.518); 2(0.482)	314	219	0.617	0.850	0.875
P2XR7UTR3B	1(0.966); 2(0.034)	313	219	0.522	0.850	0.643
P2XR7UTR3C	1(0.979); 2(0.021)	313	219	0.636	0.505	0.382
P2XR7UTR3E	1(0.02); 3(0.98)	315	219	0.147	0.161	0.161

\* The column Allele Frequencies presents the allele for each SNP (A=1, C=2, G=3, T=4) and their respective frequency.

\*\* For this SNP we observed a zero cell in both (allele and genotype) 2X2 contingency tables. p-value <0.045 was observed exact Fisher test.

For the SNP analysis, the Hardy-Weinberg (HW) equilibrium was controlled in the control samples. The Hardy-Weinberg principle (HWP) may be stated as follow: In a large, randomly mating population, in which there is no migration, or selection against a particular genotype and the mutation rate remains constant, the proportions of the various genotypes will remain unchanged from one generation to another. Take a two allele system with alleles A and a. If the proportion of A in the population is represented as p and the proportion of a as q, then p plus q represent the sum total of alleles at this locus, that is  $p+q=1$ . The HWP is useful to evaluate some population problems like marital assortment, Inbreeding, population stratification, admixture, decreased viability of a particular genotype. The SNP P2XR7v13A did not respect the Hardy-Weinberg equilibrium.

The association hypothesis was also tested using an allele positivity table known to be suitable for the detection of susceptibility alleles showing a dominant mode of inheritance (Ohashi and Tokunaga, J. Hum. Genet. 44 (1999), 246-248; Ohashi et al., Ann. Hum. Genet. 65 (2001), 197-206). Similar results were obtained using this method as those obtained using the allele frequency tables, with the exception of P2XR7v05A where the p-values were 0.253. Thus, P2XR7v05A presented a less significant association in this analysis. This difference can be attributed to the mode of inheritance.

The proportion of unipolar individuals in analysis of the German population is quite important since the American Psychiatric Association (Diagnostic and Statistical Manual of Mental Disorders-4th Edition Text Revision (DMS-IV-TR), American Psychiatric Press, 2000) has reported an increase in susceptibility for unipolar disorders in female groups. To determine whether the sexual variable could influence the association analysis, additional association studies were performed by controlling the sexual parameter. Normal individuals in the German population without gender information were omitted from the study. Then, a logistic regression model was derived by including the sex as factor. In order to obtain a model that is as stable as possible, the regression model was minimised by using the difference between log-likelihood's for models with or without interaction (Hosmer, and Lemeshow, "Applied logistic regression", John Wiley and Sons, 1989). The strategy used for handling the zero cells from contingency tables was to eliminate associated category completely. Calculations were done with SAS v8.0 SAS is a statistical software package that allows the user to manipulate and analyze data in many different ways. Because of its capabilities, this software package is used in many disciplines, including medical sciences, biological sciences, and social sciences.

The introduction of a sexual parameter did not perturb the association already observed in previous analysis. Moreover, this analysis model revealed additional results: a potential allele association with P2XR7v05B ( $p=0.064$ ), and a genotypic association for P2XR7v08A ( $p=0.042$ ) was observed.

Association studies using pooled samples was performed by merging individuals from the samples of the Saguenay/Lac St-Jean with those of the German population. Results are illustrated in table 14. The aim of this analysis is to highlight common features between both populations. However, according to differences between both samples (mainly the phenotype of affected individuals i.e. bipolar disorder in the Saguenay/Lac St-Jean samples, versus mostly unipolar disorder in the German population) some parameters were controlled, including sex and ethnicity. The modelling strategy for logistic regressions was described above.

**Table 14. Association studies using pooled samples from both populations**

<b>Locus</b>	<b>Allele analysis</b>		<b>Genotype analysis</b>	
	p-value for SNP	p-value for sex	p-value for SNP	p-value for sex
P2XR7v02A	0.8254	0.0085	0.8650	0.4531
P2XR7v05B	0.1751	0.3714	0.2034	0.5110
P2XR7v05A	0.3808	0.0266	0.0885	0.1392
P2XR7v08A	0.0452	0.0041	0.1021	0.3452
P2XR7v08B	0.3471	0.0040	0.3413	0.3617
P2XR7v11A	0.3559	0.0136	0.5888	0.4404
P2XR7v11B	0.5902	0.0093	0.3897	0.4302
P2XR7v11C	0.3731	0.0094	0.7648	0.4615
P2XR7v13A	0.0047	0.0209	<0.0001	0.4814
P2XR7v13B	0.5129	0.2352	0.9584	0.4092
P2XR7v13C	0.2466	0.0284	0.2225	0.4228
P2XR7v13E	0.8168	0.0159	0.3713	0.4990

An allelic and genotypic association was observed for the P2XR7v13A locus ( $p=0.0047$ ) which was stronger than in the separate analyses. A significant allelic association was also noted for the P2XR7v08A locus ( $p=0.0452$ ). In addition, the present analysis also demonstrate the potential relationship between SNP P2XR7v05A and the origin with a  $p\text{-value}=0.0515$  (not shown in the table) which is in agreement with previous association analysis done in both samples separately (see Table 13).

The haplotype analysis was performed using the German population. The PHASE program (Stephens et al., Am. J. Hum. Genet. 68 (2001), 978-989) was used to estimate SNPs haplotypes within exons of the P2X7R gene. Haplotypes were created for each exon having more than one associated SNP (see Table 15 for exon-associated SNPs). Case groups varied from 218-220 individuals, whereas control groups varied between 312-316 individuals. Association hypothesis was tested with the CLUMP method since many haplotypes were created for each exon. T1 and T3 statistic tests performed as described above. T2 and T4 statistics were also calculated owing to the presence of small effective cells in the contingency tables. T2 statistic is the usual chi-squared statistic applied on the contingency table obtained after collapsing columns with small expected values. T4 statistic is the largest chi-squared statistic obtained by comparing one column of the original table against the total of the other columns. One thousand simulations were used to estimate p-values.

The resulting data was analyzed with the logistic regression model (describe above) using SAS V8.0 in order to consider the sexual parameter (for these tests the sample was reduced by 14 normal individuals). However, this analysis method is limited by the reliability of reconstructed haplotypes.

**Table 15. Exon-associated SNPs**

<i>Exons</i>	<i>Associated SNPs</i>
5	P2XR7E05D P2XR7E05E P2XR7v05A P2XR7E05C
8	P2XR7v08A P2XR7v08B
11	P2XRv11B P2XRv11C
13	P2XR7v13A P2XR7v13B P2XR7v13C P2XR7E13D P2XR7E13J P2XR7v13I P2XR7v13E

**Table 16. Genotypic association with haplotypes in exon 13 of P2X7R**

<i>Exon (haplotype)</i>	<i>Allele analysis</i>			<i>Genotype analysis</i>		
	Clump*	p-value(sex)	p-value(haplo)	Clump	p-value(sex)	p-value(haplo)
5(5)	T1:0.032 T2:0.068 T3:0.054 T4:0.059	0.3133	0.1947	T1:0.193 T2:0.159 T3:0.099 T4:0.304	0.460	0.5355
8(3)	T1:0.551 T2:0.585 T3:0.646 T4:0.646	0.3813	0.3064	T1:0.812 T2:0.689 T3:0.644 T4:0.756	0.5428	0.6652
11(3)	T1:0.750 T2:0.786 T3:0.726 T4:0.726	0.0886	0.7396	T1:0.625 T2:0.919 T3:0.929 T4:0.921	0.2305	0.9494
13(15**)	T1:0.088 T2:0.079 T3:0.147 T4:0.072	0.1871	0.1264	T1:0.001 T2:0.002 T3:0.057 T4:<0.001	0.4610	0.019

\*T1 test should not be considered because of contingency tables with zero cells.

\*\*Among these 15 haplotypes, we observed 8 haplotypes where case cells have less than 3 individuals.

Table 16 illustrates a genotypic association with haplotypes in exon 13 of the P2X7R genes. Interestingly, many haplotypes for the exon 13 were observed. The

differences between statistics in exon 13 (T3 less significant) can be explained by the involvement of more than one genotype of haplotypes in the disease. A potential allelic association was also noted with haplotypes in exon 5 of the P2X7R gene.

#### EXAMPLE 4

##### **P2X7R gene structure and mRNA expression and transcript sequence**

A 1700 bp nucleotide sequence corresponding to the human P2X7R promoter was analyzed by using MatInspector V2.2 and Transfac 4.0 algorithms. This analysis showed that the P2X7R gene does not contain a standard TATA box, but has SP1 sites that can make up for transcriptional initiation. Besides the SP1 sequences, there are binding sites for the transcription factors GATA, Oct and Ikarus. These sites are thought to provide tissue specificity. Interestingly, the P2X7R promoter has binding sites that suggest responsiveness to different cytokines such as AP-1, NFAT and CEBPB.

P2X7R possesses 13 exons and 12 introns (Buell et al., Receptors Channels 5 (1998), 347), providing a basis for alternative splicing that would yield in theory different transcripts and produce different isoforms with possible different functions. No alternatively spliced variant was clearly identified. However, experiments of EST clustering allowed the description of three splicing variants. One is defined by the lack of the exon 5. This P2X7v02 variant corresponds to the clone IMAGE: 3628076 isolated from brain-derived cell lines. The P2x7v02 lacking the exon 5 produces a frame shift, thus generating a shorter polypeptide. The second splicing variant, P2X7v03, is characterized by the presence of the short intron 10 into the mRNA. This variant is supported by two high quality sequences, the cDNA clone BRAMY2008977 (AC number: AK090866) from human amygdala and the EST clone dbEST:7339877 derived from an unknown human tumor. The last variant, P2X7v04, is defined by the lack of the first exon that suggests an alternative promoter usage closed to the exon 2. A high quality EST clone dbEST:4782844 derived from a head and neck tumor supports this variant. These variants are shown in Figures 16a to 16e.

## P2X7 variants.

```

P2X7v01      MPACCS SCDVFQYETNKVTRIQSMNYGTIKWFFHVIIIFSVCFALVSDKLYQRKEPVISS
P2X7v04      MPPVD-----AFPCLPFS-----FALVSDKLYQRKEPVISS
P2X7v02      MPACCS SCDVFQYETNKVTRIQSMNYGTIKWFFHVIIIFSVCFALVSDKLYQRKEPVISS
P2X7v03      MPACCS SCDVFQYETNKVTRIQSMNYGTIKWFFHVIIIFSVCFALVSDKLYQRKEPVISS
1.....10.....20.....30.....40.....50

P2X7v01      VHTKVKGIAEVKKEEIVENGVKLVHSVFDTADYTFPLQNSFFVMTNFLKTEGQEORLCP
P2X7v04      VHTKVKGIAEVKKEEIVENGVKLVHSVFDTADYTFPLQNSFFVMTNFLKTEGQEORLCP
P2X7v02      VHTKVKGIAEVKKEEIVENGVKLVHSVFDTADYTFPLQNSFFVMTNFLKTEGQEORLCP
P2X7v03      VHTKVKGIAEVKKEEIVENGVKLVHSVFDTADYTFPLQNSFFVMTNFLKTEGQEORLCP
61.....70.....80.....90.....100.....110

P2X7v01      EYPTRRRLC SSDRGCKKGWMDPQSKGIQTGRVCVHEGNQKTCEVSAWCPIEAVEEAPRPA
P2X7v04      EYPTRRRLC SSDRGCKKGWMDPQSKGIQTGRVCVHEGNQKTCEVSAWCPIEAVEEAPRPA
P2X7v02      EYPTRRRLC SSDRGCKKGWMDPQSKGLLS-----
P2X7v03      EYPTRRRLC SSDRGCKKGWMDPQSKGIQTGRVCVHEGNQKTCEVSAWCPIEAVEEAPRPA
121.....130.....140.....150.....160.....170

P2X7v01      LLNSAENFTVLIKNNIDFPGHNYTTRNLPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
P2X7v04      LLNSAENFTVLIKNNIDFPGHNYTTRNLPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
P2X7v02      -----
P2X7v03      LLNSAENFTVLIKNNIDFPGHNYTTRNLPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
181.....190.....200.....210.....220.....230

P2X7v01      NFSDVAIQGGIMGIEIYWCNLD RWFHCHPKYSFRRLDDKTTNVS LYPGYNFRYAKYYK
P2X7v04      NFSDVAIQGGIMGIEIYWCNLD RWFHCHPKYSFRRLDDKTTNVS LYPGYNFRYAKYYK
P2X7v02      -----
P2X7v03      NFSDVAIQGGIMGIEIYWCNLD RWFHCHPKYSFRRLDDKTTNVS LYPGYNFRYAKYYK
241.....250.....260.....270.....280.....290

P2X7v01      ENNVEKRTL I KVFGRFDILVFGTGGKFDIIQLVVYIGSTLSYFGLAAVFIDFLIDTYSS
P2X7v04      ENNVEKRTL I KVFGRFDILVFGTGGKFDIIQLVVYIGSTLSYFGLAAVFIDFLIDTYSS
P2X7v02      -----
P2X7v03      ENNVEKRTL I KVFGRFDILVFGTGGKFDIIQLVVYIGSTLSYFGLV RDSLFHALGKWFG
301.....310.....320.....330.....340.....350

P2X7v01      NCCRSHIYPWCKCCQPCVVNEYYYRKKCESIVEPKPTLKYVSFVDESHIRMVNQQLGRS
P2X7v04      NCCRSHIYPWCKCCQPCVVNEYYYRKKCESIVEPKPTLKYVSFVDESHIRMVNQQLGRS
P2X7v02      -----
P2X7v03      EGSD-----
361.....370.....380.....390.....400.....410

P2X7v01      LQDVKGQEVPRPAMDFDLSRLPLALHDT PPIPGQPEEIQLLRKEATPRSRDSPVWCQCG
P2X7v04      LQDVKGQEVPRPAMDFDLSRLPLALHDT PPIPGQPEEIQLLRKEATPRSRDSPVWCQCG
P2X7v02      -----
P2X7v03      -----
421.....430.....440.....450.....460.....470

P2X7v01      SCLPSQLPESHRCLEELCCRKKPGACITTS ELFRLVLSRHVLQFLLLYQEPLLDVDS
P2X7v04      SCLPSQLPESHRCLEELCCRKKPGACITTS ELFRLVLSRHVLQFLLLYQEPLLDVDS
P2X7v02      -----
P2X7v03      -----
481.....490.....500.....510.....520.....530

P2X7v01      TNSRLRHCA YRCYATWRFGSQDMADFA ILPSCCRWRIRKEFPKSEGQYSGFKSPY
P2X7v04      TNSRLRHCA YRCYATWRFGSQDMADFA ILPSCCRWRIRKEFPKSEGQYSGFKSPY
P2X7v02      -----
P2X7v03      -----
541.....550.....560.....570.....580.....590

```

Therefore the transcriptional and translational start sequences of the human P2X7R were analyzed using Blast, Genescan and HMMgene computer software. This analysis indicated that P2X7R possesses with high probability only one translation start site. Most P2X7R expression sequence tags (ESTs; Unique cluster Hs. 193470)

having a reliable 5' end showed identical transcriptional start site. None of the ESTs showed any indication of alternative splicing. Therefore, in silico analysis suggests that there is a low probability to find different transcripts produced by alternative splicing or alternative promoter usage.

The above mentioned in silico data were confirmed by RT-PCR analysis spanning the whole predicted human P2X7R coding sequence using 14 and 19 bases (5'-ATGCCGGCTTGCTG-3'; 5'-GTAGGGATACTTGAAGCCA-3') oligonucleotides corresponding to the beginning and end of the coding sequence, respectively. Total RNA from whole brain, different dissected brain areas, thymus, spleen and kidney were isolated and analyzed for P2X7R expression. RT-PCR reactions were performed using the C. Therm One Step polymerase system (Roche Applied Science) and a protocol for touch down PCR with hot start. Briefly, Reverse Transcription was performed at 52°C according to the manufacturer's conditions. PCR reactions were executed with an annealing temperatures of 64°C for the first five cycles and of 54°C for the next 30 cycles.

A single specific band of the size of 1785 bp corresponding to the complete coding sequence of P2X7R was detected. P2X7R mRNA was detected in the whole brain, hippocampus, cerebellum, leukocytes and thymus but not in cerebral cortex, hypothalamus, spleen and kidney (Figure 2). All PCR products were cloned using the pGEM-T-Easy plasmid (Promega), selected in Top-10 bacteria (Invitrogen) by blue-white selection and tested by EcoRI digestion. Clones having fragments of the expected size were amplified and purified for sequencing. The sequence confirmed the identity of the 1785 bp clones as the complete coding sequence of wild-type P2X7R. Therefore, in all the tissues tested, wild-type P2X7R is expressed as a single transcript which includes the complete coding sequence. The presence of tissue specific isoforms is unlikely. These studies provide useful information about the P2X7R mRNA expression and transcript processing. This information can be used to synthesize riboprobes for in situ hybridization, Northern and Southern blot as well as engineering cells for the overexpression of P2X7R.



## EXAMPLE 5

### P2X7R expression in the mouse brain

The expression of P2X7R was further studied by immunohistochemistry of serial sections of complete mouse brains using a polyclonal antibody directed against an internal peptide of P2X7R (Santa Cruz Biotechnology). The brains from stress-free mice were shock frozen, cut into 16  $\mu\text{m}$  slices and fixed with paraformaldehyde for 5 minutes. The sections were blocked for 30 minutes at room temperature with 1:10 horse serum. All antibodies were diluted in TBST buffer (Tris-buffered saline with 0.05% Tween-20). The first antibody was used in a dilution 1:200 and incubated overnight. All washes were performed with TBST buffer. As a secondary antibody, an anti-goat IgG biotinylated (Vector Laboratories) was used and detection was performed using the streptavidin-biotin-horse-radish peroxidase complex system (Vector Laboratories) in combination with diaminobencidine. Slides were counterstained with toluidine blue using standard procedures. The same procedure in the absence of the primary antibody was performed as a negative control. As a positive control to test the Preservation of the tissue was verified with an antibody specific for the protein Patched1 (Santa Cruz Biotechnology). Patched1 was used as positive control since it stains all relevant brain structures and is not affected by stress or antidepressants. Very specific staining pattern was detected, consistent with the specific subcellular localization of P2X7R in brain cells. Negative controls were completely devoid of signal. Positive control with Patched1 showed identical signal intensity and distribution in all samples, indicating that all tissues were equally well preserved and processed.

Proceeding from frontal to caudal, P2X7R protein was observed in the glomerular layer of the olfactory bulb at low levels (Figure 3). P2X7R was also present at very low levels in a restricted area of the periventricular hypothalamic nucleus (figure 3). Ependymal cells surrounding the lateral ventricles also showed a faint staining (Figure 3). A stronger signal was detected in restricted areas of the hypothalamus, where the signal was present in single cells of the polymorph layer, the lacunosum moleculare and the oriens layer (Figure 4). In more posterior areas of the hippocampus, the signal was present in the molecular layer, stratum radiatum and

near the CA3. In a further caudal position, P2X7R was expressed in the subcomisural organ (Figure 4). Therefore, the basal P2X7R expression in the brain of stress-free mice is restricted to areas that had been previously associated with depression, stress, learning and memory.

### EXAMPLE 6

#### P2X7R is modulated in mice treated with an antidepressant

Further validation of role of P2X7R in affective disorders was performed by examining its expression pattern in response to stress and treatment with antidepressant drugs. A treatment schedule which has been proven to produce antidepressant effects on the behavioural level was administered to mice which were characterized as antidepressant-responsive by using a variety of behavioural paradigms suitable to detect anxiolytic and antidepressant effects of classical antidepressants like the selective serotonin reuptake inhibitor paroxetine. Paroxetine was delivered by gavage to naive male mice over a time period of 28 days at a dosage of 10 mg/kg bodyweight twice per day. In parallel, a control group of mice was given vehicle solution (i.e. without paroxetine) using the same treatment regiment while a second control group of mice was left undisturbed and stress-free (i.e. untreated) during the same period of the experiments. At the end of the long-term treatment, part of the mice of each experimental group were tested in the dark/light box (test of anxiety behaviour) and in the Porsolt's forced swim test (test of depressive-like behaviour) to confirm the effectiveness of the treatment (Figure 5). Passive stress coping behaviour decreased after long-term treatment with the antidepressant paroxetine. The other part of the experimental groups (i.e. mice without test experience) were decapitated, brains rapidly removed and frozen at  $-80^{\circ}\text{C}$  until usage.

The expression of P2X7R in the brains of mice under stress-free conditions, and mice under mild stress produced by the vehicle application, and mice under paroxetine treatment was evaluated using three different brains from each group. Serial slides from each group of animals were analyzed in parallel by immunohistochemistry using the same materials in order to produce completely

comparable results. No significant change in P2X7R expression in the olfactory bulb was seen in response to stress or to paroxetine treatment (Figure 6). However, in the periventricular nucleus of the hypothalamus, paroxetine produced a slight inhibition of P2X7R expression (Figure 7). No significant change was observed in the ependymal cells from different brain areas (Figure 8). The most dramatic changes were observed in the hippocampus, where P2X7R was strongly inhibited by stressful handling whereas paroxetine treatment produced a marked stimulation above basal levels (Figures 9, 10 and 11). This effect was observed all along the hippocampus but was more evident in the polymorph layer near the dentate gyrus. In the subcommissural organ, P2X7R expression remained unchanged by the different treatments. Therefore, P2X7R expression is strongly regulated in two specific brain areas involved in depression and stress. Other brain areas, which showed low levels of P2X7R and are not directly involved in depression, did not show changes.

In the samples from mice treated with paroxetine and showing a strong P2X7R expression, it was possible to analyze the distribution of P2X7R in more detail (Figures 10 and 11). The P2X7R protein was not only present in cell bodies but also was clearly detected in projections innervating the granular layer of the dentate gyrus (Figure 12). This subcellular localization of P2X7R is consistent with a role in neurotransmitter release and long term potentiation.

Since some reports (Muria et al., *Biochem. J.* 288 (1992) ,897-901; Ferrari et al., *FEBS Lett.* 447 (1999), 71-75) suggest that chronic and high dose stimulation of P2X7R may cause apoptosis in some cell types, the hippocampus of the above described animals were analyzed for the co-localization of apoptotic cells and P2X7R expressing cells, in consecutive sections, using TUNNEL staining and immunohistochemistry. In correlative sections, only few apoptotic cells were detected and they were present along the granular layers of the hippocampus where no P2X7R expression was observed (Figure 13). No significant differences in the numbers of apoptotic cells were observed between the different treatment conditions. Therefore, the location and number of apoptotic cells did not correlate with the location and number of cells expressing P2X7R and rules out an involvement of P2X7R in the induction of apoptosis in the hippocampus.

Thus, P2X7R expression is considerably restricted to specific brain areas involved in depression. Moreover, P2X7R expression is inhibited by stress and strongly stimulated by antidepressant treatment in these specific areas. Therefore, P2X7R fulfils all criteria required for the actions of antidepressants according to the highest standards in the field of depression research. In addition, these results suggest that modulation of function of P2X7R is associated with chronic stress, which serves as a model for several aspects of affective disorders.

### EXAMPLE 7

#### The behavioural effect of P2X7R inhibition in mice

To demonstrate that P2X7R inhibition acts as a causative agent for affective disorders, P2X7R function was specifically inhibited in distinct regions of the brain without affecting any other brain function. This was achieved by delivering double stranded small interference RNA molecules (siRNA) into restricted areas of the brain.

According to the observed expression pattern of P2X7R in the hippocampus (Figures 9, 10, and 11) and the known involvement of the hippocampus in depression, the dentate gyrus (hippocampus) was selected as target region for siRNA application. Male, naive mice were bilaterally implanted with a guide cannulae (23 gauge, length 8 mm) by means of a stereotactic instrument. The coordinates, in relation to bregma, were -2.0 mm posterior,  $\pm 1.0$  mm lateral, and -1.0 mm ventral. Following a recovery period of 5 days, the mice were divided into three experimental groups: vehicle (veh), control double stranded RNA (control), and P2X7R specific double stranded siRNA (siRNA). Sequences used for P2X7R siRNA are 5'-GUGGGUCUUGCACAUGAUCTT-3' and 5'-GAUCAUGUGCAAGACCCACTT-3'. Both sequences were annealed and injected together as a double stranded RNA. On day 6 after surgery, mice were slightly anaesthetized with Isofluran and injections of siRNA were carried out. The concentration of the control and siRNA was 0.1 nmol/ $\mu$ l, and a volume of 1  $\mu$ l per side was infused using specifically adapted injection systems (30 gauge, length 9 mm). The anaesthesia for the infusion was of

short duration and the mice were awake immediately or few seconds after the manipulation.

Once delivered into the brain the siRNA molecules specific for P2X7R were taken up by brain cells and specifically induce the degradation of the complementary P2X7R mRNA with high efficiency. As a result, P2X7R function was specifically inhibited for a short period without affecting any other brain function. In this regard, injection of vehicle or control siRNA did not result in any obvious changes in normal behaviour, i.e., food and water intake, or motor behaviour in the home cage.

The effects of P2X7R inhibition on depressive-like behaviour was assessed 24 hours and 48 hours after infusion of siRNA, control or vehicle according to the standard test paradigm, the Porsolt's forced swim test (Porsolt et al., Arch. Int. Pharmacodyn. 229 (1977), 327-336; Porsolt, Rev. Neurosci. 11 (2000), 53-58). The parameter used to evaluate depressive-like behaviour is the time the animal is floating in the water, a behaviour which is associated with behavioural despair as the animal does not make any effort to actively cope with the stressful situation. Compared to vehicle application, no influence of control double stranded RNA (5'-CAACUUCAUCUUCUACGCGTT-3') on floating behaviour (passive stress coping) was detected. In contrast, compared to controls, mice infused with P2X7R specific siRNA showed a significant increase in passive behaviour, which is construed as depressive-like behaviour. This interpretation becomes moreover evident when the effects of antidepressants on passive stress coping behaviour in the forced swim test are visualized (Figure 14). Passive stress coping behaviour increased after acute intrahippocampal (bilateral, dentate gyrus) of siRNA targeting P2X7R. The Porsolt's forced swim test is a standard test used to assess the effectiveness of antidepressants and it has been proven by many studies that the test is selectively sensitive for these effects, given that the right animal model is used. The paradigm has been widely used to test pharmaceutical compounds and to validate animal models of depression, which show an increase in passive behaviour as do the mice where P2X7R has been inhibited (siRNA).

At the end of the experiment, the mice were sacrificed and the brains were examined to confirm the location and efficiency of the siRNA injections. For this purpose the brains were cut into sections and the slides were stained by immunohistochemistry using the above mentioned protocols. Brains from mice injected with the specific double stranded siRNA, with control double stranded RNA and with vehicle were examined in parallel. Under these conditions, the specific siRNA directed against P2X7R injected near the dentate gyrus induced on average an 80% inhibition of P2X7R protein expression as compared to the samples from mice injected with vehicle or with control double stranded RNA. Both the number of cells expressing P2X7R as well as the intensity of the expression were strongly reduced (Figure 15). The injections with siRNA did not produce any sign of local inflammation or infiltration at the hippocampus. Thus, P2X7R expression is specifically and locally inhibited by siRNA application in vivo. This inhibition produced behavioural changes indicating a causative role for P2X7R in affective disorders. These results in combination with those mentioned above support and confirm the observation of mutations in P2X7R being associated with affective diseases in humans and that modulation of P2X7R activity has antidepressive effects.

### EXAMPLE 8

#### Drug screening assay

Methods for identifying P2X7R agonists were established using an immortalised mouse hippocampal cell line expressing the endogenous P2X7 gene. Briefly, the expression of P2X7 was confirmed by culturing the cells at 37°C/5% CO<sub>2</sub> in DMEM with 10% foetal calf serum (Gibco). Upon reaching 80 % confluence, cells were collected in PBS and homogenized by repeated passage through a syringe (18G needle). The amount of total protein was measured by the Bradford assay (Sigma; diluted 1:5, O.D. measured at 595 nm) according to the manufacturer's recommendation. Protein homogenates were then mixed with an equal volume of loading buffer (50 mM Tris-Cl pH 6.8; 25% glycerol; 7,2 mM bromophenol blue; 2% SDS; 200 nM β-mercaptoethanol) and subsequently denaturated in boiling water for 10 minutes. 20 mg of each sample were loaded onto a 10% polyacrylamide gel

containing 0,4 % SDS. Electrophoresis and Western blot transfer were performed according to conventional protocols described, for example, in Sambrook, Russell "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Laboratory, N.Y. (2001). Membranes were then blocked with 5 % dry milk and incubated with an antibody against P2X7R (1:1000 dilution; Santa Cruz Biotech) followed by incubation with a horse anti-goat peroxidase-coupled secondary antibody (1: 10000 dilution; Santa Cruz Biotech). Membranes were then incubated for 1 hour at 37°C in Lumi-Light Western Blotting Substrate (Roche Applied Science) followed by a 10 minute exposure on a BioMax MR Film (Kodak).

A 70 kD band corresponding to the expected size of the P2X7R protein was detected in HT-22 cells demonstrating expression of the endogenous mouse P2X7 gene (Figure 17). A second mouse hippocampal cell line (HT-39) did not express P2X7.

Since P2X7R is an ATP-gated ion channel which allows the entry of calcium and sodium ions into cells, a method for identifying P2X7R agonist was established by monitoring calcium influx into HT-22 cells. Cells were first loaded with the fluorescent dye Oregon green AM ester (Molecular Probes) for 30 minutes at room temperature, washed 2 times with DMEM/10% foetal calf serum to remove excess dye and cultured for 15 minutes in the presence of 100  $\mu$ M 2- and 2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate (BzATP:  $C_{24}H_{24}N_5O_{15}P_3$ ). BzATP is a known agonist of P2X7R (North and Surprenant, Annu. Rev. Pharmacol. Toxicol. 40 (2000), 563-580). Calcium movement into the cells was visualised under a fluorescent microscope with a fluorescein filter (wavelength 492/517 nm). Oregon green AM ester is a fluorescent dye that binds to intracellular calcium. Accordingly, an increase in green fluorescence was observed in cells treated with BzATP (Figure 18) signalling an activation of P2X7R which results in an influx of calcium into the cells and an increase binding of Oregon green dye to intracellular calcium. Alternatively, Oregon green AM ester can be replaced by Fluo-3, fluo-4, fluo-5F, fluo-5N, fluo-4FF, Fluo-4 dextran, Fluo-3 AM, Fluo-4 AM, Fluo-5F AM, Fluo-5N AM and Fluo-4FF AM (Molecular Probes). Calcium influx in HT-22 cells can also be measured in 96-well and 384-well microplate using the Calcium Plus Assay Kit (Molecular Device) or FLIPR ® Calcium Assay Kit for Fluorometric Imaging Plate

Reader Systems (Molecular Device). HT-22 cells can be replaced by any cells expressing P2X7R, including cells that have been genetically modified by introducing an exogenous P2X7 gene.

The specificity of the P2X7R agonist on calcium influx was confirmed by pre-treatment of HT-22 cells with 100 mM Oxidized ATP (oATP; Sigma) for 1 hour before the addition of BzATP. oATP is an irreversible inhibitor of the receptor (Chen et al., J. Biol. Chem., 268 (1993), 8199-8203). Activation of P2X7R by the agonist was inhibited by oATP (Figure 18) as illustrated by the absence of green fluorescence in the cells.

Yet another method of measuring P2X7R activity involves the entry of ethidium bromide into P2X7R expressing cells. Activation of P2X7R by an agonist allows the entry of ethidium bromide which binds nuclear DNA and emits a fluorescence signal. Alternatively, the propidium dye YOPRO-1 can be substituted for ethidium bromide. An increase in fluorescence can be used as a measure of P2X7 receptor activation. Therefore, the assay can be used to test and quantify the effect of an agent or compound with agonist properties on P2X7R. In the present example,  $10^3$  HT-22 cells were seeded per well in a 96-well flat bottom microtitre plates and incubated at 37°C/5%CO<sub>2</sub> in DMEM medium containing 10% FCS until the cells attached to the culture surface. Once attached, cells were incubated for 60 minutes in DMEM medium containing 10%FCS,  $10^{-4}$ M ethidium bromide and increasing concentrations of BzATP (1  $\mu$ M, 10  $\mu$ M, 100  $\mu$ M, 500 $\mu$ M, 1 mM). The number of fluorescent cells which have integrated the ethidium bromide to the DNA can then be counted using a fluorescent microscope (Zeiss, Germany). Concentrations above 100 $\mu$ M BzATP increased the number of fluorescent nuclei signalling activation of P2X7R (Figure 19a). Alternatively, ethidium bromide fluorescence can be measured using a Perkin-Elmer fluorescent plate reader (excitation 520 nm, emission 595 nm, slit widths: Ex 15 nm, Em 20 nm). From the readings obtained, a pIC50 figure can be calculated for each candidate agent or compound. Accordingly, a P2X7R agonist is defined as an agent or a compound with an EC50 equal or below 300 micromolar, whereas the term EC 50 is defined as the concentration eliciting 50% of maximal response to an agonist (North and Surprenant, Annu. Rev. Pharmacol. Toxicol. 40 (2000), 563-580).



The specificity of an agonist for P2X7R can be evaluated by pre-incubation of the cells for 60 minutes with 100  $\mu$ M o-ATP before adding the agonist and ethidium bromide dye. Under these conditions, activation of P2X7R by the agonist is inhibited by oATP resulting in a reduction in the number of fluorescent cells (Figure 19b).

## EXAMPLE 9

### Activation of P2X7R with agonists has antidepressive effects

To demonstrate that activation of P2X7R has therapeutic effects on affective disorders, the P2X7R agonist BzATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate ( $C_{24}H_{24}N_5O_{15}P_3$ )) was administered to a selected DBA/2O1a mouse strain that displays characteristics of being highly anxious, responding to antidepressants, and showing anxiolysis after subchronic antidepressant treatment (Lucki et al., *Psychopharmacology* 155 (2001), 315-322). BzATP is a compound with strong specificity to P2X7R (North and Surprenant, *Annu. Rev. Pharmacol. Toxicol.* 40 (2000), 563-580). In the present example, the P2X7R agonist was directly injected into the hippocampus of mice. However, a P2X7R agonist agent or compound could also be delivered orally, subcutaneously, intravenously, intra-arterial, intranodal, intramedullary, intrathecal, intraventricular, intranasally, intrabronchial, transdermally, intrarectally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly.

Four months old male mice were bilaterally implanted with guide cannulae (23 gauge, length 8 mm) by means of a stereotactic instrument (David Kopf Instruments). The coordinates, in relation to bregma, were -2.0 mm posterior,  $\pm 1.0$  mm lateral, and -1.0 mm ventral. After surgery, the mice were allowed to recover for 10 to 12 days. Following this recovery period, mice were injected with 1  $\mu$ l vehicle solution (0.5% DMSO, Sigma) or 50  $\mu$ M BzATP (Sigma, prepared in 0.5% DMSO) in each side of the brain over a period of 60 seconds. Injections were performed using a 9 mm-30 gauge needle inserted into the guide cannulae and connected via tubing to a 10  $\mu$ l Hamilton syringe.

The behaviour of individual mice was assessed using the Porsolt's forced swim test 24 hours after injection of vehicle solution or BzATP. A pre-exposure of 5 minutes to the test was done 10-15 minutes after vehicle or BzATP injection. The forced swim test is a standard test that measures primary stress-induced reductions in avoidance or escape, termed behavioural despair. The test is used to determine the effectiveness of antidepressants, test new pharmaceutical compounds and validate animal models of depression (Porsolt et al., Arch. Int. Pharmacodyn. 229 (1977), 327-336; Porsolt, Rev. Neurosci. 11 (2000), 53-58; Rénérice et al., Behav. Brain Res. 136 (2002), 521-532; Page et al., Psychopharmacology 165 (2003), 194-201; Kelliher et al., Psychoneuroendocrinology 28 (2003), 332-347). The test consists of placing a mouse for a period of 5 minutes into a glass cylinder containing water. Under such circumstances, the mouse cannot touch the bottom of the cylinder and is thus forced to swim. Time, latency and frequency of struggling versus floating are scored as behavioural parameters. Floating (i.e. movements made only for keeping balance and breath) is a passive behaviour associated with despair and represents a depressive-like symptom since the animal does not make any effort to actively cope with the stressful situation. Increased struggling (i.e. active attempts to escape) indicates active coping behaviour that can be interpreted as an improvement of depression-like symptoms. For example, treatment with serotonergic antidepressants reduce the total time spent floating (Borsini, Neurosci. Biobehav. Rev. 19 (1995), 377-395; Redrobe and Bourin, Psychopharmacology 138 (1998), 198-206).

The P2X7R agonist BzATP was found to increase active escape attempts (i.e. increase in time and frequency of struggling, decrease in latency of struggling) while a decrease in passive behaviour (i.e. decrease in time and frequency of floating, increase in latency of floating) was measured when compared to control mice injected with vehicle solution (Figure 20). Observed results were verified statistically using Mann-Whitney U and one-way MANOVA tests. The differences in time struggling, latency of floating and frequency of floating were found to be statistically significant. While latency and frequency of struggling and time floating results were not supported statistically, they still represented a tendency towards improvement in stress coping behaviour. These results demonstrate that a P2X7R agonist can lead to improvements in depressive-like symptoms.

Since conclusions drawn from the forced swim test can be influenced by unspecific effects of an agent or compound on animal activity (i.e. increase in struggling behaviour can be the result of hyperactivity instead of increased active coping behaviour), the potential effect of BzATP on locomotor activity was assessed by the open field test (Crawley "What's wrong with my mouse: Behavioral phenotyping of transgenic and knockout mice", Wiley-Liss (2000)). Locomotor activity in mice treated with control vehicle solution or 50  $\mu$ M BzATP was assessed 24 hours after injection by placing individual animal in a dark-grey wooden box (30x30x40 cm). Locomotor activity was monitored for a period of 30 minutes using a video camera. Overall distance travelled by the animals during the testing period was then analysed by means of VideoMot2 computer software (TSE GmbH, Bad Homburg). No difference in locomotor activity was measured between mice treated with control vehicle solution and BzATP (Figure 21). Therefore, the application of BzATP did not induce hyperactivity. These results confirm that activation of P2X7R by an agonist agent or compound leads to improvements in depressive-like symptoms and is not the result of an unspecific effect on animal activity per se.

Several reports suggest that activation of P2X7R can induce apoptosis and cell death in vitro (Di Virgilio et al., Cell Death Differ. 5 (1998), 191-199, Virginio et al., J. Physiol. 519 (1999), 335-346). To test whether P2X7R activation in the hippocampus resulted in cell death, apoptosis levels were quantified in the brain of the mice treated with BzATP. Mice were sacrificed at the end of the behavioural experiments, the brains were removed, shock frozen and sectioned into 16  $\mu$ m slices. Brain sections were then studied for apoptosis using the DeadEnd fluorometric TUNEL system according to the manufacturer's recommendation (Promega Corporation). The TUNEL system measures the fragmented DNA of apoptotic cells. Positive control for the assay are made by pre-treating brain sections for 10 minutes with 1 unit/ml of DNase I.

Very few apoptotic cells (i.e. less than one cell per brain section) were observed in brains of mice treated with control vehicle or the P2X7R agonist (Figure 22) when compared to positive control sections pre-treated with DNase. Moreover, no

significant differences in the numbers of apoptotic cells was observed between the control animals and mice treated with BzATP, indicating that activation of P2X7R did not result in cerebral cell death in vivo.

### Claims

1. A nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:
  - (a) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 5'UTR region corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;
  - (b) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of the ATP-gated ion channel P2X7R, wherein in the exon as indicated in column "Exon" of the following Table A the amino acid residue as indicated in column "Amino acid residue" of Table A corresponding to the position as indicated in column "Position in wild-type" of Table A of the wild-type ATP-gated ion channel P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue

**Table A**

<b>Exon</b>	<b>Amino acid residue</b>	<b>Position in wild-type</b>
exon 3	R (Arg)	117
exon 5	G (Gly)	150
exon 6	E (Glu)	186
exon 6	L (Leu)	191
exon 8	R (Arg)	270
exon 13	I (Ile)	568
exon 13	R (Arg)	578

- (c) a nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in exon 5 or 8 corresponding to position 32548 or position 37633 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1,

wherein at said position said nucleotide is replaced by another nucleotide

- (d) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted;
- (e) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R, wherein in the intron as indicated in column "Intron" of the following Table B the nucleotide as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1 is replaced by another nucleotide

**Table B**

<b>Intron</b>	<b>Replaced nucleotide</b>	<b>Position in wild-type</b>
intron 1	G	3166
intron 1	C	24778
intron 1	C	24830
intron 3	A	26308
intron 3	G	26422
intron 4	G	32394
intron 4	T	32434
intron 5	A	32783
intron 6	G	35641
intron 6	A	35725
intron 6	T	36001
intron 7	G	36378
intron 7	T	36387
intron 7	G	36398
intron 9	C	47214
intron 11	T	47563
intron 12	C	54307
intron 12	G	54308

- (f) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 3'UTR region corresponding to position 54925, 55169, 55170, 55171 or 55917 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as

depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;

- (g) a nucleotide sequence comprising at least 20 or 21 nucleotides and comprising the mutations or deletions as defined in any one of (a) to (f);
  - (h) a nucleic acid sequence comprising a nucleotide sequence as shown in any one of SEQ ID NOs: 13 to 51;
  - (i) a nucleic acid sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NOs: 5 to 12;
  - (j) a nucleotide sequence which hybridizes to a nucleotide sequence defined in any one of (a) to (g) or to the nucleotide sequence of (h) and having a mutation as defined in any one of (a) to (f); and
  - (k) a nucleic acid sequence being degenerate as a result of the genetic code to the nucleic acid sequence as defined in (j).
2. The nucleic acid molecule of claim 1 derived from mouse, rat or human.
  3. The nucleic acid molecule of claim 1 or 2 which is DNA, RNA, PNA or phosphorothioates.
  4. A vector comprising the nucleic acid molecule of any one of claims 1 to 3.
  5. The vector of claim 4 which is an expression vector, a gene targeting vector and/or a gene transfer vector.
  6. A host transformed with a vector of claim 4 or 5 or transformed with the nucleic acid molecule of any one of claims 1 to 3.
  7. The host of claim 6 which is a mammalian cell, an amphibian cell, a fish, an insect cell, a fungal cell, a plant cell or a bacterial cell.
  8. The host of claim 7, wherein said mammalian cell is selected from the group consisting of CHO cells, HEK293 cells, COS-7 cells or PC12 cells.

9. The host of claim 7, wherein said amphibian cell is an oocyte, preferably a *Xenopus* oocyte.
10. The host of claim 9, wherein said oocyte is a frog oocyte.
11. The host of claim 6 which is a non-human transgenic organism.
12. The host of claim 11, wherein said non-human organism is a mammal, amphibian, a fish, an insect, a fungus or a plant.
13. A method for producing the polypeptide encoded by a nucleic acid molecule of claims 1(b) or 1(d) comprising culturing/raising the host of any one of claims 6 to 11 and isolating the produced polypeptide.
14. A polypeptide encoded by the nucleic acid molecule of claim 1(b) or 1(d) or produced by the method of claim 13.
15. An antibody specifically directed to the polypeptide of claim 14, wherein said antibody specifically reacts with an epitope generated and/or formed by the mutation in the ATP-gated ion channel P2X7R selected from the group consisting of:
  - (i) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein the R (Arg), G (Gly), E (Glu), L (Leu), R (Arg), I (Ile) or R (Arg) residue corresponding to position 117, 150, 186, 191, 270, 568 or 578 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue; and
  - (ii) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted.



16. The antibody of claim 15 which is a monoclonal antibody.
17. An aptamer specifically binding to a nucleic acid molecule of any one of claims 1 to 3 or to the polypeptide of claim 14.
18. A primer or pair of primers capable of specifically amplifying a nucleic acid molecule as defined in any one of claims 1 to 3.
19. The primer of pair of primers of claim 18, which is selected from the group consisting of SEQ ID NOs.: 52 to 111.
20. A composition comprising the nucleic acid molecule of any one of claims 1 to 3, the vector of claim 4 or 5, the polypeptide of claim 14, the antibody of claim 15 or 16, the aptamer of claim 17 and/or the primer or pair of primers of claim 18 or 19.
21. The composition of claim 20 which is a diagnostic composition.
22. The diagnostic composition of claim 21, optionally further comprising suitable means for detection.
23. Use of the nucleic acid molecule of any one of claims 1 to 3, the vector of claim 4 or 5, the polypeptide of claim 14, the antibody of claim 15 or 16, the aptamer of claim 17 and/or the primer or pair of primers of claim 18 or 19 for the preparation of a diagnostic composition for the detection of an affective disorder.
24. A method of diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2XR7 protein expressed in the cells of said individual is non-functional, shows an altered ATP-gating in comparison to

the wild-type P2XR7 protein or is over- or under-expressed in comparison to the P2XR7 protein level an unaffected individual.

25. A method for diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R gene sequence or encoded protein thereof comprises a mutation in comparison to the wild-type P2X7R sequence.
26. The method of claim 25, wherein said mutation is a mutation as defined in claim 1 and/or a nucleotide replacement or deletion selected from the following Table C indicating in column "Region of P2X7R" the region of the P2X7R genomic nucleotide sequence in which the replacement or deletion occurs, in column "Nucleotide" of Table C the nucleotide which is replaced by another nucleotide or the nucleotides which are deleted and in column "Position in wild-type" of Table C the corresponding position in the nucleotide sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1

**Table C**

<b>Region of P2X7R</b>	<b>Nucleotide</b>	<b>Position in wild-type</b>
5'UTR	T	362
5'UTR	T	532
5'UTR	A	1100
5'UTR	A	1122
5'UTR	C	1171
5'UTR	T	1351
5'UTR	G	1702
5'UTR	T	1731
5'UTR	C	1860
5'UTR	C	2162
5'UTR	C	2238
5'UTR	A	2373
5'UTR	G	2569
5'UTR	G	2702
Intron 1	G	3166
intron 1	C	24778
intron 1	C	24830
exon 2	T	24942

exon 3	C	26188
exon 3	A	26308
exon 3	G	26422
intron 4	G	32394
intron 4	T	32434
exon 5	G	32493
exon 5	G	32506
exon 5	C	32507
exon 5	C	32548
intron 5	A	32783
intron 5	T	35309
intron 5	C	35374
intron 5	A	35378
exon 6	G	35438
exon 6	T	35454
intron 6	T	35549
intron 6	G	35641
intron 6	A	35725
intron 6	T	36001
intron 6	A	36064
intron 6	deletion of GTTT	36091 to 36094
intron 6	C	36108
intron 7	C	36374
intron 7	G	36378
intron 7	T	36387
intron 7	G	36398
intron 7	C	37439
intron 7	T	37513
exon 8	C	37604
exon 8	G	37605
exon 8	G	37623
exon 8	C	37633
intron 9	C	47214
exon 11	G	47383
exon 11	C	47411
intron 11	T	47563
intron 12	C	54307
intron 12	G	54308
exon 13	C	54399
exon 13	A	54480
exon 13	C	54523
exon 13	deletion of CCCTGAGAGCCACAGGTGCCT	54562 to 54582
exon 13	A	54588
exon 13	C	54664
exon 13	G	54703
exon 13	A	54804

exon 13	G	54834
exon 13	G	54847
3'UTR	G	54925
3'UTR	C	55169
3'UTR	A	55170
3'UTR	A	55171
3'UTR	C	55917

27. The method of claim 26, wherein the occurrence of the mutation in the ATP-gated ion channel P2X7R gene is determined by PCR or immunological methods.
28. The composition of claim 20 which is a pharmaceutical composition.
29. The pharmaceutical composition of claim 28, optionally further comprising a pharmaceutically acceptable carrier.
30. A method of treating an affective disorder comprising administering a therapeutically effective amount of the nucleic acid molecule as defined in any one of claims 1 to 3 or a therapeutically effective amount of the polypeptide of claim 14 to a subject suffering from said disorder.
31. Use of the nucleic acid molecule of any one of claims 1 to 3, the vector of claim 4 or 5, the polypeptide of claim 14, the antibody of claim 15 or 16 and/or the aptamer of claim 17 for the preparation of a pharmaceutical composition for the treatment of an affective disorder.
32. A pharmaceutical composition comprising a nucleic acid molecule comprising a nucleotide sequence which encodes a functional ATP-gated ion channel P2X7R and which is selected from the group consisting of:
  - (a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as depicted in SEQ ID NO: 3 or 4;
  - (b) a nucleotide sequence comprising the nucleotide sequence as depicted in SEQ ID NO: 1 or 2;

- (c) a nucleotide sequence which hybridizes to the nucleotide sequence of (a) or (b); and
  - (d) a nucleotide sequence which is degenerated as a result of the genetic code to the nucleotide sequence of (c).
33. A pharmaceutical composition comprising a compound the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells or comprising a nucleic acid molecule the expression of which in cells or the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells.
34. The pharmaceutical composition of claim 33 which is an antisense nucleic acid, a ribozyme, a co-suppressive nucleic acid, iRNA or siRNA.
35. A method of treating an affective disorder comprising administering a therapeutically effective amount of the nucleic acid molecule as defined in any one of claims 32 to 34 or a therapeutically effective amount of the polypeptide encoded thereof to a subject suffering from said disorder.
36. Use of the nucleic acid molecule as defined in any one of claims 32 to 34 for the preparation of a pharmaceutical composition for treating an affective disorder.
37. Use of a modulator of P2X7R activity for the preparation of a pharmaceutical composition for treating an affective disorder.
38. The use of claim 37, wherein said modulator is an agonist.
39. The use of claim 38, wherein said agonist is selected from the group consisting of ATP, ATP-4 and BzATP (2'-3'-O-(4-Benzoylbenzoyl)adenosine 5'-triphosphate ( $C_{24}H_{24}N_5O_{15}P_3$ )).

40. The use of any one of claims 23, 30 or 36 to 39 or the method of any one of claims 24 to 27 or 35, wherein said affective disorder is selected from the group consisting of major depression, generalized anxiety disorder and bipolar disorder.
41. The use or the method of claim 40, wherein said major depression is selected from the group consisting of major depression, dysthymia, atypical depression, premenstrual dysphoric disorder and seasonal affective disorder.
42. The use or the method of claim 40, wherein said generalized anxiety disorder is selected from the group consisting of panic disorder, phobias, agoraphobia, social phobia, specific phobia, obsessive-compulsive disorder, post-traumatic stress disorder, separation anxiety disorder, mania, hypomania and cyclothymic disorder.
43. The use or the method of claim 40, wherein said bipolar disorder is bipolar disorder type I or bipolar disorder type II.
44. A kit comprising the nucleic acid molecule of any one of claims 1 to 3, a vector of claim 4 or 5, a host of any one of claims 6 to 11, a polypeptide of claim 14, an antibody of claim 15 or 16, an aptamer of claim 17 and/or a primer or pair of primers of claim 18 or 19.
45. A method for identifying compounds which are capable of specifically interacting with the polypeptide of claim 14, comprising the steps of
- (a) contacting a polypeptide of claim 14 with a compound or a candidate mixture of compounds to be tested; and
  - (b) determining whether said compound or a candidate mixture of compounds is capable of specifically interacting with said polypeptide.

46. A method for the characterization of compounds which are capable of altering characteristics of the polypeptide of claim 14, comprising the steps of
- (a) contacting a polypeptide of claim 14 with a compound or a candidate mixture; and
  - (b) determining whether the compound or a candidate mixture alters a characteristic of the polypeptide of claim 14.
47. A method of screening for compounds which are capable of interacting with the polypeptide of claim 14, comprising the steps of
- (a) contacting a polypeptide of claim 14 with a compound or a candidate mixture of compounds;
  - (b) measuring and/or detecting a response; and
  - (c) comparing said response to a standard response as measured in the absence of said candidate molecule.
48. A method for the production of a pharmaceutical composition comprising the steps of the method of any one of claims 45 to 47 and comprising a further step, wherein a derivative of said identified, characterized and/or screened molecule is generated.
49. A method for the production of a pharmaceutical composition comprising the steps of the method of any one of claims 45 to 48 and formulating the molecules identified, characterized, screened and/or derivatized in pharmaceutically acceptable form.
50. The method of claim 49, wherein the pharmaceutical composition to be produced further comprises neuroprotective substances, nootropic substances, brilliant blue, piperidine or piperazine derivatives thereof, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate or 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate.

51. The method of any one of claims 45 to 50, wherein said compound(s) or candidate mixture(s) of compounds comprise(s) antagonist(s), partial antagonist(s), partial agonist(s) and/or agonist(s) for an altered ATP-gated ion channel P2X7R.
52. A method for diagnosing an affective disorder of an individual comprising:
- (a) isolating DNA from cells obtained from an individual;
  - (b) determining all or part of the nucleotide composition of the P2X7R gene; and
  - (c) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism(s) mutation or allelic variation.
53. A method for diagnosing an affective disorder of an individual comprising:
- (a) isolating RNA from cells obtained from an individual;
  - (b) converting said RNA into cDNA;
  - (c) determining all or part of the nucleotide composition of the P2X7R gene; and
  - (d) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism(s), mutation or allelic variation.
54. A method for diagnosing an affective disorder of an individual comprising:
- (a) isolating RNA or protein from cells obtained from an individual ;
  - (b) determining the levels of P2X7R RNA or protein; and
  - (c) comparing the levels of P2X7R RNA or protein with the corresponding levels from a normal individual not afflicted with an affective disorder.



### **Abstract**

The present invention relates to nucleic acid molecules, preferably genomic sequences, encoding an ATP-gated ion channel P2X7R which contain a mutation in the 5'UTR or 3'UTR regions, a mutation in exon 3, 5, 6, 8 or 13 or in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 or a deletion in exon 13, which allow to diagnose affective disorders. The invention further relates to polypeptides encoded by said nucleic acid molecules vectors and host cells comprising said nucleic acid molecules as well as to methods for producing polypeptides encoded by said nucleic acid molecules. The present invention also provides antibodies specifically directed to polypeptides encoded by said nucleic acid molecules and aptamers specifically binding said nucleic acid molecules.

Additionally, primers for selectively amplifying said nucleic acid molecules are provided in the present invention as well as kits, compositions, particularly pharmaceutical and diagnostic compositions comprising said nucleic acid molecules, vectors, polypeptides, aptamers, antibodies and/or primers. Moreover, the present invention relates to methods for diagnosing affective disorders associated with a non-functional P2X7R protein, an altered ATP-gating of the P2X7R protein, an over- or underexpression of the P2X7R protein or associated with the presence of any one of the aforementioned nucleic acid molecules or polypeptides encoded thereby. Additionally, the present invention relates to uses and methods for treating affective disorders employing a functional or non-functional ATP-gated ion-channel P2X7R.

The present invention also relates to uses of modulators of P2X7R activity for treating affective diseases.

Furthermore, the present invention also relates to methods for identifying and characterizing compounds which are capable of specifically interacting with or altering the characteristics of the polypeptides of the present invention as well as to methods for the production of pharmaceutical compositions.

5,2Mb

1,6Mb

12/1




Figure 1a

2131




Figure 1b




Figure 1c

4131




Figure 1d

s/31




Figure 1e

6131





Figure 2

Ependymal cells of the  
lateral ventricle  
of a stress-free mouse:



hypothalamus  
of a stress-free mouse



periventricular  
hypothalamic nucleus

olfactory bulb  
of a stress-free mouse




glomerular layer

Figure 3




8(31

Subcommissural organ  
Of a stress-free mouse



dentate gyrus/hippocampus  
of a stress-free mouse



granular layer  
polymorph layer

Figure 4

9/31




Figure 5

10/31






Figure 6

11(31


paroxetine-treated  
mouse (stressful handling)



vehicle-treated mouse  
(stressful handling)



stress-free mouse



periventricular  
hypothalamic nucleus

Figure 7

12131

paroxetine-treated mouse  
(stressful handling)



vehicle-treated mouse  
(stressful handling)



stress-free mouse






Figure 8

13/31

paroxetine-treated mouse  
(stressful handling)



vehicle-treated mouse  
(stressful handling)



stress-free mouse



**Figure 9**

14131




Figure 10

15/31




Figure 11



16131




Figure 12

17131




Figure 13

18131





Figure 14

**Figure 15**

20/31

P2X7v01 MPACCSVDVFQYETNKVTRIQSMNYGTIKWFFHVIIFSYVCFALVSDKLYQRKEPVISS  
P2X7v04 MPPVD-----AFPCLPFS---FALVSDKLYQRKEPVISS  
P2X7v02 MPACCSVDVFQYETNKVTRIQSMNYGTIKWFFHVIIFSYVCFALVSDKLYQRKEPVISS  
P2X7v03 MPACCSVDVFQYETNKVTRIQSMNYGTIKWFFHVIIFSYVCFALVSDKLYQRKEPVISS  
1.....10.....20.....30.....40.....50  
P2X7v01 VHTKVKGIAEVKEEIVENGVKLVHSVFDTDADYTFPLQGNSEFFVMTNFKTEGQEORLCP  
P2X7v04 VHTKVKGIAEVKEEIVENGVKLVHSVFDTDADYTFPLQGNSEFFVMTNFKTEGQEORLCP  
P2X7v02 VHTKVKGIAEVKEEIVENGVKLVHSVFDTDADYTFPLQGNSEFFVMTNFKTEGQEORLCP  
P2X7v03 VHTKVKGIAEVKEEIVENGVKLVHSVFDTDADYTFPLQGNSEFFVMTNFKTEGQEORLCP  
61.....70.....80.....90.....100.....110

Figure 16a

```

P2X7v01 EYPTRRTLCSDDRGCKKGWMDPQSKGIQTGRGVVHEGNQKTCEVSAWCPIEAVEEAPRPA
P2X7v04 EYPTRRTLCSDDRGCKKGWMDPQSKGIQTGRGVVHEGNQKTCEVSAWCPIEAVEEAPRPA
P2X7v02 EYPTRRTLCSDDRGCKKGWMDPQSKGLLS-----
P2X7v03 EYPTRRTLCSDDRGCKKGWMDPQSKGIQTGRGVVHEGNQKTCEVSAWCPIEAVEEAPRPA
      121.....130.....140.....150.....160.....170

P2X7v01 LLNSAENFTVLIKNNIDFPGHNYTTRNILPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
P2X7v04 LLNSAENFTVLIKNNIDFPGHNYTTRNILPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
P2X7v02 -----
P2X7v03 LLNSAENFTVLIKNNIDFPGHNYTTRNILPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
      181.....190.....200.....210.....220.....230

```

Figure 16b

```

P2X7v01 NFSDVAIQGGIMGIEIYWDCNLD RWFHCHPKYSFRRLDDKT TNVSLYPGYNFRYAKYYK
P2X7v04 NFSDVAIQGGIMGIEIYWDCNLD RWFHCHPKYSFRRLDDKT TNVSLYPGYNFRYAKYYK
P2X7v02 -----
P2X7v03 NFSDVAIQGGIMGIEIYWDCNLD RWFHCHPKYSFRRLDDKT TNVSLYPGYNFRYAKYYK
      241.....250.....260.....270.....280.....290

P2X7v01 ENNVEKR TLIKVFGIRFDILVFGTGKFDIIQLVVYIGSTLSYFGLAAVFIDLIDTYSS
P2X7v04 ENNVEKR TLIKVFGIRFDILVFGTGKFDIIQLVVYIGSTLSYFGLAAVFIDLIDTYSS
P2X7v02 -----
P2X7v03 ENNVEKR TLIKVFGIRFDILVFGTGKFDIIQLVVYIGSTLSYFGLV RDSL FHALGKWFG
      301.....310.....320.....330.....340.....350

```

Figure 16c

P2X7v01 NCCRSHIYPWCKCCQPCVVNEYYYRKKCESIVEPKPTLKYSFVDESHIRMVNQQLLGRS  
P2X7v04 NCCRSHIYPWCKCCQPCVVNEYYYRKKCESIVEPKPTLKYSFVDESHIRMVNQQLLGRS  
P2X7v02 -----  
P2X7v03 EGSD-----  
361.....370.....380.....390.....400.....410  
  
P2X7v01 LQDVKGQEVPRPAMDFTDLSRLPLALHDTPPIPGQPPEEIQLLRKEATPRSRDSPVWCQCG  
P2X7v04 LQDVKGQEVPRPAMDFTDLSRLPLALHDTPPIPGQPPEEIQLLRKEATPRSRDSPVWCQCG  
P2X7v02 -----  
P2X7v03 -----  
421.....430.....440.....450.....460.....470

Figure 16d



24(31

P2X7v01	SCLPSQLPESHRCLEELCCRKKPGACITTSELFRKLVLSRHVLQFLLLYQEPLALDVS
P2X7v04	SCLPSQLPESHRCLEELCCRKKPGACITTSELFRKLVLSRHVLQFLLLYQEPLALDVS
P2X7v02	-----
P2X7v03	-----
	481.....490.....500.....510.....520.....530
P2X7v01	TNSRLRHCAIRCYATWRFSGQDMADFAILPSCCRWRIRKEFPKSEGQYSGFKSPY
P2X7v04	TNSRLRHCAIRCYATWRFSGQDMADFAILPSCCRWRIRKEFPKSEGQYSGFKSPY
P2X7v02	-----
P2X7v03	-----
	541.....550.....560.....570.....580.....590

Figure 16e

25131




Figure 17

26/31




Figure 18




Figure 19a

28131





Figure 19b



30131




Figure 21




Figure 22



SEQUENCE LISTING

<110> NeuroNova AG  
<120> A Method for Diagnosing and Treating Affective Disorders  
<130> XXX  
<160> 111  
<170> PatentIn version 3.1  
  
<210> 1  
<211> 56580  
<212> DNA  
<213> Homo sapiens  
  
<220>  
<221> exon1  
<222> (3000) .. (3124)  
<223>  
  
<220>  
<221> exon2  
<222> (24841) .. (25009)  
<223>  
  
<220>  
<221> exon3  
<222> (26134) .. (26202)  
<223>  
  
<220>  
<221> exon4  
<222> (30958) .. (31030)  
<223>  
  
<220>  
<221> exon5  
<222> (32481) .. (32577)  
<223>  
  
<220>  
<221> exon6  
<222> (35416) .. (35496)  
<223>  
  
<220>  
<221> exon7  
<222> (36113) .. (36242)  
<223>  
  
<220>  
<221> exon8  
<222> (37541) .. (37677)  
<223>  
  
<220>  
<221> exon9  
<222> (45470) .. (45560)  
<223>  
  
<220>  
<221> exon10  
<222> (47229) .. (47295)  
<223>  
  
<220>  
<221> exon11  
<222> (47380) .. (47529)  
<223>  
  
<220>  
<221> exon12  
<222> (50438) .. (50539)  
<223>

<220>  
 <221> exon13  
 <222> (54392)..(54889)  
 <223>

<400> 1  
 tgtggtccca gctactcagg aggctgtggt aggggggatca ctggagccca gaagtttgaa 60  
 gctgcagtga gctatgattg cgccactgca ctccagcctg gacgacagag caagaccag 120  
 tctctaaaat aaattaatta aattaatttt ttaaaaagaa agataagata aagagtcata 180  
 gaagtacaaa tgagaaacca ggagaacatg ggaccagga aagtcatgga aaccaaggga 240  
 aaatactggt ccaagatgca gaaagtagcc agtgatgttg aatgcatcca agagatcaag 300  
 taacataagg aataaaatat atccaccag tctggcaact gagaagtctt tggtaacctc 360  
 attgagaaca gaatccatgg agtggaggag gtagcagtga tgaaatagcg gacaagggct 420  
 gaaggagtgg agaggggaga gcatgaggga ccagtgcag agtgcagcag aaagacaaca 480  
 aacaggccgg gtgtagtggc tcacgcctgt aatcccaaca ctttggtagg ctaaggcggg 540  
 cagatcgctt gtgaccagga gttcgagacc aacctagcca acatggtgaa accccatctc 600  
 tactaaaaac acaaaaatca gctgagcgtg cgcatggctg taattgcaga tactcatgag 660  
 gctgaggcac aagaatcgct tgaaccagg aggcaggggt tgcagtgacc caaagtcttg 720  
 ccactgcact acagcctggg caacaggggg agactccatc tcaaaaaaac aaaaaaaaaa 780  
 aagaaagaaa gaaaagacaa agaaagaaag aaagaaagaa agaaagaaag aaagaaagaa 840  
 agaaagaaag aaagaaagaa agaaagaaag agaagaagg aaagaaagac aagaaaatag 900  
 atgtctttga gtggtagcaa gagcaacact gaagtgattt gcctattctg gacatttcct 960  
 ataaatggag tcgtaaaaga tgtatccttt tgtgtctggc ctcttttact taagcgtagt 1020  
 gttttcaagg ttcattccatg ttggagcatg ggtcagtgtt tcattccttt ttatggctga 1080  
 ataattttcc cttgtatgga ctttcgtgct tttgtgtatc catccatcag ctgatggaca 1140  
 tttgacttgt ctgcctgttc actgccatat ctactattcc tgggtgctctt gggaatgagt 1200  
 gaatgaatag cataaacaga gctgtccaag gtcacagagc tggtaagatg tgaagccagg 1260  
 atcggaagcc aggccattag tcccctagag cctatgttct aagcatcagg ctttacctgt 1320  
 gaatctctc tttttacaga tgaagatgac tgtatcactc agattcccgg caggaaagca 1380  
 atggcatact caagtggggt aactaatgat ggaaccattt acaaagggtg ggacagagtt 1440  
 aagaaaaagc aataggagat agtgagcttc ctggggctgg taagagtggg gagcccttac 1500  
 cactcccagg actaaaggag ggagtgggtc ccagaagccc tgcctatatg caactgagaa 1560  
 gggcagggcc agggagtcac gtccatcctc actgctctcc agtctcctga actggaagcc 1620  
 agaaggtag gggaacctg atgcagtgtg tatgtgtgag aaagtaaat tagtttagac 1680  
 tgaaaaactg aaaatctacc cggccactta gcaggctgga ataacagaaa tggatcaagc 1740  
 cagctgtaaa gataacaggg aacaataatt ctctgtagct gtaaagtgat aatacaacc 1800  
 tgcattcttg agtgactgct gaaacattgt cctttaaaat cagagacctt cagaaacttc 1860  
 gctgtttgaa attacatgac taagactgaa atattccaat tttgcctgga agatttaagt 1920  
 catcttgaca cagagaagca gcctcaattt acaactcagg agcagagctt cagataaaga 1980  
 ttttctggac acatttgaca tgtatcttag ctatgttgct tcctaggaaa cagggccctg 2040  
 ggtcctcttt gcaatccaga ctgaagttga ctgctttgta caaacctgtt ttgctttgag 2100

tccatcaaaa	catgacttca	tttagatfff	atctcaactc	cactttcctc	ggaatcctat	2160
actaaattgc	tgttttcctt	tgtttggtga	tgtgcgtagc	tcttctggtg	ggtggtgtcc	2220
ctcactgaat	aggtcaacaa	acctaacttt	gttggtactgc	cactgtgtcc	ctggtgatct	2280
ttggctgatt	ggtctaggtc	atagatcgac	ctgccggggt	gcagaggagg	gtggagagta	2340
actcagaggg	tcaagcatga	aagatctggc	agaaaaataa	agccctcca	ccccaccac	2400
ccctaccct	gcaaactctga	tttccccac	caactgcaga	ccagagtatt	ataagggcg	2460
gtggaagaag	agggggagat	cttcatttac	ccagagctcc	tatacatcag	gggctgaata	2520
aagggttgta	gaaatgaatg	aatcaatctc	tgagtggggc	ttcaggcagt	ggaaagatct	2580
cagtcctfff	ctgaggcata	atggaagctc	ccagtcttgt	gacatttgca	aggctgcccc	2640
tttctcccaa	gagacatgag	acaaaaaag	tgaaaggaaa	ggggggaaaa	gggagaattc	2700
tgaaaatgcc	catcctctga	acaccatctt	tgtgtaggca	tctgggggag	gccagctggg	2760
gtgaggtcat	ctgccagcca	ggcccgtagg	acttggcgct	tcttgtttat	cacagccaca	2820
tgtggggcca	ctgccagggc	ccgccccaac	tctgcagtca	ttggaggagc	ttgaagttaa	2880
agactcctgc	taaaaaccag	tacgtttcat	tttgagttta	ctgggagggg	gcttgctgtg	2940
gccctgtcag	gaagagtaga	gctctgggtc	agctccgcgc	agggagggag	gctgtcacca	3000
tgccggcctg	ctgcagctgc	agtgatgttt	tccagtatga	gacgaacaaa	gtcactcgga	3060
tccagagcat	gaattatggc	accattaagt	ggttcttcca	cgtgatcatc	ttttcctacg	3120
tttggttaagt	gggatctggg	gaggaccag	atctctgcag	tggccgacag	cacagaaagc	3180
cccagcgggc	agcttcaggc	gcacattctg	aatctcacat	ggttttcgaa	tctgagacgt	3240
gctctcacag	ccagctgggc	gggagggagg	aagcagcagc	aggcaagagg	aaacggtgcc	3300
aggctgcagc	agagagaagc	cacaggacaa	gcgggattcc	tttctgctct	acttcaggcc	3360
cgccagggcg	cgcagggcag	ggcgtgcctg	gggaaggtag	gaaagcgag	ggcaacaccc	3420
tggatcccca	gggaggaggc	gaggatctca	gggcacgcct	ggtgatcatg	ctggcatctg	3480
agtcaccatg	cttgggagga	ataggaccag	gcttgaaaat	gtgttataac	tttaggtcct	3540
caccaacgtc	aggaaggccc	tgctttttgg	tttttgtttc	ttctaaaaga	aacttactga	3600
gatataatff	atacaccata	caattgaccc	atttaaaggg	taccatttaa	tgattttcag	3660
attattccca	gagttgtgca	accatcccca	aatcaatff	tagaatatff	taatcgactc	3720
aaaaggaatc	ccacactcct	tcacatcat	ttccaacact	gttcctcctc	cttcccaccc	3780
atcaatffcc	tttctgcctc	tatggatttg	ccgattctgg	acatttcata	taaatggaat	3840
cacataatat	gtggtcctff	gtggcactta	gcgtgttttc	aatgctcatc	catgtttag	3900
catgtgttga	tacttcattc	aatffttfff	tttaaagaga	cggggtctca	ctatffttgtc	3960
caggctggtc	tcaaactcct	ggactcaagt	gatctgcctg	cctcggcctc	ccaaagtgtc	4020
aggattacag	gcgtgagcca	ttgcacccgg	ctgatacttc	attcctfftt	acggctgagt	4080
agtactccat	tgcatggata	gaccactfff	ttctatccat	tcatccattg	atggacattg	4140
gggttgtttc	tctffttttg	ctatcatgaa	taatgccaca	tgaacatttg	tgtacaaggt	4200
tttatgtgga	tatatattct	cctttctctc	gaatatgtac	ctaagagtaa	aaattgctag	4260
gtcatatgtt	aactatgtff	cacctttggg	gggaatgtgg	agctgaatff	cacagcagct	4320
gcagttfftt	acattcctat	cagcagagta	tgagggatcc	aatffctcca	catcctcacc	4380

aacgcttggt atcgctctgc tttttgggtt ttgttggtgt cattttgttt ttgtctttga	4440
gatgaagtct tgctctgttg cccaggctgg agtgcagtgg cgcaatcttg gctcactgca	4500
acctccacct ccccggttca agcgattctc ctgcctcagc ctcagagta gctgggatta	4560
caggtgtgcg tcaccactcc tctaataatt ttgtattttt agtagagatg aggtttcgcc	4620
atgtaggcca ggctggtctc aaactcctga cctcaagtga tccgcccacc ttggcctccc	4680
aaagtgctgg gattacaggg atgagccgcc acgcccggct gattgtctgt cttttttatt	4740
atagccatgc tagtgggtgt gaagtggtag ttcatgtgg ttgtgatttg aatttccctg	4800
atggtgagtg cctcttattc tctgtgctga ttgataatga tgatgaaggc aatttgatc	4860
tatagagtgg cagtgtagtt tactaagagt tagggtaact tatttcatag tactggctat	4920
gtcttctggg ccaagtcatt aacttctctg agcctcagtt tctgcatctg ttcataggg	4980
tgtggcaatt aacaaaaaaa aaaaggcatg aacagccctt atcatgatga ctgacatagg	5040
ataagagctc cataactagt atctattttt aaaaaataatc tttttaagtc tgggagtggt	5100
ggctcacacc tgtaatccca acactttggg aggccgaggg gggaggatca cgaagtcagg	5160
agtttgagac cagcctggcc aatatgggtga aaccccatct ctactaaaaa tacaaaaatt	5220
agtggggagt ggtggtgcac acctgtaatc ccagctacta gggaggctga ggcaggagaa	5280
tcgcttgaac ccggaggcgg aggttgagct gagccgagat caagccactg cactccagcc	5340
tgggtgacag agcaagactc catctcaaaa taataataat agtaataatt tttttgatta	5400
tataatagta tatatgtata taaaatacat gtatgtattt ttatctatat cctctgctct	5460
gacctcaaaa gtaaccacgt ccaagttcag gatttgaaat ctggaaacgt ggattcaaaa	5520
atccttcacc tctttgagcc ttggtttcat catctgtaaa atggggagaa ttgttgatag	5580
gaatattaaa tgaactaata aatgcaaagc tgtttgagaa atatatggca tatagtaatc	5640
cctgattaag tgtagttctt tattattaat aatgctatta ttaggattat tattattcga	5700
ttcatatgtt tactgttcaa caaatattga atgataaaca tatatgctgg gtccggcatg	5760
gtggcccatg cctgtaattc cagcactttg ggaggccaag gcgggcaggt cacttgaggt	5820
caagagtttg agaccagcct ggccaacgtg gtggaaactc cgtctgtgct aaaaaataca	5880
aaattagccg ggcaggtggtg tgggtgcctg taatcccagc tactcgggag gctgagacag	5940
gagaatcact tgaaccagg aggtggaggt tgcagtgagc caagattgca cactgcact	6000
ccagcctgag ccacagagca agactctgtc tcaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	6060
tatatatata tatatatata tatatatata tagtattttt agtagagatg gggttttgcc	6120
atctcttata tatttttata tatttatatt ttttatatat ttatatatat ttatatgtat	6180
tttatatatt tatatatatg tgctgaaatc taaaataaga catatcactt aaaccagcta	6240
tcacaggacc aatggtaaag agcataatta gtaaatgccc cttctgcat taatacgtag	6300
attgtatgtc acaatttcat gtagcaggtg atggggcttc ctgacattct tgggcccagc	6360
atacaatctt tttatttttt ttttagagac aggtctcat ttgtttgccc taggctggag	6420
tgcaagtgtca ttatcacagc tcgccacagc ctcaacttct caggctcaag tgatcctccc	6480
gcctcatcct cccaagtgc tgggaccata ggcacagcc atgatgtctg actaattttt	6540
tatttttttt ttgtggagatg ggggtctccc tatgttgctc aggttagtct caaactcctg	6600
gactcaagt atcctcctgc tttggcctcc caaagtgtg ggattatagg catgagccac	6660

tttgtctggc	ctacaattcc	ttttctttct	ttctttcttt	tttttttttt	tttttgagtt	6720
gagtctcact	ctgtcaccta	ggctgtagtg	cagtgggtgtg	atctcagctc	actgcaacct	6780
atacctccca	ggcccaagca	atcctcccac	ctcaacctcc	cgagtagctg	ggaccacagg	6840
tgtacaccac	cacgcccagc	taattttttg	tatttttggg	agagacaggg	tttcaccata	6900
gttggtccagg	ctgggtcttga	actcctgagc	tcaagcaatc	caccgcctt	ggcctcccaa	6960
agtgtggga	ttacagggtg	gggtcaccat	gcccagcctt	caatttcttc	agattattgt	7020
ataataataa	aatttctaaa	aacagattac	tatgagaaca	ccttgtaatg	ataaatattt	7080
ttctttcaca	tggtgaaatc	ataacttaca	tgtatttaac	tccctcaaat	tcagttatac	7140
atgttccatg	gggacatgga	gagagagaga	acaatataaa	taatgcaatg	attcagcaac	7200
cgggttcgct	ggagtaggtg	tgagatggga	tactggcatc	tgggttccc	tccgtcaatc	7260
tctgttccca	cgtgcttttt	agcgagcatc	tgtgtgtctt	acttattaat	ctaggtacat	7320
atttttcage	cacatgaccc	ctaaaataag	tcaactcctt	caccttcacc	tggagcaaaa	7380
ctgtaaaaac	agcaacttgt	tctagatctg	gaggcaaaat	tggttccagc	agaccactg	7440
ggaagtggga	ggtgtttgct	atgctttgtg	gacaatttaa	gtgtttaaaa	ggcaattttg	7500
atgagccatg	attttcatct	taagtgccaa	ctttagaaga	taaccttcat	ataaaagttt	7560
ctctacagtt	tgtatttttaa	aatataaaaat	ggaagaaaaa	actatcacca	gctttgcttt	7620
tttctctctg	gttaatggac	tatctattaa	cgtctcacta	aatggccaaa	tggagtctca	7680
cagtacacag	tttgggaaat	gctgccttaa	gttgctctgt	gaagccttaa	gttcgtgtaa	7740
ccatgactta	gccatctgtg	aaatgggtag	acccatgggg	cataggaatt	ggggaatgta	7800
ttttgcatca	agtaggactg	gtgtggacca	cagggtgaaa	tcagacaggg	ctcgagcaat	7860
gttcaggata	cagcagcaag	tgacagggtc	tggccagctg	tccctactgg	ggctgcaagg	7920
actaagagcc	aggataagga	gacagggcct	cgggtattga	aaatgggata	caaagcaggg	7980
actggttatg	gcggaaatta	accaagaggc	tgttcagctt	cttctattct	cacaaccaat	8040
tcagatctgg	attcaaaggc	agggatgcag	ctacaagagg	agtttgtgtt	ggaggtctgt	8100
gtagatagga	tgctaagcag	gtctgtgtca	gccaggatcg	tttgggcagt	aagtgcaga	8160
aaccaactt	acgttaacca	caaagcctca	aaggggaatt	tattggatca	tgtaactggc	8220
cagttgaaga	gatgggtgctg	tcctcggacg	tggatgaaga	taagagttca	aactgtgtca	8280
tcaggacgga	tccgtctccc	tacatcctgg	ctccatttgc	ttctatttgg	cttccttctt	8340
gaacaactct	gtatgtacaa	gatatcctca	cattctcttc	ttttaacaac	ctcagaggag	8400
caaacaagtc	tttctgata	gctctgctaa	gaaagtccca	ggaaagactc	tgattgatca	8460
gatgtgagtt	acagatccat	ccctgaacca	atcactgtgg	ccagggccaa	ggtgtcatcc	8520
cattggccag	cctgggtcat	gtgtggctc	ccgtagcagg	aagtagggag	aagacgaggg	8580
agccgtgtga	gccctacgtg	aaccacatcc	agtggatttt	ccttaggaag	gaagattctg	8640
tgaccagacc	aaatggaaaa	ggaagtgggg	agggtaggca	gacaaaaatc	ggtgatctat	8700
tacaaggctc	ttccaaaagc	cagaaattag	ggatttctaa	gtgtacctct	ctagtctctg	8760
cttgctcag	ccctgcagct	gggtgtgga	gactccatca	gcaattgctg	acccctatt	8820
cttacctcgg	ccccctctg	agtcccacca	ggagccctgc	tcctcatctg	ggacatgctt	8880
cttggcccat	ggaggttgcc	agctcacctt	ggtactttac	tactatttgg	tttctgtgac	8940

caccaagctg ttgcttaaac caagccgcct atggtggcac cactttggag aattcttttt	9000
tattttttga gacaggttct cactctggtt accgagactg gagtgcagtg gtatgatctt	9060
ggctcactgc agcctcgacc tcccgggttc aggtgatttt cccacctcag cctccccagt	9120
agctgggact attggtgcat gccactacac cgggtcatt tttgtatttt tagtagagac	9180
ggggttttgc catgttggcc gggctggtct tgaactcctg acctcaggtg atctgcccc	9240
ctcggcctgt caaagtgctg aaattacagg cgtgagccac cgcacccggc tggataattc	9300
tttagatttt attccaagtg ggttgagaag ccatgggaag attgtaagca aaggaataaa	9360
agttgggtgc attctctttg gaggatttcc ctcttcctg gagaccagtc ctcacagaag	9420
cttttccaat ggccacagaa gactaaggaa gattgagaga ggcacaggaa acaattccac	9480
tcagagctgc tgcttcgcca ctgcccagtc agcctccact ggcgccaaca cgagcctatg	9540
ctgtcattcc tgggcccagg ctggctgtcg ctgtggagat ggtgctgctg tgctagtttg	9600
tggctgttaa aatggagctc ctggctggac acctgtttcc cagtgggtgg ccacaaacct	9660
gtgagtctgg gcctcctcca gctgagtaga agtgagccca gcacagaggc cggggaggcc	9720
ccagacgcca ggaggagcaa atccaaatcc aaccaagca tggccctct gacctgacc	9780
atcttttttt ttttttttt ttttgagaca gagtctcct ctgtcaccca gactggagt	9840
cagtggcacg atcttggctc attgcaaccc ccacctcctg ggttcattgc attctcctgc	9900
ctcagcctcc ctagttagga ctacaggcat gcaccactac acccagataa tgtctgtatt	9960
tttagtagag atggggttct gctatgttgg ctaggctggt ctgaactcc tggcctcaag	10020
cgatccatcc gtcttggctt cccaaaatgc tgagattaca ggtgtgagcc actgcgcca	10080
gcctgcccct gactgccttc agaaaagtag aagcaggagg tctgccacat tctcatccca	10140
tccttgacac ctgggtgatg tgctagagct tgaggggaag gtcagaggta acccagttcg	10200
ggaagtgacc ccgtaggagg agaagaatcc agccacgtga cacacagtac gaagagaaca	10260
gtaaatgcaa agacctggaa acgaggaaca gcaaggacag catgtcccag aacctgggga	10320
gagaggaggg agcagccgga gctgagggca gctcagggga agagttgcta tttgattttc	10380
agtgcattgg aagccactga aaagttttat gcagagaagt aaaaccatct cactcatgtt	10440
tctagaagcc ttatgggcca ctgtgcagag catagagtgt ggtgggacaa gaatgacggt	10500
gaggaaccag tgagaagggt cctggagtca ccagatgcga ggagatagag gcttgggcta	10560
gaatgggctg gcggagctgg agagaggtgg acaaagctga gatgtgtttt ggaggcagaa	10620
atgatagatt tgccaggagg tgaaagatgg ctctagattc ctgaggaacg atgatgcctt	10680
ttaccaacct gaagaagacc aaggagaggc ttgaggacta acacggaact gtgaaggagg	10740
ccaggcaaaag gtggagactg ccgtgagctg tgatcacgcc actgcaccct cgctgggtg	10800
acagactgag accctgtctt aaaaatata tgtaataaaa aataataaat aaagggccag	10950
gcgtggtggc tcaacgcctg taatcccaac actgggaggc tgaagcaggt ggatcacctg	10920
aggtcagaag ttcaagacca gcctggtcaa catggcgaaa tcccatctat aataaaaaa	10980
caaaaattag ccagacatgg tggcacatgc ctgtaatccc agctacttgg gaggtgagg	11040
caggagaatc acttgaacct gggaggcaga ggttcagtg agcctaaatc acaccactgc	11100
actccagcct gggcaacaga gtgagacccc ataaataaat aaataaatat ataaataaag	11160
gatatcaggc aagcaattaa aacacctgcc tactaccatc tatccgcatc ctttttttca	11220

tgtctaacac caaaacacca aaatagctgg aaacatatca tgtcaaaaaga taagcatgat 11280  
 tctttcaaat gaatggctgt ggacttatga aaagctgcct ccgcgtctga ttttcttgtt 11340  
 tcaaggggta atgatggagg tttctctcct gcacctgcag gcctccgggg atgctgcccc 11400  
 agtgcagcct cccaccctgg gctggagcct cccagccccg ctgcatgggt agcacagtgt 11460  
 tgggaagagg aaggagcagt ttgtctggct ttttcacagg cccacagctg gccggtacag 11520  
 tggcagagaa ggcgaggctg aggtggggct ggggaagcca gcaggggcct gatcatggag 11580  
 cgccttgagc tgcattccac aggtactgag aaagcatgga gagctgtccc ccagcccca 11640  
 gcatcttatt agaaaatatt gaaacataac agaaaacatg aagggtttta tgttcatatt 11700  
 ttcatttttag agagatcatt ctgcttgaat tataacacct actgagtact gctgaggact 11760  
 taccatgtgc cgggttctgt tcttagcatg tgacatgtat tggcctattt gattctccta 11820  
 cataatcctg gaagatttgt tgtgggagtg ggtgatcatg acctgaacag cctaaccatg 11880  
 tttagggaaac actctgtgtt atgcgttcca ccacacaatt catgtcacag ctgaatgtcc 11940  
 cagtttccct tgcagctagg acacaacggg agacccaagc tttccaaaca tatgccccag 12000  
 gcaacacttt gatacacaa gaataacact ggatgctggg tgtgcacaga attcaaattc 12060  
 aggaatggag ggaagtactg ccctttgacc catacaatag gggatcccct tccttgacc 12120  
 ccatggctta gagggcccca ctctgtccct ctgcagactg tgagctgccc tatggaatga 12180  
 ggggtgtctg aggtgttatt aatttctatt tgttacacag cagtggaaac taatacagtc 12240  
 ccacaggcct aaatcaaggc gtcagcaggc tgagttcctg cctgggagact ctggggaaaa 12300  
 ttcacttcca aactcatttt tgttgttggc aaaattcaat cccttgaggc tgcaggactg 12360  
 ggctctgtc ctcactggct gtcagcctgg ggccactctc agattctgaa gtccctccac 12420  
 attccctgcc acatggtcct ctccatctcc aagcctgcta tgacacaacg cattgttctc 12480  
 atgcttcaaa tctctgactt ctctcccggg accagaagaa gaatactctc ttcttttaag 12540  
 gggctcatgt gattgggtca ggtcaactct atcatatcat ggagtctcac cgtcttgccc 12600  
 aggtgggtct tgaacacctg ggctcaagtg atcctcctgc gttggcctcc caaagtgtg 12660  
 ggattacagg cgggaactgc tgtgcccagc caaggccagc atcttcagac ctctctgccc 12720  
 tgttttcaca tgggtctctc ctctttgtgt gtgtgtcaaa tctccttctg cctctcttat 12780  
 aaggacgtgt gtaatagcac tcagggccca cctggatgac acagggcatc ctgccatct 12840  
 caaaatcggt aactttggcc aggtgcagtg gctcatgcct gtaatctcag cactttggga 12900  
 ggctgagaca ggtggatcac ttgaggtcag gagttcaaga ccagcctggc cagcatgggtg 12960  
 aaaccctgtc tctactaaaa atacaaaagt tagctgggca cgcacctgta gtcccagcta 13020  
 ctgaggagac tgaggcaaaa gaatctctag aacctgggag gcagagactg cagtgaacca 13080  
 agattatgcc actgcacttc cagcctgggg gacagagtga gacttcatct caaaaaaaaa 13140  
 aaaaaaaga tgactgggca cgggtggctca catctgtaat cccagcattt gggaggccca 13200  
 ggtgggcaga tcacctgggt tcaggagttc attcaagacc agcctggcca acatggtgaa 13260  
 agtctctctc tactaaaaaa aaaaaaata caaaaattac ccagatgtgg tggcaggtgc 13320  
 ctgtaatccc agctacttgg caggccgagg caggagaatc acttaaactt gggaggcaga 13380  
 ggggtgcagtg agctgagatc actctgctgc actccagcct ggacgacaga gcgaaactcc 13440  
 atctcagaaa aacaaattaa ccaggcatgg tgctgcgtgc ctgtagtccc agctactggg 13500

aaggctgagg tgggaggatt gcttgatcct gggagggtga ggctgcagtg agctgtgttt 13560  
gtgccactgc actccagcct gggcgacaga gtgagaccct atctgaaaaa aaaaaaatcg 13620  
gccaggcgcg gtggctcacg tctgtaatcc cagaactctg ggagactgag gtgggtggat 13680  
cacctgaggt caggagtctg agaccagcct gaccaacagg gtgaaacccc gtctctacta 13740  
aaaatacaaa aattagctgg gcatgggtggc aggcacctat ataatcccag ctactcggga 13800  
ggctgaggca ggagaatcgc ttgaactcag gaggcagagg ttgcagtga cagaggtcat 13860  
ggcattgcac tccagcctgg tggcagagtg agactccgtc tcaaaaaaaaa gaaagtctct 13920  
ttttacaact tttttgaggt ataacttaca tatcagaaaa tcaccggttt taaatataca 13980  
tttcaacgac ttttagtaaa tttcccaagt tgtgcaacca tcatcacaat ccagtttcag 14040  
aacatttcat cagcccagta agagccctca caccattaa cagtcaactcc ccactccac 14100  
ttcctcctgg tggttgctca acccaggtaa ccatgaatct attctctgcc tctggagatt 14160  
tgtcttttct ggacaatctt tcttcttaaa ctccagcagag tgagttctgt tgtttagaac 14220  
tagaaacccct gcccaataga atactaatat cctcttgttt tgcaaggagg ttagagtctc 14280  
gctctgtcgc ccaggctgga gtgcagtgg atgatctgc ctcaacttcga cctctacctt 14340  
ccggattcaa gcgattctcc tacctcagct tctgagtag ctgggattac agacgtgcac 14400  
caacacacct ggctaatttt tgtactttta gtgagatgg ggtctcacca tgttggccaa 14460  
gctggtcttg aactcctgac ttcaaagtag ccccccgcct tggcctccca aagttctggg 14520  
attacaggca tgagccactg caccagccc taatatcctc ttattacaaa agaagaaaca 14580  
atgactcaga aagattaagt cacttaccca aggtcatgta gtcagtaagc aatagagctg 14640  
agactgaacc tcaactcaaa gtctatgctc tcaactaaagt attgcagtgt agagagtaga 14700  
ggggacaggg tgagactgga ggaaggagg gcathtagga gactattggg atgtccagat 14760  
ctgagagctg ttggcgacct ggactgggga ggtgatagag gcagaggaga gaagtagatg 14820  
aatttgagca tattaggaag tgaaacagta ggatttgga actgaacggg tgagtggaca 14880  
agaaggagag gatgagtcaa agagtgcacc atgcttctgg cttgggcaac tgatgggtac 14940  
tggaataact tgctgggata ggaaatgcag gaaaaacagc caatttggg tggaagatga 15000  
ggtcaggact aaaggcattg agtttgaggt gtctatggaa tgtacaaatg gagatgttga 15060  
atggatggtt ggatatacga atctgcagct gagatgtagg tttctatata accagcatat 15120  
agacggcaac cagaacctg gaagaagggt caaccgacca gcaagagaag atgtagatga 15180  
gaaagcaaaa gggcctagaa cacgcccaca aggagtgtta acctttaagg cgtgggagcg 15240  
cagaagtccc caaaagattc tgaggagtag ccacggcgat ccagtgaag tcaggcacia 15300  
gtggactcct gcaagctaag ggaaggaggc gaggaagtg gggcatggc acagtcaagt 15360  
tcaactgctg ctgagagttc caggaagggt aggtccatgg tgtatctatt ggatttagtg 15420  
acaacgacat tgttgtttac ttggcaagag aggtctttaga gaagggatgg gggtagggagc 15480  
cagactagag tgggtagagg agtgagtga tcaagatgag gaagtggatg tgatgactcg 15540  
ccctttcttt cttctttctt tctttttttt ttttttgaga tgcagtcttc ctctgtcgcc 15600  
caggctggag tgcgatggtg tgatctccac tcaccacaac ctctgcctcc tgggttcaag 15660  
tgattctcct gcctcagcct tccaagtagc tgggattaca ggcatgtgcc accacacccg 15720  
cctaattttt tgtattttta gtagagacgg ggtttcacca tgttggccag gctgggtctca 15780



aactcctgac ctcttgatcc acctgcctca gcctcccaaa gtgctgggat tacaggcgtg 15840  
agccaccgta cccggcctgg tgactcactc tttcaaggga ctcagctata aagaagagag 15900  
gcatggagtg gtggtaggat ggattaagct ggggagggga ggcattttta agatgagaaa 15960  
tgatgtttaa atgcagatgg aaatgggaca ggagagaaag ggaagttgat gggataaatg 16020  
aaaattatca tcatcatctt cctcaccatc accagcagca accacaagga aagctcgttc 16080  
tgctctggga ctgccctggt agacatttct tccttatact gagccacaat ctatctccct 16140  
gtaactttcg cctttgggta gagttctgcc ttgtgtccct cttctactct tcaaattggct 16200  
gcacgtgct actatgtctg tcctcttctt caaactcaca gccccatcc tacgggccct 16260  
gtgctcccag cctggtgacc cgaccctcta ctgtcccagc caagttcaca tgaccatcct 16320  
cttatttgte aagcctctct tttttttttt tttttttttt tttttgggag gagtcttact 16380  
ctgtcccagg ctggagggca tggcacaatc tcagctcact gcaacctctg cctcctgggt 16440  
tcaagtgatt ctctgcctc agcctcctga atgctgggat tacaggcata catcgccatg 16500  
cctggttaat tttgtatatt ttagtagaga cgggggttca ccatgttgge caggctggte 16560  
ttgaactcct gacctccagt gatccccctg cctcagcttc ccaaagtgtt gggattacag 16620  
cgtgagccac tgagccgggc caagcccctc tgaagaggca gtgtgcacac tcttgctttg 16680  
ctgatagaag catgaaatgg ttcaagtttt tgtgatttgg gaaaatgaac tcagctataa 16740  
aatgtgcaca cccagaagtt gtcctatgga cttaatgtct taggggtgag aagaagcttg 16800  
tgcagggatt cctggcaaca tgaatgcggg atggacagac agtgagcaca gtgttgagga 16860  
gcacagcctg agtgcatctc cctgagatca ggaacaggac aaggatgcc gctttcaccg 16920  
ctgctgttcg acattgtact ggaagttcta gccagagcaa ttaggcaaga aaaagcaata 16980  
aaaggcatcc agattgtaaa ggatgaagta aaactatctt tagtcacagg tggcacaatc 17040  
ccatataggg agaattccaa agaattcaca agaaaactgt tagagcaaat aaactaatc 17100  
agtcaagttg caaggaacaa gatcaaccac aaaaattaat tgtgtgggtt ttttttacac 17160  
cagcaatgaa ctttgaaaaa ggaaattcga aaaagtattc cattcactgt agcatttaag 17220  
aaagctgagg ctgtgaagct acatggctgt gtttgagct ggtgctgcta tttaccatgt 17280  
gaccttgga cagcttctca cttcattcat gacaccgtt attcagctat taagtggggg 17340  
taataatagt accttcctta tagagttggt tttttgtttt tgtttttgtt tttggtgggg 17400  
gagagtaagg attgcaagag ttaatatatg ttagcaaaat ggacttagaa gagtgggttg 17460  
cacatggtaa ggactatatt agggttacca aaaaatgttt ataattggga aaattgaaca 17520  
aaggttccag taaatatcca tcccagaggc tgggtgacaa attatcacia tgtacctttt 17580  
agcttgaacc atatggaact gcctatatc tatagctatg gcctacaaaa tgcaaatgg 17640  
gcagtttttt ttgttgttgt tgtttgtttg tttgtttgtt tgtcttgaga cagtctcagt 17700  
ctgtcaccag gctgaagtgc agtggcacga tcttggctca ctgcaacctc tgcctcccag 17760  
gttcaagcga ttctcctgcc tcagcctccc aagtagctgg gattacaggt gtgcaccacc 17820  
acgcccagct aatttttgta tttttagtag agatgggggt tccaccatgt tggccaggcg 17880  
ggtcttgaac tcctggcctc aattaatctg cctgcctcgg cctcccagag ttctgggatt 17940  
acatgagtga gccaccgagc ccagccaaaa tgggcccgtt catatgttca atttaaacac 18000  
taagatactc ttttagccact gaaaatgact gtgtatcata gatctatctt tatcacagat 18060

ctatctttac tgtcatagaa agaactaata acatattatt gaacaaaaat aacaagttac 18120  
 aaaacagctt gcagccacgt ttgtaacatc agtatatcta tttgtgtgtc taggtgtgga 18180  
 cataaagaga tatctagaag aactcccat tgcctatggg aatgacttgg gctgggatct 18240  
 tttttattta tttatttatt tttttattta tttatttatt ttttttgaga cagagtctca 18300  
 ctctgtcacc caggctggag tgcaatggca tgatcttggc tcaactgcaac ctccgctcc 18360  
 caggttcaaa cgattctccc acctcagcct cccaagtagc tggaattata ggcacccatc 18420  
 accgcaccta gctaattttt gccttttttag tagaaatggg gtttcgccat gttgcccagg 18480  
 ctggtctcaa actcttgatc tcaaattatc cacctgcctc agcctcccaa agtgctggaa 18540  
 ttacaggcgt gagtccaccg acccagctgt gatcttaatt tataattttt ctttttaacc 18600  
 cttgattttt accctgtttg actgacttta ggagtctgac ccccaaaact gcaaaagaat 18660  
 aaattagtgt tgttgaaagc cactaagttt gtggtaat ttttagagcaa caaccagaaa 18720  
 ctaatacaat cgggattaaa cttgccttgg ggggtctggc tctccccac tttcttctca 18780  
 gagcatctct tgggtcctcg ccccttctg agccccacc atggccgggt ggaccagcga 18840  
 tggacctgat cggattctcc tcttgggatt ttaaataaga gatgcagagg ctgcacgcca 18900  
 agcttctctg tgccgcaact caggcacagt cctggcgcta ctgttcttta ggggtccatg 18960  
 aagatgtttg ttggcctggc ttaggtgtga gctaagagat acaagggata cactttggag 19020  
 gctcagccct ttgatgttac ctttgcgctg gaagcccagg ggctgtaaga cggagaggaa 19080  
 ctttgcgtga agggagggtcc tccctgcaga gcttgaagag caatctaacc atgctggcat 19140  
 ggggcccatc ccacctgtga tgccagtga agtctgtgac agtcaactcat attcagagca 19200  
 tgtggattga gactctgaag aacgacttta tgggtgggac cttcatctcc atctgggtgg 19260  
 tccagagcaa gcagcctcag ggtcccatgg gcccaacctc acctccatc cttatttccc 19320  
 tgccactcca gggctcctcc tcagccagac ttgcctgtga gtgccccttg atggctgtat 19380  
 tagtctgttc tcacgctgct aataaagaca tacccaagac caggtaattt ataaaggaaa 19440  
 gaggtttaat gggccacag ttccacatgg ctgggaaggc ctcacaatca tggcagaggg 19500  
 cgaaaggcat gtcttacatg gtggcaggca aaaagtgaat gacaaccaag caaaagaaga 19560  
 aacctcttat aaaaccatca gatcccatga gacttactca ctaccacgag aacagtatgg 19620  
 gggaaactgc ccccatgatt caattatctc ccaccaggtc cctcccacaa cacatggaaa 19680  
 ttatgggagc tataattcaa gatgaaattt ggatggggac acagccaaac catatcagt 19740  
 gccctggctt ctctccatt agctctgttc tacttcccaa aatctttcca gctttgttct 19800  
 tttttagaga ctggagtctc actgtgttgc ccaggettgt ctggaactcc tgggctcaag 19860  
 ggatcctcct acctcggctt aacaagtagc tgcgaaatata ggtgcacacc accacacccc 19920  
 cacaacttac ccacccctcc cggctcagt aaatccctag gtccttcaca ctcccgtag 19980  
 ctccaggtac cagcggcttt gccttctaca gaatgcataa cattatagcc tcttagagca 20040  
 aaaaggctcat ggggatggg aaggctggga gttctctctt cattcaggga ttaaacctac 20100  
 cacctggagg ggtgacatgg ctcaccttag tcaccaagca aggtcgggag agtatttctg 20160  
 ggatggctcat tagcactttc ttcccaaagg gaggatacta caaactcact ttcattgatca 20220  
 cagtatcccc cccgacaggt aaagtgtgca aactcgtttt cattcccttt ttgtctttga 20280  
 cgcagctctg aagcatttta attcttaaaa ttacatgtt catgatgaca cagagacacg 20340

ttttctgttt	aggagagggc	ctgattccat	taaggctgat	cgatcatatg	ttgcacattc	20400
tgggataaat	atgagagatc	agaattgagc	ttcccaaagt	gtgggtcaca	tgaccagtg	20460
ttgttccagg	agaacgttgc	aacatccagc	atgagaaagc	gctgcatgca	ccatgtatgt	20520
gttgggatgc	tgggctgaaa	tgtattaatg	agcaagtgtt	atcatatttg	tctatggaat	20580
tctatttaat	agctatgtct	cgtatgtgct	gattgggtct	caaactctggg	tatcttacct	20640
caacatcttt	cttgaataat	cttcttttta	aataaatgtt	ggccgtgtgc	actggcttat	20700
gcctgtaatc	ccaacacttt	gggaggccaa	ggcgggtgga	tcacctgagg	tcaggagtgc	20760
aagaccagcc	tgaccaacat	ggagaaaccc	catctctact	aaaaatacaa	aaattagctg	20820
ggcagatgg	tggatgcctg	taatccagc	tacttgggag	gctgaggcag	gagaatcact	20880
tgaacctggg	aggcggagg	tgcagtaagc	tgagatcgcg	ccattgcact	ccagccttgg	20940
caacagagt	agactccgtc	tcaaataagt	aataaataaa	tggtatagaa	atactaaagt	21000
gtattgtgca	taatataaca	cccacgtacc	accatccaac	ttaagaaata	aagcattact	21060
aatccagttt	tgctccctgc	aaactcatcc	ttcatcccat	tctctttcct	tttgctttat	21120
aaacatcctg	cagccaggca	cagtggctgc	acctgcagcc	tgagctactt	gggaggctga	21180
gatgggagga	tcacttgagc	ccaggagtgt	gtggctgtcg	tgtgctgatt	gcacctgtga	21240
aatagccact	gtgttccagc	atgggcaaca	tagcaagact	ccatctctaa	atgaaatcct	21300
tttttttttt	tttctttttg	agacagagtc	tcactctatc	accaggctg	gagtgcagtg	21360
gcgccatctc	ggctcactgc	aacctccgtc	tccagggttc	ctgcctcagc	ctccggagta	21420
gctgggacta	caggcacatg	ccaccacaac	tggctaattt	ttgtattttt	agtagagacg	21480
gggtttggcc	agggttggtt	cgaaccaggg	acctcagatg	atccacctgc	ctcggcctcc	21540
caaagattac	aggcatgagc	caccatgccc	agcctaaaag	aaatcctgaa	tttgacattt	21600
ttaattccct	tggttggttt	aatatttcta	cagcacatat	atgtctacat	atataaaaaa	21660
tataattttt	tattgggttg	aaattcacat	aacgtaaaat	taaccactgt	aaacatgaac	21720
aattcagtg	catttaggac	attcatagta	ctgtgcaacc	accacctcta	tctagctcca	21780
gaacattttc	atcaccccaa	aaggacactc	catgcccccc	aggagtcact	ccccatttct	21840
tccactctg	tcccctgcaa	cgaccaatct	actctctgtc	tctgtggatt	cacctattct	21900
ggatatttca	tataaatgga	atcatacaat	atgtggcctt	ttgtgactgg	tctctttgac	21960
ttagctgggt	gtggtggcac	gagctaccca	cctacttggg	aggctgaggc	aggagaatcg	22020
cttgaactcc	agccataaat	tatattgttt	tcaaggttca	tcaatattac	tcgccagggc	22080
tggagtgcag	taatgtgacc	ttggccccct	gcaacctcca	cctccccggt	tcaaaggatt	22140
ctctgectc	accctcccga	gtagttggga	ttacaggcgc	cccccccccc	caccgggcta	22200
attttttgta	tttttagtag	agagaggggt	tcaccatgtt	ggatcatgctg	gtcctgaact	22260
cctaattctca	agtatccac	ccaccttggc	ctccaaaat	gctgagatta	cagggtgtgag	22320
acactatgtc	aggcctagtc	aattcatttt	cactgetgca	aagcatctca	ttttaaaaaa	22380
ttattttatt	attatttttt	tgagacaggg	tctcactctg	ttgcccaggc	tggagtacag	22440
gggtgcaatc	acagcttact	atagccttga	cctccccggc	ccaagcaatc	ctccactgt	22500
ggcctccaga	gtagctggga	ctacaggcat	gtgccaccac	acacagctaa	tttttgttct	22560
ttgttttttg	ttttgttttg	ttttgttttt	tgagatggag	ctttgtctct	gttgcccagg	22620

ctggagtgca gtggcgcgat ctgagctcac cacaacctcc atctcccagg ttcaagcgat 22680  
 tctcctgcct cagcctcccg aataggtggg attacaggct cagccaccg taccagtta 22740  
 atttttatatt ttagtagaga tgaggtttca ccatattggc caggatggcc tcaaactcct 22800  
 gacctcaggt gatccatccg ccttggcatc ccaaagttct gggattacag gcgtgagcca 22860  
 ctgtgcctgg ctcaaagcat ctcatgtat gaatccaaca atttatttat ccattcgttg 22920  
 tttgagtaca tgtaggttat ttcttgcttt ccctcttaca agcaatgctg cagaaacatt 22980  
 cttgtccata gctcctccgg cacatgtaca agagcttctc tagggcaca atgtaagagt 23040  
 tttattgctg ggccatgggg aatgctcttc ttcaaatgta cttgctattg tcaaactctga 23100  
 gctatctttt caggagccta ctactggttc tcagccctgg ctatgcatta gaatcacttg 23160  
 ggaactttaa aaaaatatgt gggatatctg ataactcccc cgaccctgg aatcctagtc 23220  
 tccagggatg ggactgggga cttgaaagct ccctgggtga ttacacagcc aaggtggaga 23280  
 acagctggga ctgagtcttg gccactctg tctcagcttc ccaaggtcag ctcatagagg 23340  
 gcagccatca tcttagactt gctaataatca ctgagcagct gagcagatag aagggactaa 23400  
 ataaataaag tctgaatgag taaataaagc atttaaaaca aaccctttag ggctgggtgt 23460  
 ggtggctcac acctgtaate ccagcacttt gggaggccaa ggtgggtgga tcccttgagg 23520  
 ttgggcatte aagaccagcc tgaccaacat ggcgaacct tgtctctact aaaaatacaa 23580  
 aaattagcaa agtgtggtgg tacgcgctg taatcccagc tactcggaag gctgaggcag 23640  
 gagaatcact tgaaccggc aggcagaagt tgcagtgagc tgagatggag tcaactgcact 23700  
 ccagcccggtg ctacagagca agactccatc tcaaaaaaaa cataataaaa caaacctttt 23760  
 aggctgggtg cgggtggcgca tgccggtaat cctggtaact atctcctact gttgtgacga 23820  
 ttaaaggaga tgggtgcatg agagtacttg gcacatatga agtgcctaat aaagagtaac 23880  
 tctgcagcc tgggtgtggtg gccacacct gtaaccacaa catgttgga ggctcaacat 23940  
 gttggcagga ggattgcttg aggccaaggg tttgagacca gcctgggcaa cataaagaga 24000  
 cccccgtctc tataaaaaat aataaaaaact agctgagcat ggtgatacat gcctgtagtt 24060  
 ccagctactc aggaggctga ggtgggagga tcaacttgag ctcgaggagt ggaggctgca 24120  
 gtgagctacg atcacgccac tgccctccag cctgggcaac agagcaaggc cttttctctg 24180  
 gaataaaaaa aaaagagagg ctgggcgcag tggctcacac ctgtaatccc agctacttgg 24240  
 gaggtgagg caggagaatc acttgaacct gggaggtgga agttgcagtg agccaagatc 24300  
 gcaccattgc actccagtct gggcaacaag agcaaaactc tgtttcaaaa aaaaaaaag 24360  
 agagagagag agagtagctg ccattttcaa tgttattgat aaagcagggt ggtgtttctg 24420  
 aaaatgcacc agatgcagcc ttagactgga aggtgctgat gtgttactga gcctccagac 24480  
 aaagctggcg aaccaccggg ggcgagactt tatgacttga gaagctgtcc tatctatgag 24540  
 ggtccatgtt aggaatatt aaagaatagg attgggggca gtagctttcc tgtagcaagt 24600  
 gctagctatg tgccagatac tgggcaaagg agtctgcaga cgtctctcat ttcatcctca 24660  
 caatccatga ggcaggatg actattcttc ccattatcca gagagggaaa acaagtcaca 24720  
 cggaagcaag tcacgcagca gagctaggat tggaaacagaa gtgcctgcat cctccaacgc 24780  
 ctgcatccca acccgctgtg ctatgcctcc cgttgatgct ttcccatgtc tgccatttag 24840  
 ctttgcctctg gtgagtgaca agctgtacca gcggaaagag cctgtcatca gttctgtgca 24900

caccaagggtg aaggggatag cagagggtgaa agaggagatc gtggagaatg gagtgaagaa 24960  
 gttggtgcac agtgtctttg acaccgcaga ctacaccttc cttttgcagg tgagcacctc 25020  
 gtagcattct cccaggctcg tcgctgggtca ccgtcgccag ggcctagctc ctttccccta 25080  
 ggatctacag cctcactcca gaaaaaacgc tggctctatt ttaaaagctc tgtgaacatc 25140  
 ccaactgaga aaacctaaaa attgcaaaac tgggtgagat ccaggagatg attcttttgc 25200  
 ttatcaagtc ctacagggtt tctcaaaata gcctcatgtc ctggtgcatt gaaaacagga 25260  
 tttgattcac tcgcgcttga tttaccgcga gctttttaga cgcacagcca ccaagccagc 25320  
 aaggctctgc gagattcaga aggacagaga actttctacc aaagagcacc agcccttcta 25380  
 gattgtgtag catttcatca acaaaaatga ctccaggaga ccattcttgg gtacatTTTT 25440  
 ccacaagaa aggatgtaaa tgacaaaaag aagaaaggca tcggtcactg agagaaggcg 25500  
 tgtgacttct agatttgcaa gcagggaaaa tgaaagcaga attgccaaaa aagagaaatt 25560  
 aaagtagggg tgaggagtcc aaaaaacaga taagggggcg agagggacaa aggggcacgt 25620  
 ttgcggttg gggggctaaa atgctttgca aagacccatg aagcccaagc tgctgtttgt 25680  
 ttctaacagt gggtcattaa aatcctgtga ttgctcaggt ggtgggatgc cctagtgc 25740  
 tttcattaaa gtctgggaaa atctgaacag tgttgatgag aaggctgctc ccctaccctc 25800  
 gcctcccag gtctcctgag tttcaattta atgagatttt tactgcgtaa aaaaaaaaaa 25860  
 aaacaaaaac aaaaacatga aaaggctgct tttgagactg cattggtaaa tgactcttca 25920  
 acccattcaa acgtccttc actctccctc agctcagcag ggctgctcgt ccagctttga 25980  
 tattaagccc ttggcatatt ccaagttgcc cacagatcct gatttctaga agcttagaaa 26040  
 agtgagagg ttgcccagc aagctggatt attataatta agtagttctc ttttcaaagg 26100  
 ccttgcatTT tcttagcctc tccttctcca caggggaaact ctttcttcgt gatgacaaac 26160  
 tttctcaaaa cagaaggcca agagcagcgg ttgtgtcccg aggtaaggag gggacctgga 26220  
 gtggtgggtc aggtcttaag agttcctggg ggaggtgcaa gtcggaagaa gcagaaatgc 26280  
 ggacctggg gtgtatttga gcccacaaat atctactgag cacctgcctc ttttgtggg 26340  
 tagggctggg ctgagtggag gaaggagag aacacatggc agtgtctctc catggccacc 26400  
 agatctttgt ctgcctgect cggctctctg ttctagcact agggacctgt gcagacggca 26460  
 acctgctct tctatctct ggctctcttc gcagcttatg acaatggtga ttcttggcat 26520  
 ttgcacaaca tttcacattt tacttgaggc tcacaatgac tttaggagaa aaatggcatt 26580  
 ttacaaacca gaacacagat tcagagatgt tgtgacttcc ccagggtca cacagcttgc 26640  
 tgcaggcaga gccagggtta ggacctgagt gtctcactc acagcttctc ctaccatggt 26700  
 tcacagtga ctgtctgcag ggctggtctt ggaaaaccct ggcacacagc tctcttgctc 26760  
 attcatccac ttcttagagt ctacgctct ccatcacgtg catgtgcccc catgactccc 26820  
 tgccaccacc ccagacaatc tttcccaact tggctttgct ctctccaaag agccgcctct 26880  
 ttctggaaga gcttcttccc tatcaaatca tatttgtttt attctgggct aggcacggtg 26940  
 gctcacacct ataactctag cactttggga ggccaaggca ggaggatcac tggagcccag 27000  
 gagttcaaga ccagcctggg caacatagt agacctgtc tctaaaaaca aaacaaaca 27060  
 aaaactgtat ttgtctatt ctttacctca ccactctgct ctctctctt aaaccagctt 27120  
 ttcttetaag tttatctata ccagcgtgtt actgctaggc attacctgta actcccaacc 27180

tcccagcctc ccctctcagg taacgtaact gtgctgtacc agggaccatt tccaacactg	27240
accgttgccag cagcgctgggt gataccaccagc cagatcctcc actgatgagc aggaccagcc	27300
ctgccgggtgg agttttgtct acggggccac tctggcaaca aggtctagct tcctgagtgt	27360
ccaggacctc tgggcaaaag tgctgtcttt gactacagtc tttgccatca tttgtacca	27420
aattcagaac caagataaca ttcccttgggt gtctatcttg ataactctcc gtctaaaagg	27480
agagctttct acatttggtt tgtcctcatc accccttaac tgtccctgct gctcagatga	27540
aatctctgac agccaggcca gccaagccct cccttctgtc tcccttccca gcagtgttcc	27600
gtctccaatc cagtttgggg gcctgtttcc cttttgtccc atcctttgggt catctctaca	27660
ccacactcta cctccatgta cacataaaga cacttttggga cttattgaga cttggaagaa	27720
aaagagttag gatagcccag agattaagag cctgggctca gaagctagac cgctaaaggt	27780
tcaaaactta aacttcttaa ctctgtgacc tcaggcagggt gactcttccac ctctctgtgc	27840
caagttgtgt gcatctgtaa aatggggctg agaacaaaac ctactgttcc ggattattga	27900
aacaaatgag tgaataacctg tgacacgctt aaaacaagge atggtacatt ataagcactc	27960
aatgaatgtc agttgttatg attcaaatga catgttactc ctcagatagg atgtcactca	28020
gagtgtactgg aacccaacag gttccatcga agcacttcca gccagagag cacatgtgac	28080
tctttgggat ttaacttat gttccagggt gggttgggtt ggtctaattg ggcatttgcc	28140
tcctctctgc ccatcaacac ccctctatct cgggtgtgtc cggtaaatac gaccaagtcc	28200
cagttacaca ccaacatgaa tagatatttg ctccatttat ctgtggctgc aaaacaaact	28260
accctgatca tgggtgggctt aacacaagca ttttattaac tctcacagtt cagtgggtga	28320
ctgagttcag ctgggtgggtt cttctgctgg tctcactagg gatctctcaa ggagctgcag	28380
tcagatggca gctgggggtt gcatcccttg gagcctcaat tggcaccag ggatagctag	28440
acatctcttc ctctctggga acccctaaga gttaccctct cttcaaggtc tgtccgcatt	28500
gtctctccag cagggtagtc aaactttcta atagtggctc aggttcctca aaacaaaaca	28560
aaaaaataga agcttccagg ccttcttaag gcttaggcct ggaactggca acagcattcc	28620
ttctgtgtgt tctattgggt ttgcagtcac agcacaggct cagcttcaga aagtagggga	28680
aacactccac ctctcaatgg agaaagtgc caagaatttg tggccatctt taatccatgc	28740
aacaatcaca aaagagaaga ttaaaacaaa actaatgtgc tgttttttaa aaaaaaatt	28800
agaattcctt ttggagagca aaataaaact gatttctctt ctttcctgag attcatctcc	28860
agttgataat gaaaatgtaa ctttccatga gccacaatga atgatcacac attcattcct	28920
tccatgacta tcaagacctc ctatttgcca ggcacttggg ctgcaaagat atgaccttgg	28980
ctctgtattc aaaggggttca taatctaate tggggagtca caaggggagg agaccaacgg	29040
catgccatgt gacaaggcag cctgggaaat aaagagatca tgagagagtt tgaaaaetac	29100
aagaacatt cattgagcat gtactatgtg ctaaaccctt tcttttcata actcacttaa	29160
tcttcacaac aaccaggcaa gacaatttgt tggctgggca tgatggctca cgcctataat	29220
cccagcactt tgggaggtcg aggcaggcag atcatgaggt caggagttca agaccagcct	29280
ggccaagatg gggaaacacc gtctctacta aaaatacaaa aattagccag gcatgggtgg	29340
gcgtgcctgt aatcccagct actcaggagg ctgaggcagg agaattgcct gaactcagga	29400
ggtggagggtt acagtgagcc aagatcacac cattgcactc tagcctggtc gacagagcaa	29460

gactctgtct ccaaaaaaaaaa aaaaaaaaaa aaaaaagaca aatctgttca ccctgctttt 29520  
 acagatgaac agtgagctgg ccaaggtctc tcagctagaa aatggtgcag ctactgctga 29580  
 accagatata cctgggtcca agacctaggt ccttgacctg aatcactgta gaagccacaa 29640  
 attgcacaga tgacaaagag cagaggccat tttcacacct atgtcatgtt ctggcttccc 29700  
 caagactggc catgaagcct gctccctaga aaacaggaag agaaatatgc ataaaccagg 29760  
 gcagctcatt tctctcttcc gtgttctgtc cttcagccct gagaccagc tggctctcatt 29820  
 ttcttggggc aggcgagtgt cctggccaat gtctctcagt tccatgtgtc ctgctgagtc 29880  
 tccgcacctg tttgccctgc tcagtccttt cagccacagg gccacttggg ccaagccacc 29940  
 tgtcccccat ccagccagc atccctgaaa gtgateccca gcttcccaga cctctgcttt 30000  
 tcccattttt atttatttgt attgggatat aattcacata ccataaaccc caccatttta 30060  
 aagtgtacat acgggttcagt ggttttttagt ataatacacag agttgtgcaa ccatcaccac 30120  
 cgtctaattc cagaatattt tcttcttctt ttctttcttt ttttttctt tctttctttc 30180  
 tttttctttc ttttcttctt ccttcttctt tttctttttt taggtggagt tttgctcttg 30240  
 tcaccagggc tggagtgcag tgacgcaatc tcagctcact gcaacctcca cctccagggt 30300  
 tcaagtgatt ctctgcctc agcccccta gtagctggga ttacaggcgc acaccaccac 30360  
 atctgataat ttttgtattt ttagtagaga cgggggtttca ccatgttgag caggctgggtc 30420  
 tcagactcct gacctcagggt gatctgcccg ccttggcctc ccaaagtgtt gggattacag 30480  
 gcgtgataag ccacggcgcc cagccccag aacattttca tcacctacaa aggaaacccc 30540  
 aaatccagta gcagtcactc ccattctctc ccttccccctg tccctggcca cagtctactt 30600  
 tctgtctcta tagatgccta ttctggacat ttcctataaa tagaattgta tatggtgtgg 30660  
 ccttttgtgt ctgtcttctt tcaactcagca tcatgttctc caggctccatc catgttgtag 30720  
 cctgtgtcat tgcttcatcc ttcttatggc taaataagat tctgtgtatg aatgtaccac 30780  
 attttatttg tccattcatc cgtcagtggc cacttgcatt gtttccactt ttttggcgat 30840  
 tctgagtagt gctgtataaa gcattcgtgt gcacattctg gtggatatcg aatcacttct 30900  
 ccacatctta gtaacacacg tcacttactc cccactctgt catccttcta tctgcagtat 30960  
 cccaccgca ggacgtctg ttctctgac cgaggttgta aaaagggatg gatggaccgg 31020  
 cagagcaaag gtaccttctg tttcttttcc cgagacccta ggggtggatg gtctggcatc 31080  
 ttggtgacat ttgtgatgcc caggtcagggt cttcagcctc tgctctcagc tgccctcttc 31140  
 caccatcacc aagccatagg cgagtctgcc catgcttcgg ctctgtcccc agcagaccag 31200  
 ctgctgactg taaacatgac tccagttttc cagttagaga agaagctcct aaaaacctag 31260  
 caggttcagg attctaactg gtagaaaatt cacatggcct atagcatcat ctgagtattc 31320  
 taaactttcc ccctgaattt cctcaaagggt tgaggacat gaacttttac ccccaggga 31380  
 cctggcagca ataccatata taacctgcag aatttttttt gttttttatt ttattttatt 31440  
 ttttaaacat tttttgact gttttatttt gattttgatt ttgattttat ttatatctaa 31500  
 gtgcagtgtt attgcgatac ctgcagaatt tctttatctc acattttaac ttaaaaaggc 31560  
 acagggcagc gagcgcagag gctgggtgcct gtaatcccag cactttggga gggtagggca 31620  
 gatggatgct tgaggtcagg ggttcgagaa cagcctggaa aacatgggtga aaccccgctc 31680  
 ctactaaaaa taaaaaatc agccagacat ggtggcacac gcttataatc ccagctactt 31740

gggaggctga gacgtgagaa tcacttgaac ctggaaggca gaggttgag tgagccaaga 31800  
 tcatgccact gcactccagc atgggtgaca gagcgagacc cctttaaaaa aaaaaaaaaa 31860  
 aggcacaggg caattttaaa aatactgcaa atagtaaaaa aaaaaaaatc agtgggtata 31920  
 atgcaaacac acacaaaaag gcatatgcc attactgcat tctactccat actgtatgtg 31980  
 tatttgagtt agtataaaaag ttattttaac attgctcact atttaattaa ttctcccttg 32040  
 gaaactgatt aatcatcctg gcactccagg aagatgtgcc atgctgattt catggctttg 32100  
 cacatcctgg gcaggctgtg tacccttgga gggacttggt cccctttgag aggccatggt 32160  
 ctagtccatt tatactaagt gagagcatac acctgttccg ctcccctcat gggcaccttt 32220  
 tcttataaag aaacaaaaga gccagcagaa tccacagtct ttctgtgttc tctctgatct 32280  
 ttattatggt ttgcttggtt gccttgccct gtgttcgttg tggtaggat gggcttgatg 32340  
 gaagctgaag ctgcgtgggt tggaaagcct ggtcaaagcc tagtctctcg cccgggttga 32400  
 gttaatgatg tccctcctgg agaacgtcct ctctgcagtt ctttcacatc tgtggttcta 32460  
 cgatgctttg acccctatag gaattcagac cggaagggtg gtagtgcag aagggaacca 32520  
 gaagacctgt gaagtctctg cctgggtgcc catcgaggca gtggaagagg cccccgggtg 32580  
 agtcgcatgg ggagacagac acagtggccc tcagcggcga ccagatgagg ccttgccgag 32640  
 gctgcttggg ccttcccctc tcagcacagc cctgcaaagt cctgggtcct accggcttg 32700  
 ggaccctgc gctctggatg cactgcttgg cacaaactag tatctctggg agggccatgg 32760  
 tggttggtaa actgttgtaa caatcctgta ccaactggta aatagctact accctgagca 32820  
 tccttgggtg tccctggccc ctctcttccc ccagatcttc cagggtaccc ccagacccc 32880  
 tcctgtagtg ccacagcagg atcccttctg acttgtcagt gtccatactg agtgatcaag 32940  
 gatagaaagg aaggaggagg atggaaggga aggacgaagc gaggaagag aagggaagg 33000  
 ggaggaaaaa gcaaaagggg tgagggtaaa agggggggga aagggaagtt tctcagatta 33060  
 aatgcttaca atgacataca gatttggtgg tcccttgat tgatgcttcg cttcaatata 33120  
 caaagtcaca atgttaaact tcagaagcca caagggtga tgtatttcag cagagaatag 33180  
 ttagaaagac ctggattcaa ttcttaactc taacaccatt ttgctgtgtg tccttgggaa 33240  
 aatggcttaa cctctctgag ttctagtgc ctcacctgta aaagcagaat aataatttca 33300  
 ccaacttcat agggctgttg taaggattaa atgagatgat acttgtagag ttattgtaag 33360  
 gtaagcccca tgcattgcctg gcttacacac acacacacac acacacacac acacgcacac 33420  
 acgcacacac acacacacac acacaatcta ccctagaag tgtggtggtt ctagaccagc 33480  
 actgtccaat tgaacttgat gcagtgtgg aaatttctgt atctgtgctg tccaataggg 33540  
 cagctactag gtacatgtgg ctattgagta catgaaatgc gactactgaa tttttgaaag 33600  
 agatgataga ttagatagag aagatagat agatagata atagataata gatagataga 33660  
 caggtagata gatagataga tagatagata gatagataga tagatagata gatagagttt 33720  
 tgctatgttg cccaggctgg ttttgaactc ctgggtcaa gcgacccctc tgccttggcc 33780  
 tcccaaagtg ctgggttac aggtttgagc cattgtccc agcctgaatt tttaattaaa 33840  
 tttaaattta aatagccaca catgtctagt ggctaccata ttggacagcg cagttctaga 33900  
 ccgatgtgat tcaggatcat tccctcagca tcgtggggca aagagaaaac tgccccaagc 33960  
 tggcctgtag aaggctcagg cgaaggtttc ccaatgccgg gatggggggg gcgctcagca 34020



gcatcacccc ttatgattct caatcgctaa tagctccact caggttcatt tctcggtcag 34080  
 gggcatttct ttgggaatca cccagctctg ggagatacag cagcctccac tcaggtagtc 34140  
 cttgttcaag acaagcggcc cttgactgac tgcagtttca gttccagctc tgctatcaac 34200  
 tcactcatta aataaactgc atctccagtg tgcctgcctc tgggctggat tttgacgtga 34260  
 cctgggcaag caactccctg aacttcagtt tctcatatat tatatgaatt agctaagatg 34320  
 gttcgtttaa tcattcattc aacacatcca tcaccacgta gtaggtgtta gatatttatt 34380  
 tcatacgtaa ctacgcataa gagactttgc taagtttttag gtaaaataca agtcccagat 34440  
 acggagcaag tctcaaccac tgtacatacc tgaatgtgta attacatcac tgtaagaggt 34500  
 gccacagtaa atgccactgg gtcttgtgtt agtccattct cacacaaaga actacctagc 34560  
 caggtgccgt ggctcacgcc tgtaatccca acactttggc aggctgaggc aggcggatca 34620  
 cttgagggtca ggagtttgat accagcctgg caaacatggt gaaaccccat ctctattaaa 34680  
 aaatgcataa attagccagg tgtggtggca cacgcctgta atccccgcta ctccggaggc 34740  
 tgaggcagga gagtcgcttg aacccgggag gtggagggtg cagtgaccgc agatcgcgcc 34800  
 actgcactcc agcctgggtg acagagttag actccatctc agaaaaaaat aaaaataaaa 34860  
 ataaagaact acctgagacc aaatacttta cgaaaaaaaa gaggtttaat tgactcacag 34920  
 ctccacaggc ttaacaggaa gcctcaggag acttacaatc atggcagaag gcgaagggga 34980  
 agcaaacaca tcttaccatg atggagcagg aggcgagttt cgggggatgt gccgcacact 35040  
 tttaaatgat cagatctcgt gagaactcac tcactatcac gagaacagca aggaggaagt 35100  
 ccgcccccat gattcagtca cctcccacca ggccctcct ctggcacatg gggattacaa 35160  
 ttcaagatga gatttgggtg gggacacaga gccaaacat atcagatctc aagaaggag 35220  
 aaattcttct tggaggagct ggaggggctt tgtggagagt ttcagaatgc tttgccact 35280  
 aggtttgctg tatccatttc tcttcatgta tcccaaagac caagccaaga aaccagaagc 35340  
 ctctggtccc actggcccat gggctccctc ggtccccacc gtcactaatg gccattttgc 35400  
 atgtctctct cccaggcctg ctctcttgaa cagtgccgaa aacttcaact tgctcatcaa 35460  
 gaacaatata gacttccccg gccacaacta caccacgtaa gtgcccaggc tgcttggtg 35520  
 tcttagttat ctactgctga gtaataaatt atcccaaacc tcagaagcct gaaacaacaa 35580  
 acgcctattg tctcccacgg tttctgtggg tcaggaatct gggaatgact ttgctgctg 35640  
 gttctggctc aaggtctgtc aggttgtagc caagctgtca accagggctg cagtcatttc 35700  
 taggcttgac tggggctgga gaacactttt ccaagctctc acacagttgc tcgtgggaga 35760  
 gctcagttcc tcaccacgtg aacctcgccc tagaccactt gagtatcctt ggtatatggt 35820  
 ggctggcttc tcccagagca agtgacccea gagagacaga gcaagcaacc aagagtataa 35880  
 ccaagatgga agccacagtc tttgggggga gacccaaca cttctgcat atgccattgg 35940  
 tcacacagat caacctggt ccagtgtgag aggccactgc ccaggggtcc caggaggcag 36000  
 tgatcatttg gggctttcat ggaacctctc caccacactg gctactcct gggaaagaga 36060  
 cagatctgtt ttcaatcgag atgtttgttt gtttgccttt aattatgcac aggagaaaca 36120  
 tcttgccagg tttaaacatc acttgtacct tccacaagac tcagaatcca cagtgtccca 36180  
 ttttccgact aggagacatc ttccgagaaa caggcgataa ttttccagat gtggcaattc 36240  
 aggttggtgg tgctttgtac actgggatgt ggggctgtgt gtctagggat ggaggatgtc 36300

aacagccaa gaggccgggc cactgggtct tcataatgtg gctcacattt actgagcatt 36360  
 tagtaaattcc acccgctgcg ctaaggtctt tacctacgct acctcgtaa atcccaaac 36420  
 aatccttatg agtgagagct acttgggtga ttcccttccct gtggctgctg tagcaagtta 36480  
 tcaaagctta gtggcttcaa acaacacata tttgcttatg ttgccagaga tcagaagttg 36540  
 gagatgattt tccctgagcc agggcggtgc tccctccggg actttaaggg agaattccagt 36600  
 tcctcagctt ttccaccttc tggagctgca ttccttgcat ttcttcaaag ccagcagcat 36660  
 aacatcttgc ctcagtggcc actttcactc cctatcctgt gtccaatctc cctttgcctc 36720  
 tgtcttaciaa agagagagag catttacaag agggggcatt taaggaccaa ctggataatc 36780  
 caggataatc tcccatctca agatccttca tttaggctgg gcacgggtggc tcatgcctgt 36840  
 aatcccagca ctttggggagg ctgaggtggg tggatcacct gaggtcagga gttcaacacc 36900  
 agcctggcca acatggtgaa agcccatctt tactaaaaat aaaaaaaaaa aaaaaaatag 36960  
 cggggcatga ttgcaggctc ctgtaatccc agctactcgg gaggtgaga caggagaatc 37020  
 gcttgaacct gggaggcaga ggttgcagtg agccgagatc gcaccactgc actccagcct 37080  
 aggtgacaag agcgaaactc catctcaaaa aaaaaaaaaa aatccttcat gtattcgcat 37140  
 ctgcaaaagag ctttccctag gggagtacta ggaggtaaag cagaaaagat atttgataga 37200  
 gtgccctgaa ttccagtcta ataagtttgg acttgatctt taatgggggc gtggggggca 37260  
 ttaaagggtg ttgggtacag gagtggctctg ttgaaagttg tatttttagga caatgagttt 37320  
 aacagtgatg tgtcccagac gggggtaggg agagttagga gatgcgattg tggctgccac 37380  
 aataacactt gtgcgagtta ggtggggctg tacatatggt tcttcaatca gcatttttcc 37440  
 tctaaaaacc ttaagcaatc ctggctatgc agggagatgt ctggcggttg cgtaactcac 37500  
 acccagcagc catagagact gtcccttgtt gatccttcag ggcggaataa tgggcattga 37560  
 gatctactgg gactgcaacc tagaccgttg gttccatcac tgccgtccca aatacagttt 37620  
 ccgtgcctt gacgacaaga ccaccaacgt gtccttgtac cctggctaca acttcaggtt 37680  
 actccaaggc ccagggtcaaa ctcacccagt ggctgaatcg cattcccagg aactgggtgag 37740  
 actaattttg gtttccaagg caacaagatg aatgaaaaaa gactttctct aagaactagg 37800  
 tgataactga atttttttcca taatttttta aaattctcaa aagagatgca cactctttat 37860  
 ttttttactt attttttttt ttttgaaatg gagtctcact ctgtcaccca ggctgaagtg 37920  
 cagtggcgcc atctcagtca ctgcaaactt ccgcctccca gggtcaagcg actctcctgc 37980  
 ctcagcctcc caagtagctg agattacagg cggatgcaca ctgtttataa aacaaaacta 38040  
 ttgggaaaca gaaaagcata gagggggatc aaaatcaccc ataattcccc taccctgaaa 38100  
 taatcaataa caaccctcgg gggaatttct ctcactgtga ccaattattt catacagctc 38160  
 ctatgagata atagcatata tatatatatc ttgtggtatt ctgcagggtt ttccatacca 38220  
 cagccactca aaattctttg taaccatcac attaatgatc ataacattcc atttttagg 38280  
 tgaacaaata acaactgcta caattcagga agtgttttct tttcttttct tttcttttct 38340  
 ttttttttt tagatggagt cacactctgc ttgccaggc tggagtgcag tggcatgatc 38400  
 tcagctcact gcaacctctg cctcctaggt ccaagcgatc ctcccacctc ccaagtttct 38460  
 gggaccacag gcatgtgcca ccaccccag ctaatttttg tatattcagt agagatgggg 38520  
 tttcactgtg ttggccagtc tggctcgaac ctcttgacct caagtgatct tcccaccttg 38580

gcttcccaaa gtgctaggat tacaggcatg agccactgtg cctggcccaa ggagggtttt 38640  
ccatatacca agcactcccc atcgccatcc ctaaactctcc caacaaccct ggaaggaaga 38700  
tattgtttct ggaagatgat ttgcccaaga cccacagctg atagtacatg ttgcataatt 38760  
ctaaccacag tcaactctgac cccacactca cactccatcc ctteettccc atctcatgat 38820  
tttctcacct acgcctccat gattgaatat ttgagttgct tccagttttt ctattacaag 38880  
taaccacagt gtgcatcttt gcacataaac ttctctttga attccagggt acttcttag 38940  
gataaatttc tagacttatt gaatcaaagg ttgtgaacat tttatcatat gctttttatt 39000  
tttaaaaata tctatagtta taatgtttca tttttttttc tgagacagag tctcactctg 39060  
tcacccaggt tggagtggac cgggtgcaat tatagctcac tgcaacctct gcctcccagg 39120  
cccaagtgat cctcccacct cagcctcctg agtagctagg actacagggt cacgccacca 39180  
tgcccagcta attttttaaaa tttttttgta gagtcgggggt ctactatat tgcccagggt 39240  
ggtctcaaac tcctggctca agcaatccgc ctgccttggc ctctaaagc gttgggattg 39300  
cagggtgtaag ccactgcacc tggcctataa ttttcatttc taggattttt atttggtgct 39360  
ttttcaagt catctattat tgttcagtg agtccattc ttaccttaag gatcctactc 39420  
cttctgtcca ttctactgta tcattccttt cataccgacc tattatctga agtaacttgg 39480  
gtgggagttc tcctcgtggg ctttgaaata ctgtcttcag tagaaaagat cttatgcaaa 39540  
gttctgcatg tgctgtgtgt tatggagtat tccttatata ttactcaaag cagctccata 39600  
aactggcagg cagccctta aatgtgtttc tgtttctgtt tttttgtttt ttgggtctgca 39660  
tgatgtttta aaacttgaga caggccaggt gcggtggctt atgcctgtaa tcccagcact 39720  
ttgggaggcc aagggtgggag gatcacttga acccggggggt ttgaggctag cctgggcaac 39780  
atagggagac cccatctcta caaataataa tttttaaaac atcagccagg tgtgggggca 39840  
tgacacctctg gtcccagcta ctcaggaggc tgaggcagga ggattgcttg aacctgggag 39900  
gtggaggctg cagtgaaccg tggttgtgcc actgcactcc agcctgggtg acagagttag 39960  
acccttacat ggtggccact ggctggagct gagtatcagt ggtcctattt agaaaggggc 40020  
tgggctttct ggttcatcac tgtccccgcc actccttagt gcttatacct ggcccacatc 40080  
actcatttct gtcacttgcc tggcccctgt gtagacattt gagtttgaaa cccttgactc 40140  
aaaggcaggc tgatgctttt tgcctcctct ggatcaatgg aatttcatcc aaggcaatgg 40200  
aaccagctcc taattgtgat aactttttga tcatccctg ccactgtgct gaataggctt 40260  
atggccatta agaagagagg acaggatgaa tgggtccctg actgactgcc cttccagggt 40320  
gttttttttag ctgtcagcag aagcatgctg ggcagtgtac caatcagggt tcagcaagtg 40380  
tccttccagc gactgagttg agcacaataa catctgctcc tggagagacc tgagccctct 40440  
gaaggcctca gccagttagt atgttaatgt tcttttagaa caaagtggta gagcatttgt 40500  
tgtctggaat gagcaccaga agaaatctct ggccaaaaat aaattgttga gggccgggag 40560  
cagtggctta tacctgtaat cccaacactt tgggaggccg aagcaggagg atcacctgag 40620  
gtcaggagtt cgagaccagc ctggccaaca tggtgaaacc ctgtctctcc taaaaatata 40680  
aaaattagct ggacgtggtg acagggtgcct gtaatacagc tactgggagg ctgaggcagg 40740  
gagaactgct tgaaccgga aggtggaggt tgcagtgagc cggggctcatg ccacttcact 40800  
ccagcctggg caacagagca aagctctgtc tttaaaaaca acttaaataa ataaataaat 40860

tgttgaggtc tgatgagtaa gtggacaagt tttttccag cagacacaca aaagagaagg 40920  
 aaattacagg ttatacgagg tttttcagaa aatataactt tctaaaacat aggaagtga 40980  
 agaagttgat cacattacag aattctgttg tttagaaaat gacctgtggc gaaatgtcct 41040  
 tattcagtga ataggtgatt ccgcttatgc acgacctgtg tgaagtggat caggccaccc 41100  
 agaatgcacg acgcgcttct caggcccagc aggagtatgt gtctgtgtta atttctgtg 41160  
 gctattatga ctaattgcca caaatgtggt ggcttaaaac aacagaaatt aatcttctta 41220  
 tagttctgga agccagaagt ttggaatcaa gatgtcagca gggccacact cgctctgatg 41280  
 ctctacggga gggctctctc ttgcctcttc cagcgtctgg tggctccagg cattccataa 41340  
 ctttatagca gcgtcccaca aatctctgcc tccatcctca catggccttc tccactgtgt 41400  
 ctctatgtct tcaatctctt tttttttttt tttttttttt gaggcagggt ttcactccag 41460  
 tctcctagac tgaagtgcaa tggcgtaatt tcgggtcact gcaacctctg cctcccgggc 41520  
 tcaagcgatt tgatctctcc tttatcttat aaagatacta gtcattggat ttggggctta 41580  
 ccctaaatcc aggataatct catcttgaga tgtttaactt aattatatct gcaaactg 41640  
 tttttccaaa taaggtcata tcacagccac tagggattag atacttgaac atatcttatt 41700  
 tgggggctca acccattcca gtgtacgaaa aacactcttg ttcaaggccc gatgtttctc 41760  
 agggcatagc cactgacta cctgcatcag aataatcact tggtagctgt actggaaata 41820  
 cagactccta gaaacatctc agagcttctg caaccactct ttgagtgagg ggctcaggag 41880  
 tctgcctctg aacacactca cccaagtga ttctttcttt ctttcttttt tttttttttt 41940  
 tttgagatgg agtcttgctc tgtccccag gctggagtgc agtggcgtga tctcggctca 42000  
 ctgcaagctc cgctccccgg gttcacacga ttctcttgcc tcagcctccc gagtagctgg 42060  
 gactacaggt gcccgccacc acacccggct aattttttgc atttttagta cagacggggt 42120  
 ttcaccatgt tagccaggat ggtctcgatc tcctgacctc gtgatctgcc tgcctcggcc 42180  
 tcccaaagtg ctgggattac aggcattgagc caccgcgccc agccatcacc ccaagtgatt 42240  
 cttgccttca gtttaagagc cactgtaaca agactatgga agcagaaatt cagtgctta 42300  
 ctacacaatg ttaaccttcc caggcaaacc aactcacata gggagataat gccaatccca 42360  
 gggcaggcag tggcaatgca tgcttgcttg cgaattaaaa aacaaatcac tgcctgggca 42420  
 cgttggctca tgcctataat cccaacaatt tgggaggctg aggtgggtgc atcacttgag 42480  
 ctcaggagtt caagccagcc tgggcaacac agtgagacca tcctctctat aaaaaaattt 42540  
 taaaaattag tcgggtgtaa tggtttgac ctgtagtccc agctgccggg aggtgaagt 42600  
 gggaggatga cttgagcccg ggaggcggag gccacagtga gctgtgttca agccactgca 42660  
 ctccagcctg gatgacagag ccagatcctg tctcaaaaaa aacaaaaaca aacaacaaca 42720  
 acaaagataa atcactcaat acatcagcaa gagaaaaagc tctcttgaaa tagtcacatg 42780  
 caaagaaatt gaattccctc cagtcagaaa gagccactaa agtgcctgag aatatctgat 42840  
 cgatttcaat gtcaggtttt gagaggtttt ttaaaaacag tttcagatgt ttcttactat 42900  
 tttttggcag aacatctgcc atctgcttcc ttctctcctt acatcttgta actagacggg 42960  
 gaattaataa ctcagaaaaa ataaacatgt atatgtaact ttatctaaaa agaatacatca 43020  
 aagtgtgggtg agaagaaggg gcagatttaa agtttttatg aagcgttcat ttaagcctc 43080  
 cttaattatt cttgaaaaac aaaacacacc acttttctct actgcagcac tggtaggggt 43140

tgcatatcac gggtgactgt gatgatttgt gcctggcgct taatttttaa agttagtact 43200  
 gagtgatgac agagagaggt caatgccact caaagaatat tttgtttgtt tgtttgtttg 43260  
 tttgtttgtt tgagaggcag tcttactctc tcactcaggc cggagtgcag tgggtgtgata 43320  
 tcggctcact gcaacctctg cctcctgggt tcaagtgatt ctccctgcctc agcttcctga 43380  
 gtagctggga ttacaggcgt ctgccaccac atccagctaa tttttgtatt tttagtagag 43440  
 atggggtttc atcatgttg ccaggctggt ctcaaactcc tgacctgagg tgatctgcct 43500  
 accttggcct cccaaagtgc tgggattaca ggcgtgagcc attgcacctg gccaagaaga 43560  
 atttattaca tacttttccc aagagatggg gccacaccac accataccag gccacaccag 43620  
 gccatgccat gctagacagg gccacaggag gaagtaccag atccgctcaa gaggcagaat 43680  
 aaagggtaaa gaatggtcca gagctttatt gtgttactca gtggaagggc aaggcaggac 43740  
 tagggaaaca gcttaggggt aactactttg aataatgcca gtaggttctg agttacagga 43800  
 atggtctcta aatgtctggc cctcacccta cctgtcccta aggagaaata ctggagagtt 43860  
 agaaaaggag gtggttgagg gattggttg agggtttgta atatgatttt cacaccctca 43920  
 caaaagctgg attgcagagg agatgtaaac aacttcagcc ttgggaggcc aagatgaaag 43980  
 gatggcttga agccaggagt tcaaggctgc aacaagctat gattacacca ctgcactcca 44040  
 gcctggttga aggaatgaga tcctgattct acaaaacatt tttgaaaaaa acttttttat 44100  
 tttttctttt cctttctttt tttttttttt tttttgagac agtcttgctc tgtcaccag 44160  
 gctagagcgc agtgatacga tctcggctca ctgcgacctc catctcctgg gttcaagtga 44220  
 ttctcgtgcc tcagcgtcct gagtagctgg gattacaggt gcctgccacc atgcacggct 44280  
 aatttttgta ttttttagtag agatgggggt tcaccatgtt gtccgggctg gtcttgaact 44340  
 cctgacctca agtgatccac ccgcctcggc ctcccaaagt gctgggatta cagggtgtgag 44400  
 ccacggtgcc cgacaaaaaa atttaaaaat aaaaattagc cacctgtagt ggtgcacgct 44460  
 tgtagtccca ccagctact tgggaggctg aggtgggagt atcacttggg ccaggagggt 44520  
 ggaggctgca gtgagctctg atcatgccag tgctctccaa cctgggtgac agaacaagac 44580  
 cccacctcaa aacaaaaaaa aacaaaaaac acaacttcag ccttttagttt ggccctgtga 44640  
 ttaatgattg ccaaataggc atacacagaa tctaagaaaa tacagtttgc tgagggtgtc 44700  
 ctgtctccag tccagtaatt agtatgcaag atttaccacc gccactccc actttgttct 44760  
 agatgcaaaa cctcttcttc ccccttaagg aatagtcata ttgcttgaag tttttttttt 44820  
 ttaatttctc tgcttgttgt taatcctgtg ttggtttaaa atgtgcattt taatcttaag 44880  
 cgacaagctg attttccgtc actctgagat gatgcaggta cagggtgtga catggggagg 44940  
 ggggactgct ctttgggtcc aggtgggttg agagagaccc agggcttttg attgtgttcc 45000  
 ttcccctgcc acctgtcaca gagccagggg acagaccagg aggactagac aggccactgt 45060  
 tttggctttt ccttcataa aataccagca ttttttgcca gatgcagtgg ctcatgcctg 45120  
 taaccccaaa actttgggag cctgaggcag gcaaatcact tgagctcagg agttcgagac 45180  
 cagcctgggc aacatggcaa aagcccgctc ctacaaaaaa tacaaaaaat tagccgagca 45240  
 cgggtgtgca tgctgtagt ccagctacc tgggagactg aggtgggaga atcatctgag 45300  
 cctagcaagt caaggctgca gtgagtggta atcctgctac tgcactccag cctgggtgac 45360  
 agcgtgagac cctgtctcaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa ccccaaac 45420

cagcactttc aaagggatct tacaaatata gatccttttc ttectacaga tacgccaaagt 45480  
 actacaagga aaacaatggt gagaaacgga ctctgataaa agtcttcggg atccgttttg 45540  
 acatcctggg ttttggcacc gtaagtctcg tttcccagct cggggcaccg gcatectatg 45600  
 actgtgtcct aattactgct gtggggcctc catggaggga agggttttgg tctcagcctt 45660  
 cagctagcac tgggcatttc gcacatggga taaaagagga agaagatggt ctcggtctgc 45720  
 tgccctgaa tgagtcattc gacgagtaca ggagtgggccc tggaaagcaa ttctaacatt 45780  
 cggcttttaa aatactctga tacttaacaa gagagagaaa gaaatctctc tgtaaaatag 45840  
 tttgtaagta acagaatcca attcaaattg gcttaagggg aaaaaaaaaa gttttgagtg 45900  
 ggcttttttg ggttcacgta actgaaaaaa tccagggggt atttttaagc tacaagcatg 45960  
 gttggatcca ggtgtcaaaa caacaacatc atcatcaaga acttgccacc ttccgtgtct 46020  
 tatatacgtc ctccctgggtc ttggcagcca tctctctatg aaggatcag gcagcacctt 46080  
 cgatctgacc agtttagcaa cctcagtga aagaaaacac ctctttgcca gaagggttcag 46140  
 caaagggtctc agacaaatgt cattggctct gattggccca ctatggatga catgccact 46200  
 gctgaactaa tcaactgtgac cagggtcatt ccatactccc agtggttagt gtcgaaccag 46260  
 ccctaaccta gcccgagggc cagcagagag gccaacaccc cagccccctc aacacacaca 46320  
 cacacacaca cacacacaca cacacacaca cactacatgg actgaaaaga aaattggagt 46380  
 gttgccttct gaagatggaa ggaaaagatg ccagctatgc aaaaatcaac atatcccagc 46440  
 acatcctgtc caagagcatg ttatctatgt aacacagtga taagcaggac tagaagcaat 46500  
 aagatatagc tgagagaatg caaagaccgc tgggaataaa aaccacaagg ctgggccggg 46560  
 cagggtggct catgcctgta atcccagctc tttgggaggc tgaggcagggt ggatcacaag 46620  
 gtcaggagct caagaccagc ctggccaaca tgggtgaaacc ctgtctctac taaaaatgca 46680  
 aaaaattagc tgggcgtggc ggtgggtact tgtaatcca gctacttggg aggctgaggc 46740  
 aggagtatct cttgaaccca ggaggcggag cttgcagtga gccgagattg cgccactgca 46800  
 ctccagcctg ggcaaaagtg cgagactctg tctcaaaaaa aaaaaaaaaa aaaaaccac 46860  
 gaggccaccc agacattcct tccctctgct tctgaagtaa tgtcaaacct tttggaggag 46920  
 aagcttgtct gaaacctca cttgtatgg agaaatgata gcccttcagt gggctctccc 46980  
 ttaatgtgtg ggccccagt cttgctcatg taaaaccttt atgggtccaa gcacaccctg 47040  
 cagggtgaa gcaggatgcc tgagagtcag tttcagctc cgtagtctct gcctcagcca 47100  
 gcactgaac gcatttatcc aagtcacagc atgaggctcc gctccctgat agaaccaaca 47160  
 attgcacgtt gaagcaaaag agcgttgctc tgaatttcac ctgagtaaac tctccactc 47220  
 tgtttttagg gaggaaaatt tgacattatc cagctgggtg tgtacatcgg ctcaaccctc 47280  
 tctacttctg gtctggtgag agattctctt ttccatgctt taggaaaatg gtttgagaaa 47340  
 ggaagtgact aacgcagcgc ttgtctgcat tctcccagg ccgctgtgtt catcgacttc 47400  
 ctcatcgaca ctactccag taactgctgt cgctcccata ttatccctg gtgcaagtgc 47460  
 tgtcagccct gtgtgggtcaa cgaatactac tacaggaaga agtgcgagtc cattgtggag 47520  
 ccaaagccgg tgaggccgct gtgttcacag gacaccaaga catggagaga ttccatgaaa 47580  
 tcaactcagaa atgcacgaaa attaggccca aatcacaggc ttcattcctgt agtggatacg 47640  
 tcgctggggt ctaccccgat caaccaactc tcagataaat tttttggtct tagagaagaa 47700

tggaacaaa	aatggagggg	caagatagag	ggaaggcaaa	tttttatgtc	taggacttgc	47760
caattttgtc	atttatttat	ttattttatt	atttatttat	ttattttatt	atttattttc	47820
agatggagtc	ttgctctatc	accagggtg	gagtgacag	gcatgatctc	ggctcactgc	47880
agcctctgcc	tcccaggttt	aagtgattct	cctgcctcaa	cctcccaagt	agctgggact	47940
acagggggcac	accaccacac	ctggctaatt	tttgatattt	tagtagagat	ggtgttttagc	48000
tatgttggcc	aggctggtct	caaactcctg	acttcaaagt	atccaccac	tcggcctccc	48060
acagtgtctg	gattataagt	gtgagccact	gagcccagcc	tgttttgtca	tttattaaat	48120
tggatatagc	aaaaaagaaa	aaagaaaaga	aaaataacaa	ctttggagaa	caatttggca	48180
gtgactaatg	tttaaatggg	acatacttta	cagcctggca	ttttcagttc	tccatacctg	48240
ccctagagaa	acactcacat	gagtacccgc	agtacatgag	tacaagatgt	tcaaagcagg	48300
attgtttatt	aaattatcaa	taatatatga	ttattaacag	tgaagaaata	gcatctaacc	48360
aaatatccaa	caggtgaatg	gtgaaactat	ggtaaatcca	tagaaaggaa	taccaggcac	48420
cagcttaaaa	aatgagata	gataataata	atggcaaaca	tttacacagc	acttctaaat	48480
gctttatcta	ttaactcaat	cttcacaaca	acctgatgta	gacagatgct	actattatct	48540
ctttctatat	atgaggaaat	tgagccacag	aaagggttaa	taattggccc	aaggctgggc	48600
acagtggctc	acgcctataa	tccaacact	ttgggaggcc	gaggcaggca	gatcacttga	48660
ggtcaagagt	tcgagaccag	cctggccaac	atggtgaaac	cccactctta	ctaaaaatac	48720
aaaaattagc	caggcgtggt	ggtgcatgcc	tgaagtccca	gctacttggg	aggctgaggg	48780
aggagaattg	cttgaacctg	gtagggcagag	gttgacagtga	gctgagatca	tgccattgca	48840
ctccagcctg	agtgacagag	ggaaactctg	tctcaaaaaa	aaaaaaaaaa	aaaaaattat	48900
aggacttgac	aagagcagat	ctccaagaca	ttttgttaa	agaaaaaag	caaactgtag	48960
aacaatatat	gttatatgat	atataggtta	aaaaaaaatc	actagacaca	gaagctaatac	49020
tgtatatatt	ctgcatgtat	attctatata	tacacaagtg	tacacacaca	cacacataca	49080
tatacatatg	tgtgggtgta	tatatgtaca	caaaattgta	ccagtggctg	cttctggaaa	49140
ggagcttagg	gtgggggagt	aggagtagtc	aaagagattt	tagcgtcatc	tgtattgttt	49200
tgatttgatt	aattcagact	ttatcaagca	ggtcctctgc	gttcaactcc	atgattttcc	49260
ccaaagataa	atctctggta	cctaaaaaca	aagacgattg	gctagacgtg	gtggctcacg	49320
cctgtagtcc	cagtactttg	ggaggccaag	gagggtggat	cacctgaggt	caggagtcca	49380
aaaccagcct	ggccaacatg	gagaaacccc	gtctctacta	aaaatacaaa	aaattagcca	49440
ggcatgggtg	cgcagcctg	taatcccagc	ttctctagag	gctgaggcag	gagaatcgct	49500
tgaaccagg	aggtggaggt	tgtggtgagc	caggattgca	ccattgcact	ccagcctggg	49560
caacaagagc	gaaactccgt	ctcaaaaaaa	aaaaaaaaaac	aaagacgatt	tctttgtctt	49620
tccctcatcc	aagaacatga	ttgtcctgtt	ccagcagctg	atgcacaatt	cactgtccat	49680
tgtatatgca	ttcacaattt	gaaataaaaag	ttcatctttg	cagctaaaac	taatcaccac	49740
ttcatggccc	aagatgagat	gaaatttaac	aaacatgtaa	ataatttaag	ttgcaatagt	49800
acaaatttct	ggagatactg	aatctagagt	tactgaaatt	gacagaatac	aacaaagaaa	49860
ttttatgcag	caactggggg	gtccaatgta	aaaacattaa	gcagtaagct	gtggctgtgt	49920
tgaatttaca	agttaagatg	catgggggtc	cgcctggcgt	ggtggctcac	tcctgtaatac	49980

ccagcacttt ggtaggccaa ggcgggcgga tcaactgagg tcaggagttc aagaccagcc 50040  
 tgaccaacat ggagaaaccc cgtctctact aaaaatacaa aattagctgg gcatgatggt 50100  
 gcatgcctgt aatcccagct actcaggagg ctgaggcagg agaatcattt gaacctggga 50160  
 ggcgagggtt gcagttagcc aagatcatgc cattgcactc cagcctggca acaagagtga 50220  
 actccaactc aaaaaaaaaa aaaaaaaaaa agcatggggt tccatttctg atttatcttt 50280  
 agactcagaa atcattaatt cttgggtaat gagagttttg agccagcttg ttcaatagtc 50340  
 tatcatttgg caaataggaa ttacagtgc ctttagatag gcaattcttg ataattctgt 50400  
 acaaaaatgg gttaaactttc aaaccatctt ttcctagaca ttaaagtatg tgcctttgt 50460  
 ggatgaatcc cacattagga tgggtaacca gcagctacta gggagaagtc tgcaagatgt 50520  
 caagggccaa gaagtccag taagttaaat cttttgtct tttttttttt ttttaaaaaa 50580  
 tttactgtta aatataaaca catctagaaa cttgtacaaa tcaaaactga tggattttta 50640  
 caaagtaaac atactcatat aaccggcact cagattaaac aattgaaat tactagcagg 50700  
 agtccctttt atgccccct ccaateacta ccctctcttt ctcctttttt aatttttaaa 50760  
 atttggctgg gcacggtggc tcacacctgc aatcccagca ctttgggagg ccaagggtgg 50820  
 tggatcactt gaggtcagga gttcgagacc aacctagcca acatggtgaa acctcgtctc 50880  
 tactaaaaat acaaaaatta gccgggtggg gtggtggcac acatgtaatc ccagctactc 50940  
 agaagactga ggcaggagaa ctgcttgaac ccaggagggtg gaggttgag tgagccgaga 51000  
 ttgtgccatt gcaactccagc ctgggtgaca gagcaagact ccatctcaaa aaaaaaaaag 51060  
 aaaaaaatgt atattcttaa attacaaacg aggtctcact atgttgtcca ggctggctctc 51120  
 aagcagtcct cccacctcag ctgtaccaag cccaccaact gccctctctt aaaagtagtc 51180  
 attatcctgt ttccaaagat taattttact ttggctagaa ttttctaaaa actgaatcac 51240  
 gtagtatgta agcgggtatat acgcggttga gtgtctggct tcctttactc aacattattt 51300  
 ttgtgagagt tgttcatgat gccatgtata gttcattctc attgtataat tctgttttat 51360  
 aaatatccaa cttattcagc catcctactg ttgatggaca tttgggtagt gtccagtttg 51420  
 gggctaagtc caataacgct gctatgctca acatatggca ctctactgga cagttacctc 51480  
 agagtggaat tgctgagtc taaggcagac atatgttcgg ttttaggaga tactaacaaa 51540  
 cgggtgctgaa aaatgggtgt tcacatttgc actctcccca gcagttctgg ttgctgggca 51600  
 tcttcaattt cctagggctg aattaccaca aactaagtgg cttaaaacaa cagaaatggc 51660  
 caggcatggt ggctcatgcc tgtaatctca gcactttggg aggctgaggc aggcagatca 51720  
 cctgaggtca ggagtttgag accagcctga ccaacatgga gaaaccccat ctctactaaa 51780  
 aatacaaaaa aatagccagg catggtggca ggcgactgta gtcccagcca ctcaggaggc 51840  
 tgaggcagga gaatggcatg aacctgggag gcagagcttg cagttagcca agattgcacc 51900  
 actttgagag gccaaagggtg gcggatcacc tgaggtcagg agttcaggaa cagcctggcc 51960  
 aacatggcga aacctgtct ctattaaaaa taaaaaatt agccaggcgt ggtgggtgcac 52020  
 acctgtaatc ccagctactc aggacgctga ggtgggagaa ttgcttgaac ctgggaggca 52080  
 gaggctgagg ttagctgaga tcacacctct gcactctagc ctgggcaaca gaacaagctc 52140  
 catctaaaaa aaaaaaaaaa aaaaagtctt ttctgctcct tttgaaatat cagttagttt 52200  
 tcttcttttt ttctgttagt tgaattgtac tgattgattt tcaaatatta agccaaactt 52260



gcattcctga agtaaaactca atttgaatgt gttgtactat tctttgtatt tattgctgaa 52320  
ttccattcac taatatttag gatttttaca tctcttcttg agaaagactg accaaagtgt 52380  
ttccattctt gtaatgttct tgttggatgt gtgtatgaag tgaactacag tcatgcatca 52440  
cttaatgatg gggatatgtt ctgaaaaacg catcagtagc tgattctgtg gttgtctgaa 52500  
tatcatggac tctattttaca caaacctaaa tagaatagcc tattatactt aggttatatg 52560  
gtgtagtcta ttgctcctag gctgcaaacc tgtacagcat gttactgtac tgaatacggg 52620  
aagcaactgt aacagaatgg taagatttgt atatctaaat atagaaaagg tacagtga 52680  
atatgatata aaagcttaaa aatggtacat ctgcataggg cacttaccat gaatggagct 52740  
tgtaggactg aaagtgtctc tgggtgagtc agtgagtagt gagtgaatgt gaaagcctag 52800  
gttggtaccg tgcactacag tagacttcat aaacactgta cacttaggct acactaaatt 52860  
tacttcaaaa tattttatctt tcttcaataa taaattaatc ttagcttact gtgactgttt 52920  
tactttataa atttttaaat tttttaaact ctgtacagtg gtataaaaaat attttctttc 52980  
tcaccgggag tgggtggctca tgcctgtaat cccatcactt tgggaggccg aggcaggcgg 53040  
atcacagat caggagattg agaccatcct ggccaacatg gtgaaacccc atctctacta 53100  
aaaatacaaa aaaatagcca ggcattggtg caggcgccctg tagtcccagc tactcgggag 53160  
gctgaggcag gagaatggcg tgaaccaggg aggcagagct tgcagtgagc caagatcaca 53220  
ccactgcact ccagcctggg cgacagagca agactccatc tcaaaaaaaaa aaaaattctt 53280  
tctctatata cttattctat atactttttt ctatttttta cattttttat ttttattttt 53340  
acttttttaa tatttttgtt aaaaactaag tcatggcccg gtgcagtggc tcacgcttgt 53400  
aatctcagta ctttgggagg ctgaggtggg tggatcactc gaggtcagga gttcaacacc 53460  
agcctggcca atatggtgaa actctgtctc tactaaaaat ataaaaatta gccgggtgtg 53520  
gtggtgcacg cctgtagtcc cagctactca ggaggctgag gcagaagaat cgcttgaacc 53580  
caggaggcgg aggttgcagt aagccaagat catgccactg cactccagcc tgggagacag 53640  
agcaagactc catctcaaaa aaaaaaaaaa acaacaaaca cacacacaca cacacacaca 53700  
aaaaccattt ccaatagtgc taagtcctat gaagaatgtg gaataccaca gtgtgataag 53760  
ggaatcatgt gggagaaaag ctgctagata ggggtggtcag gacaagaggt gacatctcaa 53820  
cagaggcctg gcttttctga atttcatttt ccaaaatctg taaaataggc cagggtcagt 53880  
ggctcatgcc tgtaatccca gcactttgag aggctgaggc aggtggatca cctgagggtca 53940  
ggagttccaa acaagcctgg ccaacatgtt gaaatcctgt ttctactaga aatacgaaag 54000  
aattagctgg gcatggtggc atgcacctgt aataccagct acttaggagg ttaaggcatg 54060  
aaaattgtct gaacctggga ggtgaaagtt gcagtgagcc aagatcacac cactgtgtctc 54120  
cagcctgtgc gacagagtga gacctctct caaaaaaaaa aaaaaaatct gtaaaataaa 54180  
gacaaggata cattatctca caagcgtctt caaaggcctg aatgaggcaa tgcttacaga 54240  
acacatgcat ggtcctgata tctacaccta ataaatgacg gctactataa atcatgtaat 54300  
attaaacgta actttataag ttaataaaat taaagaacct agaacctgag ggcttgtcat 54360  
ggctaataagg tttggaaact tgctttttca gagacctgcg atggacttca cagatttgtc 54420  
caggctgccc ctggccctcc atgacacacc cccgattcct ggacaaccag aggagataca 54480  
gctgcttaga aaggaggcga ctcttagatc cagggatagc cccgtctggg gccagtgtgg 54540

aagctgcctc ccatctcaac tccctgagag ccacaggtgc ctggaggagc tgtgctgccg 54600  
gaaaaagccg ggggcctgca tcaccacctc agagctgttc aggaagctgg tcctgtccag 54660  
acacgtcctg cagttcctcc tgetctacca ggagcccttg ctggcgctgg atgtggattc 54720  
caccaacagc cggctgcggc actgtgccta caggtgctac gccacctggc gcttcggctc 54780  
ccaggacatg gctgactttg ccaacctgcc cagctgctgc cgctggagga tccggaaga 54840  
gtttccgaag agtgaagggc agtacagtgg cttcaagagt ccttactgaa gccaggcacc 54900  
gtggctcacg tctgtaatcc cagcgctttg ggaggccgag gcaggcagat cacctgagat 54960  
cgggagttgg agaccgcct ggctaacaag gcgaaatcct gtctgtacta aaaatacaaa 55020  
aatcagccag acatgggtggc atgcacctgc aatcccagct actcgggagg ctgaggcaca 55080  
agaatcactt gaaccggga ggcagaggtt gtagtgagcc cagattgtgc cactgctctc 55140  
cagcctggga ggcacagcaa actgtcccca aaaaaaaaaa aagagtcctt accaatagca 55200  
ggggctgcag tagccatgtt aacatgacat ttaccagcaa cttgaacttc acctgcaaag 55260  
ctctgtggcc acattttcag ccaaaggga atatgctttc atcttctgtt gctctctgtg 55320  
tctgagagca aagtgcctg gttaaacaaa ccagaatccc tctacatgga ctcagagaaa 55380  
agagattgag atgtaagtct caactctgtc ccaggaagt tgtgtgacct taggcctctc 55440  
acctctgtgc ctctgtctcc ttgttgccca actactatct cagagatatt gtgaggacaa 55500  
attgagacag tgcacatgaa ctgtctttta atgtgtaaag atctacatga atgcaaaaca 55560  
tttcattatg aggtcagact aggataatgt ccaactaaaa acaaacctt ttcctcctgg 55620  
ctggagaatg tggagaacta aaggtggcca caaattcttt gacactcaag tcccccaaga 55680  
cctaagggtt ttatctctc cccttgaata tgggtggctc tgattgcttt atccaaaagt 55740  
ggaagtgaca ttgtgtcagt ttcagatcct gatcttaaga ggctgacagc ttctacttgc 55800  
tgtcccttgg aactcttgct atcggggaag ccagacgcca tttaaaagtc tgcctatcct 55860  
ggccaggtgt ggtggctcac acctgtaatc ccagcacttt gggagaccaa ggcggcgga 55920  
tcacttaaag tcaggagtcc aagaccagac tcgccaacat ggtgaaaccg tatctcta 55980  
aaaaatacaa aaattagctg ggcattggtgc gggcacctgt agtcctagct atcaagaggc 56040  
tgagacagga gaaacacttg aacctgggag gtggaggttg cattgagctg agatcgtgcc 56100  
actgcactcc aggctgggtg acagagcgag actccatctc aaaaaaaaaa aaaaaagaaa 56160  
aaaaaaaaatgt ctgcctatcc tgagactgcc ctgctgtgag gaagccaag cagtcacgtg 56220  
gacagtgcct gaccagcccc agctttcaag ccatccaagc ccagtcacca aacatgagag 56280  
agaagaagcc ttcaggtgat tctggactcc actaacatat gactgatacc gcatgataca 56340  
tcccagtgga gaactgcccc ataatccag aaaaccaat tgctatctta agtccctaag 56400  
tttggggctt atttgttcca cagcaacagg taactggaac agagggaag cctgatgaat 56460  
gggcacacag actcagccca taccttcctt ggttctaatag ttctcaggga gcccgacca 56520  
acctgggag cctcaggaac ttaggtttcc actggacagt tctagaaggg ctatagacca 56580

<210> 2  
<211> 2164  
<212> DNA  
<213> Homo sapiens  
<220>

<221> CDS  
 <222> (79)..(1863)  
 <223>

<400> 2  
 ggcacgagggg cttgctgtgg ccctgtcagg aagagtagag ctctgggtcca gctccgcgca 60  
 gggaggggagg ctgtcacc atg ccg gcc tgc tgc agc tgc agt gat gtt ttc 111  
                           Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe  
                           1                          5                          10  
 cag tat gag acg aac aaa gtc act cgg atc cag agc atg aat tat ggc 159  
 Gln Tyr Glu Thr Asn Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly  
                           15                          20                          25  
 acc att aag tgg ttc ttc cac gtg atc atc ttt tcc tac gtt tgc ttt 207  
 Thr Ile Lys Trp Phe Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe  
                           30                          35                          40  
 gct ctg gtg agt gac aag ctg tac cag cgg aaa gag cct gtc atc agt 255  
 Ala Leu Val Ser Asp Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser  
                           45                          50                          55  
 tct gtg cac acc aag gtg aag ggg ata gca gag gtg aaa gag gag atc 303  
 Ser Val His Thr Lys Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile  
                           60                          65                          70                          75  
 gtg gag aat gga gtg aag aag ttg gtg cac agt gtc ttt gac acc gca 351  
 Val Glu Asn Gly Val Lys Lys Leu Val His Ser Val Phe Asp Thr Ala  
                           80                          85                          90  
 gac tac acc ttc cct ttg cag ggg aac tct ttc ttc gtg atg aca aac 399  
 Asp Tyr Thr Phe Pro Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn  
                           95                          100                          105  
 ttt ctc aaa aca gaa ggc caa gag cag cgg ttg tgt ccc gag tat ccc 447  
 Phe Leu Lys Thr Glu Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro  
                           110                          115                          120  
 acc cgc agg acg ctc tgt tcc tct gac cga ggt tgt aaa aag gga tgg 495  
 Thr Arg Arg Thr Leu Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp  
                           125                          130                          135  
 atg gac ccg cag agc aaa gga att cag acc gga agg tgt gta gtg cat 543  
 Met Asp Pro Gln Ser Lys Gly Ile Gln Thr Gly Arg Cys Val Val His  
                           140                          145                          150                          155  
 gaa ggg aac cag aag acc tgt gaa gtc tct gcc tgg tgc ccc atc gag 591  
 Glu Gly Asn Gln Lys Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu  
                           160                          165                          170  
 gca gtg gaa gag gcc ccc cgg cct gct ctc ttg aac agt gcc gaa aac 639  
 Ala Val Glu Glu Ala Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn  
                           175                          180                          185  
 ttc act gtg ctc atc aag aac aat atc gac ttc ccc ggc cac aac tac 687  
 Phe Thr Val Leu Ile Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr  
                           190                          195                          200  
 acc acg aga aac atc ctg cca ggt tta aac atc act tgt acc ttc cac 735  
 Thr Thr Arg Asn Ile Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His  
                           205                          210                          215  
 aag act cag aat cca cag tgt ccc att ttc cga cta gga gac atc ttc 783  
 Lys Thr Gln Asn Pro Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe  
                           220                          225                          230                          235  
 cga gaa aca ggc gat aat ttt tca gat gtg gca att cag ggc gga ata 831  
 Arg Glu Thr Gly Asp Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile  
                           240                          245                          250  
 atg ggc att gag atc tac tgg gac tgc aac cta gac cgt tgg ttc cat 879  
 Met Gly Ile Glu Ile Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His  
                           255                          260                          265  
 cac tgc cat ccc aaa tac agt ttc cgt cgc ctt gac gac aag acc acc 927

His Cys His Pro Lys Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr	
270 275 280	
aac gtg tcc ttg tac cct ggc tac aac ttc aga tac gcc aag tac tac	975
Asn Val Ser Leu Tyr Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr	
285 290 295	
aag gaa aac aat gtt gag aaa cgg act ctg ata aaa gtc ttc ggg atc	1023
Lys Glu Asn Asn Val Gly Lys Arg Thr Leu Ile Lys Val Phe Gly Ile	
300 305 310 315	
cgt ttt gac atc ctg gtt ttt ggc acc gga gga aaa ttt gac att atc	1071
Arg Phe Asp Ile Leu Val Phe Gly Thr Gly Lys Phe Asp Ile Ile	
320 325 330	
cag ctg gtt gtg tac atc ggc tca acc ctc tcc tac ttc ggt ctg gcc	1119
Gln Leu Val Val Tyr Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala	
335 340 345	
gct gtg ttc atc gac ttc ctc atc gac act tac tcc agt aac tgc tgt	1167
Ala Val Phe Ile Asp Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys	
350 355 360	
cgc tcc cat att tat ccc tgg tgc aag tgc tgt cag ccc tgt gtg gtc	1215
Arg Ser His Ile Tyr Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val	
365 370 375	
aac gaa tac tac tac agg aag aag tgc gag tcc att gtg gag cca aag	1263
Asn Glu Tyr Tyr Tyr Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys	
380 385 390 395	
ccg aca tta aag tat gtg tcc ttt gtg gat gaa tcc cac att agg atg	1311
Pro Thr Leu Lys Tyr Val Ser Phe Val Asp Glu Ser His Ile Arg Met	
400 405 410	
gtg aac cag cag cta cta ggg aga agt ctg caa gat gtc aag ggc caa	1359
Val Asn Gln Gln Leu Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln	
415 420 425	
gaa gtc cca aga cct gcg atg gac ttc aca gat ttg tcc agg ctg ccc	1407
Glu Val Pro Arg Pro Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro	
430 435 440	
ctg gcc ctc cat gac aca ccc ccg att cct gga caa cca gag gag ata	1455
Leu Ala Leu His Asp Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile	
445 450 455	
cag ctg ctt aga aag gag gcg act cct aga tcc agg gat agc ccc gtc	1503
Gln Leu Leu Arg Lys Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val	
460 465 470 475	
tgg tgc cag tgt gga agc tgc ctc cca tct caa ctc cct gag agc cac	1551
Trp Cys Gln Cys Gly Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His	
480 485 490	
agg tgc ctg gag gag ctg tgc tgc cgg aaa aag ccg ggg gcc tgc atc	1599
Arg Cys Leu Glu Glu Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile	
495 500 505	
acc acc tca gag ctg ttc agg aag ctg gtc ctg tcc aga cac gtc ctg	1647
Thr Thr Ser Glu Leu Phe Arg Lys Leu Val Leu Ser Arg His Val Leu	
510 515 520	
cag ttc ctc ctg ctc tac cag gag ccc ttg ctg gcg ctg gat gtg gat	1695
Gln Phe Leu Leu Leu Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp	
525 530 535	
tcc acc aac agc cgg ctg cgg cac tgt gcc tac agg tgc tac gcc acc	1743
Ser Thr Asn Ser Arg Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr	
540 545 550 555	
tgg cgc ttc gcc tcc cag gac atg gct gac ttt gcc atc ctg ccc agc	1791
Trp Arg Phe Gly Ser Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser	
560 565 570	
tgc tgc cgc tgg agg atc cgg aaa gag ttt ccg aag agt gaa ggg cag	1839

Cys Cys Arg Trp Arg Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln  
 575 580 585  
 tac agt ggc ttc aag agt cct tac tgaagccagg caccgtggct cacgtctgta 1893  
 Tyr Ser Gly Phe Lys Ser Pro Tyr  
 590 595  
 atcccagcgc tttgggagggc cgaggcaggc agatcacctg aggtcgggag ttggagaccc 1953  
 gcctggctaa caaggcgaaa tcctgtctgt actaaaaata caaaaatcag ccagacatgg 2013  
 tggcatgcac ctgcaatccc agctactcgg gaggtctgagg cacaagaatc acttgaaccc 2073  
 gggaggcaga ggttgtagtg agcccagatt gtgccactgc tctccagcct gggaggcaca 2133  
 gcaaactgtc ccaaaaaaaaa a 2164

<210> 3  
 <211> 595  
 <212> PRT  
 <213> Homo sapiens

<400> 3

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15  
 Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30  
 Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45  
 Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60  
 Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80  
 Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95  
 Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110  
 Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125  
 Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140  
 Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160  
 Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175  
 Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190  
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys His Pro Lys  
 260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
 290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
 305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
 325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
 340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
 355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
 370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
 385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
 405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
 420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
 435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
 450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
 465 470 475 480

---

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
 485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
580 585 590

Ser Pro Tyr  
595

<210> 4  
<211> 595  
<212> PRT  
<213> Homo sapiens

<220>  
<221> DOMAIN  
<222> (1)..(20)  
<223> Intracellular domain

<220>  
<221> TRANSMEM  
<222> (21)..(46)  
<223>

<220>  
<221> DOMAIN  
<222> (47)..(320)  
<223> Extracellular domain

<220>  
<221> TRANSMEM  
<222> (321)..(356)  
<223>

<220>  
<221> DOMAIN  
<222> (357)..(595)  
<223> Intracellular domain

<400> 4

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80  
 Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95  
 Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110  
 Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125  
 Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140  
 Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160  
 Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175  
 Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190  
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205  
 Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220  
 Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240  
 Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255  
 Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
 260 265 270  
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285  
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
 290 295 300  
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
 305 310 315 320  
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
 325 330 335  
 Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
 340 345 350  
 Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
 355 360 365



Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
580 585 590

Ser Pro Tyr  
595

<210> 5  
<211> 595  
<212> PRT  
<213> Homo sapiens

<400> 5

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp

35                      40                      45  
 Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
     50                      55                      60  
 Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
     65                      70                      75                      80  
 Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
                     85                      90                      95  
 Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
                     100                      105                      110  
 Gly Gln Glu Gln Trp Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
                     115                      120                      125  
 Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
     130                      135                      140  
 Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
     145                      150                      155                      160  
 Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
                     165                      170                      175  
 Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
                     180                      185                      190  
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
                     195                      200                      205  
 Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
     210                      215                      220  
 Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
     225                      230                      235                      240  
 Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
                     245                      250                      255  
 Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
                     260                      265                      270  
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
                     275                      280                      285  
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
     290                      295                      300  
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
     305                      310                      315                      320  
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
                     325                      330                      335  
 Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp



Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30  
 Phe His-Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45  
 Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60  
 Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80  
 Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95  
 Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110  
 Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125  
 Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140  
 Lys Gly Ile Gln Thr Arg Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160  
 Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175  
 Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190  
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205  
 Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220  
 Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240  
 Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255  
 Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
 260 265 270  
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285  
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
 290 295 300  
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
 305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
580 585 590

Ser Pro Tyr  
595

<210> 7  
<211> 595

<212> PRT  
 <213> Homo sapiens

<400> 7

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Lys Asn Phe Thr Val Leu Ile  
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
 260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
580 585 590

Ser Pro Tyr  
595

<210> 8  
 <211> 595  
 <212> PRT  
 <213> Homo sapiens

<400> 8

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Pro Ile  
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255



Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
 580 585 590

Ser Pro Tyr  
 595

<210> 9  
 <211> 595  
 <212> PRT  
 <213> Homo sapiens

<400> 9

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp

225	230	235	240
Asn Phe Ser Asp Val 245	Ala Ile Gln Gly	Gly Ile Met Gly Ile 250	Glu Ile 255
Tyr Trp Asp Cys Asn 260	Leu Asp Arg Trp	Phe His His Cys Cys 265	Pro Lys 270
Tyr Ser Phe Arg Arg 275	Leu Asp Asp Lys	Thr Thr Asn Val 280	Ser Leu Tyr 285
Pro Gly Tyr Asn Phe Arg 290	Tyr Ala Lys Tyr	Tyr Tyr Lys Glu 300	Asn Asn Val
Glu Lys Arg Thr Leu 305	Ile Lys Val Phe Gly	Ile Arg Phe Asp Ile 310	Leu 320
Val Phe Gly Thr Gly 325	Gly Lys Phe Asp Ile	Ile Ile Gln Leu Val 330	Val Tyr 335
Ile Gly Ser Thr Leu 340	Ser Tyr Phe Gly	Leu Ala Ala Val Phe 345	Ile Asp 350
Phe Leu Ile Asp Thr Tyr 355	Ser Ser Asn Cys Cys Arg	Ser His Ile Tyr 360	
Pro Trp Cys Lys Cys Cys 370	Gln Pro Cys Val Val	Asn Glu Tyr Tyr Tyr 375	
Arg Lys Lys Cys Glu Ser 385	Ile Val Glu Pro Lys	Pro Thr Leu Lys Tyr 390	
Val Ser Phe Val Asp 405	Glu Ser His Ile Arg	Met Val Asn Gln Gln 410	Leu 415
Leu Gly Arg Ser Leu Gln 420	Asp Val Lys Gly	Gln Glu Val Pro Arg 425	Pro
Ala Met Asp Phe Thr Asp 435	Leu Ser Arg Leu Pro	Leu Ala Leu His Asp 440	
Thr Pro Pro Ile Pro Gly 450	Gln Pro Glu Glu Ile	Gln Leu Leu Arg Lys 455	
Glu Ala Thr Pro Arg Ser 465	Arg Asp Ser Pro Val	Trp Cys Gln Cys Gly 470	
Ser Cys Leu Pro Ser Gln 485	Leu Pro Glu Ser His	Arg Cys Leu Glu Glu 490	
Leu Cys Cys Arg Lys Lys 500	Pro Gly Ala Cys Ile	Thr Thr Ser Glu Leu 505	
Phe Arg Lys Leu Val Leu 515	Ser Arg His Val Leu	Gln Phe Leu Leu Leu 520	
Tyr Gln Glu Pro Leu Leu 525	Ala Leu Asp Val Asp	Ser Thr Asn Ser Arg 530	

530

535

540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
 580 585 590

Ser Pro Tyr  
 595

<210> 10  
 <211> 595  
 <212> PRT  
 <213> Homo sapiens

<400> 10

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220  
 Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240  
 Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255  
 Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
 260 265 270  
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285  
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
 290 295 300  
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
 305 310 315 320  
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
 325 330 335  
 Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
 340 345 350  
 Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
 355 360 365  
 Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
 370 375 380  
 Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
 385 390 395 400  
 Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
 405 410 415  
 Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
 420 425 430  
 Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
 435 440 445  
 Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
 450 455 460  
 Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
 465 470 475 480  
 Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
 485 490 495  
 Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Asn Leu Pro Ser Cys Cys Arg Trp Arg  
565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
580 585 590

Ser Pro Tyr  
595

<210> 11  
<211> 588  
<212> PRT  
<213> Homo sapiens

<400> 11

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190  
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205  
 Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220  
 Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240  
 Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255  
 Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
 260 265 270  
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285  
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
 290 295 300  
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
 305 310 315 320  
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
 325 330 335  
 Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
 340 345 350  
 Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
 355 360 365  
 Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
 370 375 380  
 Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
 385 390 395 400  
 Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
 405 410 415  
 Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
 420 425 430  
 Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
 435 440 445  
 Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
 450 455 460  
 Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
 465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Glu Glu Leu Cys Cys Arg Lys Lys Pro  
 485 490 495

Gly Ala Cys Ile Thr Thr Ser Glu Leu Phe Arg Lys Leu Val Leu Ser  
 500 505 510

Arg His Val Leu Gln Phe Leu Leu Leu Tyr Gln Glu Pro Leu Leu Ala  
 515 520 525

Leu Asp Val Asp Ser Thr Asn Ser Arg Leu Arg His Cys Ala Tyr Arg  
 530 535 540

Cys Tyr Ala Thr Trp Arg Phe Gly Ser Gln Asp Met Ala Asp Phe Ala  
 545 550 555 560

Ile Leu Pro Ser Cys Cys Arg Trp Arg Ile Arg Lys Glu Phe Pro Lys  
 565 570 575

Ser Glu Gly Gln Tyr Ser Gly Phe Lys Ser Pro Tyr  
 580 585

<210> 12  
 <211> 595  
 <212> PRT  
 <213> Homo sapiens

<400> 12

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn  
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe  
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp  
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys  
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val  
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro  
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu  
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu  
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser  
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys  
 145 150 155 160



Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala  
 165 170 175  
 Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile  
 180 185 190  
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile  
 195 200 205  
 Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro  
 210 215 220  
 Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp  
 225 230 235 240  
 Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile  
 245 250 255  
 Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys  
 260 265 270  
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr  
 275 280 285  
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val  
 290 295 300  
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu  
 305 310 315 320  
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr  
 325 330 335  
 Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp  
 340 345 350  
 Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr  
 355 360 365  
 Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr  
 370 375 380  
 Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr  
 385 390 395 400  
 Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu  
 405 410 415  
 Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro  
 420 425 430  
 Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp  
 435 440 445  
 Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys  
 450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly  
 465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu  
 485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu  
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu  
 515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg  
 530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser  
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg  
 565 570 575

Ile Gln Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys  
 580 585 590

Ser Pro Tyr  
 595

<210> 13  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 13  
 gtaacctcac tgagaacaga

20

<210> 14  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 14  
 ttggttaggc gaaggcgggc

20

<210> 15  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens

<400> 15  
 ccttgtaggc gctttcgtgc t

21

<210> 16  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens

<400> 16  
 ttgtgtatcc gtccatcagc t

21

<210> 17  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 17 actgccatat gtactattcc	20
<210> 18 <211> 20 <212> DNA <213> Homo sapiens	
<400> 18 aaatctaccc agccacttag	20
<210> 19 <211> 20 <212> DNA <213> Homo sapiens	
<400> 19 tgcagtggcc cacagcacag	20
<210> 20 <211> 20 <212> DNA <213> Homo sapiens	
<400> 20 catcctccaa tgcttgcattc	20
<210> 21 <211> 20 <212> DNA <213> Homo sapiens	
<400> 21 tttcccatgt ttgccattta	20
<210> 22 <211> 20 <212> DNA <213> Homo sapiens	
<400> 22 caagagcagt ggttgtgtcc	20
<210> 23 <211> 20 <212> DNA <213> Homo sapiens	
<400> 23 gagcccacag atatctactg	20
<210> 24 <211> 20 <212> DNA <213> Homo sapiens	
<400> 24 tgcttgctc agtctctggt	20
<210> 25 <211> 20 <212> DNA <213> Homo sapiens	
<400> 25 tctctcgccc aggttgagtt	20

<210> 26  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 26  
 acgtcctctc cgcagttctt

20

<210> 27  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 27  
 attcagaccg aaaggtgtgt

20

<210> 28  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 28  
 tgcctgggtgt cccatcgagg

20

<210> 29  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 29  
 gttgtaacac tcctgtacca

20

<210> 30  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 30  
 aacagtgccca aaaacttcac

20

<210> 31  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 31  
 tcactgtgcc catcaagaac

20

<210> 32  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 32  
 tgctgcgtgc ttctgggtca

20

<210> 33  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 33  
 ctggagaacc cttttccaag

20

<210> 34  
 <211> 20  
 <212> DNA

<213> Homo sapiens  
 <400> 34  
 caggaggcag ggatcatttg 20

<210> 35  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
 <400> 35  
 tccacccgct acgctaaggt 20

<210> 36  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
 <400> 36  
 tgcgctaagg actttaccta 20

<210> 37  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
 <400> 37  
 tttacctacc ctacctcgtc 20

<210> 38  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
 <400> 38  
 ccatcactgc tgtcccaaatt 20

<210> 39  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
 <400> 39  
 tcgccttgat gacaagacca 20

<210> 40  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
 <400> 40  
 gtaaactctt ccactctggt 20

<210> 41  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
 <400> 41  
 accaagacac ggagagattc 20

<210> 42  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
 <400> 42  
 taatattaaa tgtaacttta 20

<210> 43  
<211> 20  
<212> DNA  
<213> Homo sapiens  
  
<400> 43  
aatattaaac ataactttat  
20  
  
<210> 44  
<211> 20  
<212> DNA  
<213> Homo sapiens  
  
<400> 44  
ccatctcaac tggaggagct  
20  
  
<210> 45  
<211> 20  
<212> DNA  
<213> Homo sapiens  
  
<400> 45  
gactttgccca tcctgcccag  
20  
  
<210> 46  
<211> 20  
<212> DNA  
<213> Homo sapiens  
  
<400> 46  
tggaggatcc agaaagagtt  
20  
  
<210> 47  
<211> 20  
<212> DNA  
<213> Homo sapiens  
  
<400> 47  
aatcccagca ctttgggagg  
20  
  
<210> 48  
<211> 20  
<212> DNA  
<213> Homo sapiens  
  
<400> 48  
aaactgtccc aaaaaaaaaa  
20  
  
<210> 49  
<211> 20  
<212> DNA  
<213> Homo sapiens  
  
<400> 49  
aactgtccc caaaaaaaaa  
20  
  
<210> 50  
<211> 20  
<212> DNA  
<213> Homo sapiens  
  
<400> 50  
ctgtcccccac aaaaaaaaaa  
20  
  
<210> 51  
<211> 20

<212> DNA	
<213> Homo sapiens	
<400> 51	
caaggcgggt ggatcactta	20
<210> 52	
<211> 19	
<212> DNA	
<213> Homo sapiens	
<400> 52	
cgtaggactt ggcgcttct	19
<210> 53	
<211> 21	
<212> DNA	
<213> Homo sapiens	
<400> 53	
gagcacgtct cagattcgaa a	21
<210> 54	
<211> 23	
<212> DNA	
<213> Homo sapiens	
<400> 54	
ccatgaggca ggtatgacta ttc	23
<210> 55	
<211> 20	
<212> DNA	
<213> Homo sapiens	
<400> 55	
ctcctggatc tcacccagtt	20
<210> 56	
<211> 23	
<212> DNA	
<213> Homo sapiens	
<400> 56	
ctcgtccagc tttgatatta agc	23
<210> 57	
<211> 22	
<212> DNA	
<213> Homo sapiens	
<400> 57	
ggtccttagt gctagaacca ga	22
<210> 58	
<211> 18	
<212> DNA	
<213> Homo sapiens	
<400> 58	
attcatccgt cagtggcc	18
<210> 59	
<211> 22	
<212> DNA	
<213> Homo sapiens	
<400> 59	

gccatgtgaa ttttctaccg at	22
<210> 60	
<211> 20	
<212> DNA	
<213> Homo sapiens	
<400> 60	
ttcgttgtgg ttaggatggg	20
<210> 61	
<211> 22	
<212> DNA	
<213> Homo sapiens	
<400> 61	
caaggatgct cagggtagta gc	22
<210> 62	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 62	
cactaggttt gctgtatcca tttct	25
<210> 63	
<211> 20	
<212> DNA	
<213> Homo sapiens	
<400> 63	
gcaactgtgt gagagcttgg	20
<210> 64	
<211> 19	
<212> DNA	
<213> Homo sapiens	
<400> 64	
tcaaccctgg tccagtgtg	19
<210> 65	
<211> 25	
<212> DNA	
<213> Homo sapiens	
<400> 65	
caccaagtag ctctcactca taagg	25
<210> 66	
<211> 24	
<212> DNA	
<213> Homo sapiens	
<400> 66	
caataacact tgtgcgagtt aggt	24
<210> 67	
<211> 21	
<212> DNA	
<213> Homo sapiens	
<400> 67	
catcttgttg ccttggaac c	21
<210> 68	



<211> 23  
 <212> DNA  
 <213> Homo sapiens

<400> 68  
 gtgagtggta atcctgctac tgc

23

<210> 69  
 <211> 19  
 <212> DNA  
 <213> Homo sapiens

<400> 69  
 aggccactc ctgtactcg

19

<210> 70  
 <211> 19  
 <212> DNA  
 <213> Homo sapiens

<400> 70  
 ccaagtcaca gcatgaggc

19

<210> 71  
 <211> 18  
 <212> DNA  
 <213> Homo sapiens

<400> 71  
 acccagcgac gtatccac

18

<210> 72  
 <211> 19  
 <212> DNA  
 <213> Homo sapiens

<400> 72  
 aagcatgggg ttccatttc

19

<210> 73  
 <211> 24  
 <212> DNA  
 <213> Homo sapiens

<400> 73  
 gcataaaagg gactcctgct agta

24

<210> 74  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens

<400> 74  
 gcttacagaa cacatgcatg g

21

<210> 75  
 <211> 19  
 <212> DNA  
 <213> Homo sapiens

<400> 75  
 gcacctgtag gcacagtgc

19

<210> 76  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens

<400> 76	
atcaccacct cagagctgtt c	21
<210> 77	
<211> 21	
<212> DNA	
<213> Homo sapiens	
<400> 77	
gttaacatgg ctactgcagc c	21
<210> 78	
<211> 21	
<212> DNA	
<213> Homo sapiens	
<400> 78	
gcttagaaag gaggcgactc c	21
<210> 79	
<211> 22	
<212> DNA	
<213> Homo sapiens	
<400> 79	
ttgtgacatt tgcaaggctg cc	22
<210> 80	
<211> 20	
<212> DNA	
<213> Homo sapiens	
<400> 80	
tctgaagctc tgctcctgag	20
<210> 81	
<211> 20	
<212> DNA	
<213> Homo sapiens	
<400> 81	
ctcaccttct ggcttccagt	20
<210> 82	
<211> 20	
<212> DNA	
<213> Homo sapiens	
<400> 82	
cttaccactc ccaggactaa	20
<210> 83	
<211> 20	
<212> DNA	
<213> Homo sapiens	
<400> 83	
gtctgcctgt tcactgcat	20
<210> 84	
<211> 21	
<212> DNA	
<213> Homo sapiens	
<400> 84	
cagagacctt cagaaacttc g	21

<210> 85  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 85  
 agatcaccag ggacacagtg 20  
  
 <210> 86  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 86  
 ctcaactcca ctttcctcgg 20  
  
 <210> 87  
 <211> 23  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 87  
 cctttcactt ttttggtctc atg 23  
  
 <210> 88  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 88  
 gggagaattc tgaaaatgcc c 21  
  
 <210> 89  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 89  
 ggaccagagc tctactcttc 20  
  
 <210> 90  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 90  
 aggtcataga tcgacctgcc 20  
  
 <210> 91  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 91  
 aagaagcgcc aagtcctacg 20  
  
 <210> 92  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens  
  
 <400> 92  
 gcaatccaga ctgaagttga c 21  
  
 <210> 93  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 93 actctgggtct gcagttggtg	20
<210> 94 <211> 24 <212> DNA <213> Homo sapiens	
<400> 94 cctttaaaat cagagacctt caga	24
<210> 95 <211> 20 <212> DNA <213> Homo sapiens	
<400> 95 gcccatcctc tgaacaccat	20
<210> 96 <211> 21 <212> DNA <213> Homo sapiens	
<400> 96 cccttggaac tcttgctatc g	21
<210> 97 <211> 21 <212> DNA <213> Homo sapiens	
<400> 97 ggcagtacag tggcttcaag a	21
<210> 98 <211> 21 <212> DNA <213> Homo sapiens	
<400> 98 gtgggacagt ttgctgtgcc t	21
<210> 99 <211> 20 <212> DNA <213> Homo sapiens	
<400> 99 gagtccttac caatagcagg	20
<210> 100 <211> 21 <212> DNA <213> Homo sapiens	
<400> 100 gtcaaagaat ttgtggccac c	21
<210> 101 <211> 26 <212> DNA <213> Homo sapiens	
<400> 101 catgaactgt cttttaatgt gtaaag	26

<210> 102  
 <211> 22  
 <212> DNA  
 <213> Homo sapiens

<400> 102  
 gagatacggg ttcacccatgt tg 22

<210> 103  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 103  
 aattagctgg gcatgggtgcg 20

<210> 104  
 <211> 21  
 <212> DNA  
 <213> Homo sapiens

<400> 104  
 ttgagatgga gtctcgctct g 21

<210> 105  
 <211> 20  
 <212> DNA  
 <213> Homo sapiens

<400> 105  
 cactgtccac gtgactgctt 20

<210> 106  
 <211> 25  
 <212> DNA  
 <213> Homo sapiens

<400> 106  
 tcctacttcg gtctggtaag agatt 25

<210> 107  
 <211> 19  
 <212> DNA  
 <213> Homo sapiens

<400> 107  
 gggcctaatt ttcgtgcat 19

<210> 108  
 <211> 23  
 <212> DNA  
 <213> Homo sapiens

<400> 108  
 aagaacctag aacctgaggg ctt 23

<210> 109  
 <211> 19  
 <212> DNA  
 <213> Homo sapiens

<400> 109  
 ttgagatggg aggcagctt 19

<210> 110  
 <211> 17  
 <212> DNA

62

<213> Homo sapiens

<400> 110  
ttcggctccc aggacat

17

<210> 111

<211> 20

<212> DNA

<213> Homo sapiens

<400> 111  
cacagagctt tgcaggtgaa

20